


1.

A ray of light is incident on the internal boundary of a rectangular glass block in air.

Part of the light refracts out of the block at an angle of 30° .

Some of the remaining light reflects within the block to become incident on the right-hand boundary. refractive index of glass = 1.48



What is the angle of incidence of the ray at the right-hand boundary?

- A 20°
- B 42°
- C 48°
- D 70°

(Total 1 mark)

2.

In a Young's double-slit experiment, monochromatic light is incident on two narrow slits and the resulting interference pattern is observed on a screen.

Which change **decreases** the fringe separation?

- A decreasing the separation between the two slits
- B increasing the distance between the slits and the screen
- C using monochromatic light of higher frequency
- D using monochromatic light of longer wavelength

(Total 1 mark)

3.

A diffraction grating is illuminated normally.

The second-order maximum for light of wavelength 650 nm occurs at the same angle as the third-order maximum for light of wavelength λ .

What is λ ?

A 217 nm

B 325 nm

C 433 nm

D 975 nm

(Total 1 mark)

4.

Light of wavelength λ is incident normally on two parallel slits of separation s . Fringes of spacing w are seen on a screen at a distance D from the slits.

Which row gives another arrangement that produces a fringe spacing of w ?

	Wavelength	Slit separation	Distance between slits and screen	
A	2λ	$2s$	$2D$	<input type="checkbox"/>
B	2λ	$4s$	$2D$	<input type="checkbox"/>
C	2λ	$2s$	$4D$	<input type="checkbox"/>
D	4λ	$2s$	$2D$	<input type="checkbox"/>

(Total 1 mark)

5.

A narrow beam of monochromatic light is incident normally to a diffraction grating. The first-order diffracted beam makes an angle of 20° with the normal to the grating.

What is the highest order visible with this grating at this wavelength?

A 2

B 3

C 4

D 5

(Total

6. The speed of light decreases by 40% when it travels from air into a transparent medium.

What is the refractive index of the medium?

- A 0.6
- B 1.4
- C 1.7
- D 2.5

(Total 1 mark)

7. A monochromatic light wave travels from glass into air.

Which row shows what happens to the wavelength, speed and photon energy?

	Wavelength	Speed	Photon energy	
A	increases	increases	increases	<input type="radio"/>
B	does not change	decreases	does not change	<input type="radio"/>
C	does not change	decreases	increases	<input type="radio"/>
D	increases	increases	does not change	<input type="radio"/>

(Total 1 mark)

8. Monochromatic light is incident normally on a diffraction grating that has 4.50×10^5 lines m^{-1} . The angle between the second-order diffraction maxima is 44° .

What is the wavelength of the light?

- A 208 nm
- B 416 nm
- C 772 nm
- D 832 nm

(Total 1 mark)

9. In a Young's double-slit experiment, the spacing of the double slits is s and the distance between the slits and the screen on which fringes are formed is D . When monochromatic light of wavelength λ is incident on the slits the distance between adjacent fringes on the screen is w .

Which row shows another arrangement that produces a fringe spacing of w ?

	Spacing of double slits	Distance between the slits and the screen	Wavelength of the light	
A	$4s$	$2D$	2λ	<input type="radio"/>
B	$2s$	$4D$	2λ	<input type="radio"/>
C	$2s$	$2D$	4λ	<input type="radio"/>
D	$2s$	$2D$	2λ	<input type="radio"/>

(Total 1 mark)

10. Monochromatic electromagnetic radiation of wavelength 5.8×10^{-7} m is incident normally on a diffraction grating with 3.0×10^5 lines per metre.

What is the highest order maximum produced?

- A 5
- B 6
- C 10
- D 13

(Total 1 mark)

11. Which characteristics of monochromatic light change when the light passes from air into glass?

- A Speed, wavelength and frequency.
- B Speed and frequency only.
- C Speed and wavelength only.
- D Wavelength and frequency only.

(Total 1 mark)

12.

Which is a description of the pattern produced when monochromatic light passes through a very narrow slit?

A A series of equally-spaced light and dark fringes.

B A narrow central maximum with wider side fringes.


C A few bright fringes that are widely spaced.

D A wide central maximum with narrower side fringes.

(Total 1 mark)

13.

A ray of light is incident on a glass-air boundary of a rectangular block as shown.



The refractive index of this glass is 1.5

The refractive index of air is 1.0

The angle of incidence of the light at the first glass-air boundary is 44°

What is the path of the ray of light?

A

B

C

D


(Total 1 mark)

14. Rays of light are incident at the same angle θ on the core-cladding boundary of optical fibres **P** and **Q**.

The cores of **P** and **Q** have the same refractive index n .

P and **Q** are the same length L .

The core diameter of **P** is half that of **Q**.



The time for the ray to travel along optical fibre **P** is

$$\frac{nL}{c \sin \theta}$$

where c is the speed of light in a vacuum.

What is the time for the ray to travel along optical fibre **Q**?

A $\frac{nL}{c \sin \theta}$

B $\frac{nL}{2c \sin \theta}$


C $\frac{2nL}{c \sin \theta}$

D $\frac{4nL}{c \sin \theta}$

(Total 1 mark)

15.

A diffraction grating is illuminated normally with light of wavelength 6.5×10^{-7} m. When a screen is 1.5 m from the grating, the distance between the zero and first-order maxima on the screen is 0.30 m.



What is the number of lines per mm of the diffraction grating?

A 3.3×10^{-6}

B 3.3×10^{-3}


C 3.0×10^2

D 3.0×10^5

(Total 1 mark)

16.

In the diagram, P is the source of a wave of frequency 50 Hz



The wave travels to R by two routes, $P \rightarrow Q \rightarrow R$ and $P \rightarrow R$. The speed of the wave is 30 m s^{-1}

What is the path difference between the two waves at **R** in terms of the wavelength λ of the waves?

- A** 4.8λ
- B** 8.0λ
- C** 13.3λ
- D** 20.0λ

(Total 1 mark)

17.

An electromagnetic wave enters a fibre-optic cable from air. On entering the cable, the wave slows down to three-fifths of its original speed.

What is the refractive index of the core of the fibre-optic cable?

- A** 0.67
- B** 1.33
- C** 1.50
- D** 1.67

(Total 1 mark)

18.


A diffraction grating has 500 lines per mm. When monochromatic light is incident normally on the grating the third-order spectral line is formed at an angle of 60° from the normal to the grating.

What is the wavelength of the monochromatic light?

- A** 220 nm
- B** 580 nm
- C** 960 nm
- D** 1700 nm

(Total 1 mark)

19. The diagram shows a ray of light travelling in air and incident on a glass block of refractive index 1.5



What is the angle of refraction in the glass?

- A 22.5°
- B 23.3°
- C 33.1°
- D 59.4°

(Total 1 mark)

20. When light of wavelength 5.0×10^{-7} m is incident normally on a diffraction grating the fourth-order maximum is observed at an angle of 30° .

What is the number of lines per mm on the diffraction grating?

- A 2.5×10^2
- B 2.5×10^5
- C 1.0×10^3
- D 1.0×10^6

(Total 1 mark)