

M1.C

[1]

M2.A

[1]

M3.C

[1]

M4.A

[1]

M5.(a) $t = \sqrt{\frac{2s}{g}}$ or $4.5 = \frac{1}{2} \times 9.81 \times t^2$ ✓

$t = 0.96 \text{ s}$ ✓

2

(b) Field strength = 186000 V m^{-1} ✓

Acceleration = Eq / m

or $186\,000 \times 1.2 \times 10^{-6}$ ✓

0.22 m s^{-2} ✓

3

(c) $0.10(3)\text{m}$ (allow ecf from (i)) ✓

1

(d) Force on a particle = mg and

acceleration = F / m so always = g ✓

Time to fall (given distance) depends (only) on the distance and acceleration
✓

OR:

$$g = GM / r^2 \quad \checkmark$$

$$\text{Time to fall} = \sqrt{2s / g}$$

so no m in equations to determine time to fall ✓

2

(e) Mass is not constant since particle mass will vary ✓

Charge on a particle is not constant ✓

$$\text{Acceleration} = Eq / m \text{ or } (V / d) (q / m) \text{ or } Vq / dm \quad \checkmark$$

E or V / d constant but charge and mass are 'random' variables so q / m will vary (or unlikely to be the same) ✓

4

[12]

M6.D

[1]

M7.B

[1]

M8.D

[1]

M9.D

[1]

M10.A

[1]

M11.C

[1]

M12.B

[1]

M13.C

[1]

M14.A

[1]

- M15.(a)** (i) force acts towards left or in opposite direction to field lines ✓
because ion (or electron) has negative charge
(∴ experiences force in opposite direction to field) ✓

Mark sequentially.

Essential to refer to negative charge (or force on + charge is to right) for 2nd mark.

(ii) (use of $W = F s$ gives) force $F = \frac{4.0 \times 10^{-16}}{63 \times 10^{-3}} \checkmark$

$$= 6.3(5) \times 10^{-15} \text{ (N)} \checkmark$$

If mass of ion m is used correctly **using algebra** with $F = ma$, allow both marks (since m will cancel). If numerical value for m is used, max 1.

2

(iii) electric field strength $E \left(= \frac{F}{Q} \right) = \frac{6.35 \times 10^{-15}}{3 \times 1.6 \times 10^{-19}} = 1.3(2) \checkmark 10^4 \text{ (N C}^{-1}\text{)} \checkmark$

[or $\Delta V \left(= \frac{\Delta W}{Q} \right) = \frac{4.0 \times 10^{-16}}{3 \times 1.60 \times 10^{-19}} \text{ (833 V)}$

$$E \left(= \frac{\Delta V}{d} \right) = \frac{833}{63 \times 10^{-3}} = 1.3(2) \checkmark 10^4 \text{ (V m}^{-1}\text{)} \checkmark]$$

Allow ECF from wrong F value in (ii).

1

- (b) (i) (vertically) downwards on diagram \checkmark
reference to Fleming's LH rule **or** equivalent statement \checkmark
Mark sequentially.
1st point: allow "into the page".

2

- (ii) number of free electrons in wire = $A \times l \times$ number density
 $= 5.1 \times 10^{-6} \times 95 \times 10^{-3} \times 8.4 \times 10^{28} = 4.1 \text{ (4.07)} \times 10^{22} \checkmark$
Provided it is shown correctly to at least 2SF, final answer alone is sufficient for the mark. (Otherwise working is mandatory).

1

(iii) $B \left(= \frac{F}{Qv} \right) = \frac{1.4 \times 10^{-25}}{1.60 \times 10^{-19} \times 5.5 \times 10^{-6}} \checkmark = 0.16 \text{ (0.159) (T)} \checkmark$

$$\text{[or } B \left(= \frac{F}{Il} \right) = \frac{1.4 \times 10^{-25} \times 4.07 \times 10^{22}}{0.38 \times 95 \times 10^{-3}} \checkmark = 0.16 \text{ (0.158) (T) } \checkmark \text{]}$$

In 2nd method allow ECF from wrong number value in (ii).

2
[10]

M16.B

[1]

M17.A

[1]

M18.D

[1]

M19.(a) (i) required pd ($= 2.5 \times 10^6 \times 12 \times 10^{-3}$) $= 3.0(0) \times 10^4$ (V) \checkmark

1

(ii) charge required Q ($= CV$) $= 3.7 \times 10^{-12} \times 3.00 \times 10^4 \checkmark$

($= 1.11 \times 10^{-7}$ C)

Allow ECF from incorrect V from (a)(i).

time taken $t \left(= \frac{Q}{I} \right) = \frac{1.11 \times 10^{-7}}{3.2 \times 10^{-8}} = 3.5 \text{ (3.47) (s) } \checkmark$

2

(b) (i) time increases \checkmark

(larger C means) more charge required (to reach breakdown pd)

Mark sequentially i.e. no explanation mark if effect is

wrong.

or $t = \frac{CV}{I}$ or time \propto capacitance ✓

2

(ii) spark is brighter (or lasts for a longer time) ✓

more energy (or charge) is stored or current is larger

Mark sequentially.

or spark has more energy ✓

²
(Total 7 marks)