1.

Two parallel metal plates of separation a carry equal and opposite charges.



Which graph best represents how the electric field strength E varies with the distance x in the space between the two plates?







3.

4.

A particle of mass m and charge q is accelerated through a potential difference V over a distance d.

What is the average acceleration of the particle?



## (Total 1 mark)

An electron on the surface of the Earth is placed in an electric field of strength 5000 N  $C^{-1}$ .

| What is | electric force gravitational force | for the electron? |
|---------|------------------------------------|-------------------|
|         | (8)                                |                   |

| Α | 1.1 × 10 <sup>-14</sup> | 0 |
|---|-------------------------|---|
| в | 2.9 × 10 <sup>-10</sup> | 0 |
| С | 3.4 × 10 <sup>9</sup>   | 0 |
| D | 9.0 × 10 <sup>13</sup>  | 0 |

## (Total 1 mark)

An  $\alpha$  particle makes a head-on collision with a gold nucleus containing 79 protons. The distance of closest approach of the  $\alpha$  particle to the nucleus is 4.0 × 10<sup>-14</sup> m.

What electrostatic force acts on the gold nucleus when at this separation?





Two fixed parallel metal plates **P** and **Q** are at constant electrical potentials of +100 V and +70 V respectively. A proton travelling from **P** to **Q** experiences a force *F* due to the electric field between **P** and **Q**, and a change of potential energy of  $\Delta E_{p}$ .



Which line, **A** to **B**, in the table gives the direction of *F* and the value of  $\Delta E_{p}$ ?

|   | Direction of $m{F}$ | $\Delta E_{p}$ |   |
|---|---------------------|----------------|---|
| Α | towards <b>P</b>    | +30 eV         | 0 |
| В | towards <b>Q</b>    | +30 eV         | 0 |
| С | towards <b>Q</b>    | -30 eV         | 0 |
| D | towards P           | -30 eV         | 0 |

## (Total 1 mark)

An electron moves through a distance of 0.10 m parallel to the field lines of a uniform electric field of strength 2.0 kN  $C^{-1}$ .

What is the work done on the electron?

A zero

6.

0

0

0

0

- **B** 1.6 × 10<sup>-17</sup> J
- **C** 3.2 × 10<sup>-17</sup> J
- **D** 1.6 × 10<sup>-21</sup> J





The total potential at the centre of the square, a distance d from each charge, is  $\frac{5Q}{4\pi\varepsilon_o d}$ 

Three of the charges have a charge of +Q

What is the magnitude of the fourth charge?

| Α | $-\frac{7Q}{4}$ | $\circ$ |
|---|-----------------|---------|
| в | Q               | 0       |
| С | $\sqrt{2}Q$     | 0       |
| D | 2 <i>Q</i>      | $\circ$ |

## (Total 1 mark)

8.

A charged spherical conductor has a radius r. An electric field of strength E exists at the surface due to the charge.

What is the potential of the spherical conductor?





A conducting sphere holding a charge of +10  $\mu C$  is placed centrally inside a second uncharged conducting sphere.

Which diagram shows the electric field lines for the system?





в









PhysicsAndMathsTutor.com



11.

The ionisation potential for the atoms of a gas is V. Electrons of mass m and charge e travelling at a speed v can just cause ionisation of atoms in the gas.

What is V?



(Total 1 mark)

An electric field acts into the plane of the paper. An electron enters the field at 90° to the field lines.

The force on the electron is

| Α | zero.                                   | 0 |
|---|---|---|
| В | along the direction of the field.       | 0 |
| С | at 90° to the field.                    | 0 |
| D | opposite to the direction of the field. | 0 |
|   |   |   |



A positive charge of 2.0 ×  $10^{-4}$  C is placed in an electric field at a point where the potential is +500 V.

What is the potential energy of the system?

- **A**  $1.0 \times 10^{-1} \text{ J}$
- **B**  $1.0 \times 10^{-1} \text{ J C}^{-1}$
- **C** 4.0 × 10<sup>-7</sup> J



0

 $^{\circ}$ 

 $^{\circ}$ 

**D**  $4.0 \times 10^{-7} \text{ J C}^{-1}$ 



Which diagram shows lines of equipotential in steps of equal potential difference near an isolated point charge?





С

D

0



Two fixed charges of magnitude +Q and +3Q repel each other with a force F. An additional charge of -2Q is given to each charge.

What are the magnitude and the direction of the force between the charges?

|   | Magnitude of force | Direction of force |   |
|---|--------------------|--------------------|---|
| A | $\frac{F}{3}$      | repulsive          | 0 |
| В | 5F                 | attractive         | 0 |
| С | 5 <i>F</i>         | repulsive          | 0 |
| D | $\frac{F}{3}$      | attractive         | 0 |

## (Total 1 mark)

**15.** At a distance L from a fixed point charge, the electric field strength is E and the electric potential is V.

What are the electric field strength and the electric potential at a distance 3L from the charge?

|   | Electric field strength | Electric potential |   |
|---|-------------------------|--------------------|---|
| Α | $\frac{E}{3}$           | <u>V</u><br>9      | 0 |
| В | $\frac{E}{3}$           | $\frac{V}{3}$      | 0 |
| С | <u>E</u><br>9           | $\frac{V}{3}$      | 0 |
| D | <u>E</u><br>9           | $\frac{V}{9}$      | 0 |



The diagram shows a particle with charge +Q and a particle with charge -Q separated by a distance d.

The particles exert a force F on each other.



An additional charge of +2Q is then given to each particle and their separation is increased to 2d.

What is the force that now acts between the particles?



17.

Two protons are separated by distance r. The electrostatic force between the two protons is **X** times the gravitational force between them.

What is the best estimate for X?



<sup>(</sup>Total 1 mark)



Two parallel metal plates separated by a distance d have a potential difference V across them. A particle with charge Q is placed midway between the plates.



What is the magnitude of the electrostatic force acting on the particle?



(Total 1 mark)



Two charged particles **P** and **Q** are separated by a distance of 120 mm. **X** is a point on the line between **P** and **Q** where the electric potential is zero.



What is the distance from **P** to **X**?





21.

An isolated spherical conductor is charged.

The conductor has a radius R and an electric potential V. The electric field strength at its surface is E.



Point **T** is a distance 2R from the surface.

What are the electric field strength and electric potential at T?

|   | Electric field strength | Electric potential |   |
|---|-------------------------|--------------------|---|
| Α | $\frac{E}{2}$           | <u>V</u><br>4      | 0 |
| В | $\frac{E}{3}$           | <u>V</u><br>9      | 0 |
| С | <u>E</u><br>4           | <u>V</u><br>2      | 0 |
| D | <u>E</u><br>9           | $\frac{V}{3}$      | 0 |

(Total 1 mark)

**O** is the centre of a negatively charged sphere.



**K** and **L** are two points at a distance  $r_1$  from **O**. **M** and **N** are two points at a distance  $r_2$  from **O**. Which statement is true?



A small object of mass m has a charge Q. The object remains stationary in an evacuated space between two horizontal plates. The plates are separated by a distance d and the potential difference between the plates is V.



What is V?





1.5 mJ of work is done when a charge of 30  $\mu C$  is moved between two points,  $\bm{M}$  and  $\bm{N},$  in an electric field.

What is the potential difference between  $\mathbf{M}$  and  $\mathbf{N}$ ?

| Α | 20 mV | 0 |
|---|-------|---|
| В | 20 V  | 0 |
| С | 45 V  | 0 |
| D | 50 V  | 0 |

## (Total 1 mark)

24.

A parallel-plate capacitor is fully charged and then disconnected from the power supply. A dielectric is then inserted between the plates.

Which row correctly identifies the charge on the plates and the electric field strength between the plates?

|   | Charge         | Electric field strength |   |
|---|----------------|-------------------------|---|
| Α | Stays the same | Increases               | 0 |
| В | Increases      | Decreases               | 0 |
| С | Increases      | Increases               | 0 |
| D | Stays the same | Decreases               | 0 |