

1(a). Technetium-99m (${}_{43}^{99m}\text{Tc}$) is a metastable radioisotope which can be used as a tracer in medical diagnosis. It is injected into the body and decays by gamma emission into technetium-99 according to the following chemical equation.



- i. A technetium-99m tracer with an activity of 900 MBq is injected into a body. The half-life of technetium-99m is 6.01 hours.

Calculate the number of technetium-99m nuclei initially present in the tracer.

number = [3]

- ii. Calculate the time in hours taken for the activity of the tracer to have fallen to 3.0% of its initial activity.

time = hours [3]


(b). The daughter nucleus (${}_{43}^{99}\text{Tc}$) decays by beta emission with a half-life of a little over 200 000 years. Approximately 50% of it is stored in the bones, and 50% is passed out of the body.

Suggest why the presence of this remaining ${}_{43}^{99}\text{Tc}$ in the body causes little additional risk to the patient.

[1]

(c). The half-life of a radioisotope is to be determined using suitable apparatus. Each count represents one decay. The number of counts is measured for one minute every hour over a period of 6 hours. When the data has been collected, a graph of $\ln(\text{corrected count rate} / \text{minute})$ against time is plotted and shown below.

2(a). The diagram below shows the average nuclear binding energy per nucleon for a number of different isotopes.



Explain what is meant by *nuclear binding energy* of a nucleus.

[1]

(b). Suggest why the ^1_1H isotope of hydrogen has **not** been included on the above diagram.

[1]

(c). The main nuclear fusion reaction in the Sun is between nuclei of deuterium (^2_1H) and tritium (^3_1H).

This reaction can be written as shown below.



i. Explain why isotopes with low mass numbers, such as hydrogen, are those which undergo nuclear fusion.

[1]

ii. Use the diagram given at the start of this question to show that, for the reaction of deuterium and tritium, the energy released in each fusion event is approximately 3×10^{-12} J.

[3]

- iii. The Sun's mass decreases by 4.3×10^9 kg every second. Assume that the mass loss is only due to this reaction.

Calculate the number of fusion events per second occurring in the Sun.

number of fusion events per second = s^{-1} [2]

- (d). In the Sun, deuterium (${}^2_1\text{H}$) is produced from fusion of two hydrogen (${}^1_1\text{H}$) nuclei, as shown below. There is a particle missing



- i. Determine the charge of the missing particle.

..... [1]

- ii. The missing particle is a lepton. Name this lepton.

..... [1]

- iii. In the fusion reaction above, determine the total number of up quarks at the **start** of the reaction.

..... [1]

- (e). Tritium (${}^3_1\text{H}$) is another isotope of hydrogen which is formed in stars. On the Earth, tritium is a radioactive element which decays by β^- emission.

Write down the equation for β^- decay in terms of quarks.

[2]

3. This question is about the rate of decay of a radioactive source.

Which of the following statements is/are true?

The rate of decay is

- 1 dependent on the decay constant.
- 2 independent of the mass of the source.
- 3 dependent on time.

- A 1 only
- B 1 and 3
- C 2 only
- D 2 and 3

Your answer

[1]

4. What is the radius of a carbon nucleus that has 6 protons and 7 neutrons?

Assume that the average radius of a nucleon r_0 is 1.2 fm.

- A 2.2 fm
- B 2.3 fm
- C 2.8 fm
- D 1.6 fm

Your answer

[1]

5. A sub-atomic particle has a positive charge.

Which type of particle is it?

- A anti-proton
- B down quark
- C neutrino
- D positron

Your answer

[1]

6. A neutrino is a fundamental particle.

Which row of the table correctly describes a neutrino?

	Classification	Force felt
A	hadron	strong nuclear
B	hadron	weak nuclear
C	lepton	strong nuclear
D	lepton	weak nuclear

Your answer

[1]

7. Which one of these non-invasive medical scans does **not** expose the patient to ionising radiation?.

- A CAT
- B PET
- C Ultrasound
- D X-ray

Your answer

[1]

8. A pulsar is a rapidly rotating neutron star that emits radio waves.

A typical neutron star can be modelled as a sphere with mass $\approx 2 \times 10^{30}$ kg and radius ≈ 10 km.

Show that the average density of a neutron star is similar to the average density of an atomic nucleus.

- radius of a nucleon ≈ 1 fm

[3]

9. Large power stations generate an electrical power of about 1 GW.

Current methods of energy production that use nuclear fusion are unable to produce enough energy for large-scale energy production. A proposed method of controlling nuclear fusion is inertial confinement fusion (ICF). ICF uses a large number of powerful lasers to create the high temperatures required for nuclear fusion to occur.

One ICF experiment uses a network of capacitors to store the energy needed to power the lasers. When the network is fully charged:

- potential difference across the network = 24 kV
- total energy stored in the network = 400 MJ

The fusion reaction in the ICF experiment is

deuterium + tritium \rightarrow alpha particle + neutron

Calculate the number of fusion reactions that must occur for the energy released by fusion to be equal to the electrical energy stored in the network of capacitors.

- mass of deuterium = 2.014102 u
- mass of tritium = 3.016049 u
- mass of alpha particle = 4.002603 u
- mass of neutron = 1.008665 u

number of fusion reactions =**[4]**

10(a). Radiographers commonly use molecules containing fluorine F-18 as tracers in positron emission tomography (PET) scanning.

Fluorine has a proton number of 9.

F-18 decays to oxygen (O) by β^+ decay.

Write the equation for the decay of a nucleus of F-18 using nuclear notation.

[2]

(b). The β^+ particle (positron) produced travels only a short distance in the patient before it meets an electron and is annihilated.

Calculate the wavelength λ of gamma photons produced.

$\lambda = \dots\dots\dots\text{m}$ [3]

(c). X-rays and gamma-rays are produced by different physical processes.

Briefly describe both processes.

----- [2]


(d). F-18 has a half-life of 109.7 minutes.

Explain the advantage that this has for the patient but the disadvantage that this has for the radiographers.

----- [3]

[6]

(c). The diagram shows a simplified layout of a nuclear fission reactor used in a nuclear power station.



Complete the labels on the diagram

[2]

12. Which of these statements is/are true?

- 1 Antiprotons are hadrons so are subject to the strong nuclear and weak nuclear forces.
- 2 Neutrons are subject to the weak nuclear force only.
- 3 The weak nuclear force is the only force that causes a change of quark type.

- A** 1, 2 and 3
B Only 1 and 2
C Only 1 and 3
D Only 3

Your answer

[1]

17. A gamma-ray photon of frequency 6.76×10^{22} Hz creates a particle-antiparticle pair. The particle-antiparticle pair have zero kinetic energy.

What is the mass of the particle?

- A 2.49×10^{-28} kg
- B 4.98×10^{-28} kg
- C 7.47×10^{-20} kg
- D 4.48×10^{-11} kg

Your answer

[1]

18. The half-life of fluorine-18 isotope is T .
After time $t = 4T$ the number of fluorine-18 nuclei in a source is N .

How many fluorine-18 nuclei have decayed in the time interval from $t = 0$ to $t = 4T$?

- A $3N$
- B $4N$
- C $15N$
- D $16N$

Your answer

[1]

19. The activity of an alpha-emitting source is 120 kBq. The kinetic energy of each alpha-particle is 4.0 MeV.

What is the rate of energy released by the source?

- A 6.4×10^{-13} W
- B 4.8×10^{-8} W
- C 7.7×10^{-8} W
- D 1.2×10^5 W

Your answer

[1]

END OF QUESTION PAPER