

# Cambridge IGCSE™

PHYSICS

0625/42 February/March 2024

Paper 4 Extended Theory MARK SCHEME Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the February/March 2024 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

## **Generic Marking Principles**

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

**GENERIC MARKING PRINCIPLE 1:** 

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question .
- the specific skills defined in the mark scheme or in the generic level descriptors for the question .
- the standard of response required by a candidate as exemplified by the standardisation scripts.

**GENERIC MARKING PRINCIPLE 2:** 

Marks awarded are always whole marks (not half marks, or other fractions).

**GENERIC MARKING PRINCIPLE 3:** 

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond ٠ the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do ٠
- marks are not deducted for errors •
- marks are not deducted for omissions .
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the • guestion as indicated by the mark scheme. The meaning, however, should be unambiguous.

**GENERIC MARKING PRINCIPLE 4:** 

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

PMT

#### **GENERIC MARKING PRINCIPLE 5:**

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

#### **GENERIC MARKING PRINCIPLE 6:**

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thre sholds or grade descriptors in mind.

## Science-Specific Marking Principles

- 1 Examiners should consider the context and scientific use of any keywords when awarding marks. Although keywords may be present, marks should not be awarded if the keywords are used incorrectly.
- 2 The examiner should not choose between contradictory statements given in the same question part, and credit should not be awarded for any correct statement that is contradicted within the same question part. Wrong science that is irrelevant to the question should be ignored.
- 3 Although spellings do not have to be correct, spellings of syllabus terms must allow for clear and unambiguous separation from other syllabus terms with which they may be confused (e.g. ethane / ethene, glucagon / glycogen, refraction / reflection).
- 4 The error carried forward (ecf) principle should be applied, where appropriate. If an incorrect answer is subsequently used in a scientifically correct way, the candidate should be awarded these subsequent marking points. Further guidance will be included in the mark scheme where necessary and any exceptions to this general principle will be noted.
- 5 <u>'List rule' guidance</u>

For questions that require *n* responses (e.g. State **two** reasons ...):

- The response should be read as continuous prose, even when numbered answer spaces are provided.
- Any response marked *ignore* in the mark scheme should not count towards *n*.
- Incorrect responses should not be awarded credit but will still count towards *n*.
- Read the entire response to check for any responses that contradict those that would otherwise be credited. Credit should **not** be awarded for any responses that are contradicted within the rest of the response. Where two responses contradict one another, this should be treated as a single incorrect response.
- Non-contradictory responses after the first *n* responses may be ignored even if they include incorrect science.

PMT

#### 6 <u>Calculation specific guidance</u>

Correct answers to calculations should be given full credit even if there is no working or incorrect working, **unless** the question states 'show your working'.

For questions in which the number of significant figures required is not stated, credit should be awarded for correct answers when rounded by the examiner to the number of significant figures given in the mark scheme. This may not apply to measured values.

For answers given in standard form (e.g.  $a \times 10^n$ ) in which the convention of restricting the value of the coefficient (a) to a value between 1 and 10 is not followed, credit may still be awarded if the answer can be converted to the answer given in the mark scheme.

Unless a separate mark is given for a unit, a missing or incorrect unit will normally mean that the final calculation mark is not awarded. Exceptions to this general principle will be noted in the mark scheme.

#### 7 <u>Guidance for chemical equations</u>

Multiples / fractions of coefficients used in chemical equations are acceptable unless stated otherwise in the mark scheme.

State symbols given in an equation should be ignored unless asked for in the question or stated otherwise in the mark scheme.

#### Acronyms and shorthand in the mark scheme.

acronym/shorthand	explanation
A mark	Final answer mark which is awarded for fully correct final answers including the unit.
C mark	Compensatory mark which may be scored when the final answer (A) mark for a question has not been awarded.
B mark	Independent mark which does not depend on any other mark.
M mark	Method mark which must be scored before any subsequent final answer (A) mark can be scored.
Brackets()	Words not explicitly needed in an answer, however if a contradictory word/phrase/unit to that in the brackets is seen the mark is not awarded.
<u>Underlining</u>	The underlined word (or a synonym) must be present for the mark to be scored. If the word is a technical scientific term, the word must be there.
/ or <b>OR</b>	Alternative answers any one of which gains the credit for that mark.
owtte	Or words to that effect.
ignore	Indicates either an incorrect or irrelevant point which may be disregarded, i.e., not treated as contradictory.
insufficient	an answer not worthy of credit <u>on its own</u> .
CON	An incorrect point which contradicts any correct point and means the mark cannot be scored.
ecf [question part]	Indicates that a candidate using an erroneous value from the stated question part must be given credit here if the erroneous value is used correctly here.
сао	correct answer only

PMT

Cambridge IGCSE – Mark Scheme
PUBLISHED

PMT

Question	Answer	Marks
1(a)	constant speed	B1
	constant / uniform deceleration	B1
	stationary	B1
1(b)(i)	$7.1 \times 10^5$ J <b>OR</b> 710 000 J <b>OR</b> 710 kJ	A2
	$E_{\rm K}=\frac{1}{2}mv^2$ OR ( $E_{\rm K}=$ ) $\frac{1}{2}mv^2$ OR $\frac{1}{2}\times$ 18 000 × (8.9) <sup>2</sup>	(C1)
1(b)(ii)	31 000 N <b>OR</b> 31 kN	A3
	W = Fd OR (F =) W/ d OR 710 000/23	(C1)
	F = 710 000 / 23	(C1)

Question	Answer	Marks
2(a)	(Impulse =) force × time (for which force acts)	B1
2(b)(i)	$F \Delta t = \Delta \{mv\} \mathbf{OR} (F =) \Delta \{mv\} / \Delta t \mathbf{OR} (F =) \Delta p / t$	M1
	$\{2800 \times 1400\}$ (/ 1) = 3 920 000 N <b>OR</b> $\{2800 \times 1400\}$ (/ 1) = 3920 kN	A1
2(b)(ii)	$4.0 \times 10^{5}$ kg <b>OR</b> 400 000 kg	A3
	(at maximum mass) force = weight of rocket <b>OR</b> F = mg	(C1)
	( <i>m</i> =) <i>F</i> /g <b>OR</b> 3.9 × 10 <sup>6</sup> /9.8 <b>OR</b> 4(.0) × 10 <sup>N</sup> (kg)	(C1)

Question	Answer	Marks
3(a)(i)	2.0 × 10 <sup>5</sup> Pa <b>OR</b> 200 000 Pa <b>OR</b> 200 kPa	A2
	(P=) F/A <b>OR</b> 13 000 / (0.016 × 4)	(C1)
3(a)(ii)	<ul> <li>Any four from:</li> <li>friction between road and tyre</li> <li>temperature of air / tyre increases</li> <li>particles (of air in tyre) move faster</li> <li>particles (of air) collide harder with the <u>walls</u> (of the tyre) <b>OR</b> particles (of air) collide more frequently with the <u>walls</u> (of the tyre)</li> <li>force (on the tyre wall) increases, <b>AND</b> area (of tyre) is constant (so tyre pressure increases)</li> </ul>	B4
3(b)	(maximum number = ) 34	A3
	$pV$ = constant <b>OR</b> {2.0 × 10 <sup>6</sup> × 0.026} = 1.0 × 10 <sup>5</sup> × V <sub>2</sub>	(C1)
	$V_2 = 0.52 \text{ (m}^3)$ <b>OR (</b> $V_2 =$ ) {2.0 × 10 <sup>6</sup> × 0.026} / 1.0 × 10 <sup>5</sup> <b>OR</b> (number of balloons =) {2.0 × 10 <sup>6</sup> × 0.026} / {1.0 × 10 <sup>5</sup> × 0.015}	(C1)

Question	Answer	Marks
4(a)	energy transferred per unit mass per unit temperature change	A2
	(thermal) energy (transferred) per unit temperature change	(C1)
4(b)(i)	( <i>m</i> =) 2.2 kg	A2
	$m = \rho V$ in any form <b>OR</b> 910 × 0.0024	(C1)
4(b)(ii)	$7.0 \times 10^5 \text{ J}$ <b>OR</b> 700 000 J	A2
	$c = \Delta E / \{ m \Delta \theta \}$ OR ( $\Delta E =$ ) $mc \Delta \theta$ OR 2.2 × 2000 × 160 OR 2.184 × 2000 × 160	(C1)

0625/42

## Cambridge IGCSE – Mark Scheme **PUBLISHED**

Question	Answer	Marks
4(b)(iii)	1700 W	A2
	( <i>P</i> =) <i>E</i> / <i>t</i> <b>OR</b> 700 000 / 7 × 60 <b>OR</b> 704 000 / 7 × 60 <b>OR</b> 698 880 / 7 × 60	(C1)

Question		Answer	Mar
5(a)(i)	application	region of electromagnetic spectrum	
	cancer treatment	gamma rays	
	bluetooth	radio waves	
	optical fibres	infrared	
	security marking	security marking ultraviolet	
	sterilising food gamma rays		
	wireless internet	Microwaves	
5(a)(ii)	$3.0 \times 10^8 (m /s)$ OR 300 000 000	(m/s)	
5(b)(i)	three crests parallel to the barrie		
	same wavelength as wave after th	ne gap	
5(b)(ii)	central part of crest (parallel to th	e (gap in the) barrier) is straight	
	crests have curved ends		

Question	Answer	Marks
6(a)	ray from top of object to tip of image	M1
	line labelled L drawn perpendicular to principal axis at its intersection with previous ray	A1
6(b)	ray from top of O parallel to principal axis to lens <b>AND</b> ray from lens to tip of   <b>OR</b> ray from tip of   parallel to principal axis to lens <b>AND</b> ray from lens to top of O	B1
	2.1 cm	B1
6(c)	virtual	B1
	upright	B1

Question	Answer	Marks
7(a)		B1
7(b)(i)	1.5 V	A2
	Vout / VR = Rout / R OR VR = 3 Vout	(C1)
7(b)(ii)	0.51 C	A2
	$I = Q / t \text{ OR } (Q =) / t \text{ OR } 1.7 \times 10^{-3} \times 300$	(C1)

Question	Answer	Marks
8(a)(i)	three concentric circles centred on X	B1
	second and third circles further apart than first and second circles	B1
	direction of arrows clockwise	B1
8(a)(ii)	strength of magnetic field increases	B1
	its direction reverses	B1
8(b)	<ul> <li>any three from:</li> <li>P = /<sup>2</sup>R OR power (loss) = /<sup>2</sup>R</li> <li>(high voltage allows) low current (so at same power output, less power / energy lost)</li> <li>thin wires have high resistance (so more power / energy lost)</li> <li>(low) current has a greater effect (on efficiency) than (high) resistance (of the thin wires)</li> </ul>	B3

Question	Answer	Marks
9(a)(i)	most of the atom is empty space	B1
9(a)(ii)	any <b>two</b> from: 1 the nucleus is very small 2 mass of gold nucleus is much greater than mass of alpha particle 3 the nucleus is positively charged	B2
	<ul> <li>corresponding explanation to conclusion:</li> <li>1 not many alpha particles pass close to the nucleus / owtte</li> <li>2 large force between alpha and nucleus (has bigger effect on small mass of alpha)</li> <li>3 alpha particles are positively charged, AND force is repulsive</li> </ul>	B2
9(b)(i)	(in the nucleus a) neutron is changed into a proton (and an electron which is the emitted $\beta$ -particle)	B1

Question	Answer	Marks		
9(b)(ii)	22 µg	A3		
	(87 years is) three half-lives <b>OR</b> 25 / 8 <b>OR</b> 87 / 29 = 3 (half lives)	(C1)		
	1 / 8th (of the strontium remains) <b>OR</b> 25 / 8 (decays) <b>OR</b> 3.125 seen	(C1)		

Question	Answer	Marks
10(a)(i)	position A and / or E	B1
	position G	B1
10(a)(ii)	1 month	B1
10(b)(i)	110 000 (km / h)	A3
	$(v = )2\pi r / T$	(C1)
	$T = 365 \times 24h \text{ OR } 2\pi \times 1.5 \times 10^8/365 \times 24$	(C1)
10(b)(ii)	500 s	A2
	$v = s / t \text{ OR } (t =) s / v \text{ OR } 1.5 \times 10^{11} / 3.0 \times 10^{8}$	(C1)

Question	Answer	Marks
11(a)	inward force / force of gravitational attraction, is balanced by an outward force / force due to fusion reactions	B1
11(b)(i)	ratio of the speed at which the galaxy is moving away from the Earth/observer to its distance (from the Earth/observer)	A2
	ratio of speed (of a galaxy) to distance (away from observer)	(C1)

February/March 2024

0625/42

## Cambridge IGCSE – Mark Scheme PUBLISHED

PUBLISHEDQuestionAnswerMarks11(b)(ii)(age of universe =)  $d/v = 1/H_0$  OR age (of universe) =  $1/H_0$ B111(b)(iii) $4.5 \times 10^{17}$  (s)B1