Surname	Centre Number	Candidate Number
Other Names		0



## New GCSE

4473/01

## ADDITIONAL SCIENCE FOUNDATION TIER PHYSICS 2

P.M. THURSDAY, 17 January 2013

l hour

For Examiner's use only				
Question	Maximum Mark	Mark Awarded		
1.	7			
2.	9			
3.	7			
4.	10			
5.	9			
6.	12			
7.	6			
Total	60			

#### **ADDITIONAL MATERIALS**

In addition to this paper you may require a calculator.

#### INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use a gel pen. Do not use correction fluid.

Write your name, centre number and candidate number in the spaces at the top of this page.

Answer all questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the continuation page at the back of the booklet, taking care to number the question(s) correctly.

### INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

A list of equations is printed on page 2. In calculations you should show all your working.

You are reminded that assessment will take into account the quality of written communication (QWC) used in your answer to question 7.



### Equations

power = voltage × current	P = VI
resistance = $\frac{\text{voltage}}{\text{current}}$	$R = \frac{V}{I}$
speed = $\frac{\text{distance}}{\text{time}}$	
acceleration [or deceleration] = $\frac{\text{change in velocity}}{\text{time}}$	$a = \frac{\Delta v}{t}$
acceleration = gradient of a velocity-time graph	
momentum = mass × velocity	p = mv
resultant force = mass × acceleration	F = ma
force = $\frac{\text{change in momentum}}{\text{time}}$	$F = \frac{\Delta p}{t}$
work = force × distance	W = Fd

### SI multipliers

Prefix	Multiplier
m	10 <sup>-3</sup>
k	10 <sup>3</sup>
М	10 <sup>6</sup>



**BLANK PAGE** 


3


# PLEASE DO NOT WRITE ON THIS PAGE



© WJEC CBAC Ltd.












0 5





	A <b>AN</b> TA A				
	(c)	Narr	Braking force =	N [1]	
		(iii)	Use an equation from page 2 to calculate the braking force that brought ta stop.	the car to [2]	
		(ii)	What is the change in momentum of the car in coming to a stop?	[1]	
		(i)	What is the momentum of the car when it stops at traffic lights?	[1]	
	(b)	As the from	he car approaches traffic lights, they change from green to red. The car slo $12 \text{ m/s}$ to $0 \text{ m/s}$ in $3 \text{ s}$ .	ows down	
			Momentum =	kg m/s	
	(a)	Use	an equation from page 2 to calculate the momentum of the car.	[2]	
3.	The	diagra	m shows a car of mass 800 kg travelling at 12 m/s.		9
		(ii)	Give a reason for your answer.	[1]	
	(c)	(i)	Compare the speed of the cyclist in the first 40 seconds and the last 40 se	conds. [1]	









Examiner

5. The diagram shows a circuit used to investigate currents in a parallel circuit when the voltage is varied. A lamp L and a wire W are connected in parallel with a variable voltage supply. The circuit has 3 ammeters  $A_1$ ,  $A_2$  and  $A_3$  as shown.



The currents through the lamp, L, and the wire, W, depend on the voltage applied to them in the way shown on the graph below.





a)	(i)	Use the graph to find the current through the lamp when the voltmeter readin $6 V$ .	ig is [1]
		Current =	A
	(ii)	Using an equation from page 2, calculate the resistance of the lamp at 6 V.	[2]
		Resistance =	Ω
	(iii)	Using an equation from page 2, calculate the power produced by the lamp at 6	5 V. [2]
		Power =	W
	(iv)	At what voltage, were the power of the lamp and wire the same?	V [1]
	(v)	Find the current through ammeter $A_1$ at 6 V.	[1]
		Current =	A
5)	The	voltage supply in the diagram is increased from 6 V to 12 V.	
	(i)	Compare the resistances of the lamp and wire at 12 V.	[1]
	(ii)	Give a reason for your answer.	[1]

1 1

<ul> <li>It uses a radioactive source that emits alpha particles.</li> <li>The alpha particles ionise the air inside the detector causing an electric current.</li> <li>Any smoke getting into the detector absorbs the alpha particles and changes the current.</li> <li>The change in current sets off the alarm.</li> <li>(a) (i) What is an alpha particle? [I]</li> <li>(ii) Explain why the detector would not work if the radioactive source emitted gamma rays only. [2]</li> <li>(iii) Explain why, in normal use, the radioactive source in the detector is not a risk to human health. [2]</li> <li>(b) Americium-241 has a half-life of 432 years. Curium-242 has a half-life of 160 days. Both isotopes are alpha emitters. [2]</li> <li>(i) Explain why Americium-241 is more suitable for use in the smoke detector than Curium-242. [2]</li> </ul>	A sm	oke d	etector works as follows:	Exam on
<ul> <li>(ii) Explain why the detector would not work if the radioactive source emitted gamma rays only. [2]</li> <li>(iii) Explain why, in normal use, the radioactive source in the detector is not a risk to human health. [2]</li> <li>(b) Americium-241 has a half-life of 432 years. Curium-242 has a half-life of 160 days. Both isotopes are alpha emitters.</li> <li>(i) Explain why Americium-241 is more suitable for use in the smoke detector than</li> </ul>	• The • An	e alph y smo	a particles ionise the air inside the detector causing an electric current. oke getting into the detector absorbs the alpha particles and changes the current.	
rays only.       [2]         (iii)       Explain why, in normal use, the radioactive source in the detector is not a risk to human health.         [2]       [2]         (b)       Americium-241 has a half-life of 432 years. Curium-242 has a half-life of 160 days. Both isotopes are alpha emitters.         (i)       Explain why Americium-241 is more suitable for use in the smoke detector than	<i>(a)</i>	(i)	What is an alpha particle?[1]	
<ul> <li>human health. [2]</li> <li>(b) Americium-241 has a half-life of 432 years. Curium-242 has a half-life of 160 days. Both isotopes are alpha emitters.</li> <li>(i) Explain why Americium-241 is more suitable for use in the smoke detector than</li> </ul>		(ii)		
<ul> <li>human health. [2]</li> <li>(b) Americium-241 has a half-life of 432 years. Curium-242 has a half-life of 160 days. Both isotopes are alpha emitters.</li> <li>(i) Explain why Americium-241 is more suitable for use in the smoke detector than</li> </ul>				
<ul><li>isotopes are alpha emitters.</li><li>(i) Explain why Americium-241 is more suitable for use in the smoke detector than</li></ul>				
	( <i>b</i> )			
		(i)		
		······		

			]	Examiner
(ii)	An a whic	werage smoke detector contains about 0.4 micrograms (μg) of Americium h has an initial activity of 52 000 units.		only
	(I)	Name the unit of activity.	[1]	
	(II)	Calculate how long it will take for the activity to drop to 26 000 units.	[2]	
		Time = y	ears	
	(III)	Calculate the mass of Americium-241 remaining after 864 years.	[2]	
		Mass remaining =	µg	
				12



Speed (mph)	Thinking Distance (m)	Braking Distance (m)	Total Stopping Distance (m)	
60	18	55	73	1
70	21	75		1
80	24	97.5		1

# **BLANK PAGE**

# PLEASE DO NOT WRITE ON THIS PAGE



Question number	Additional page, if required. Write the question numbers in the left-hand margin.	Examiner only
		1

