Surname	Centre Number	Candidate Number
Other Names		0



New GCSE

4473/02	

ADDITIONAL SCIENCE HIGHER TIER PHYSICS 2

P.M. THURSDAY, 17 January 2013

l hour

For Examiner's use only				
Question	Maximum Mark	Mark Awarded		
1.	8			
2.	12			
3.	6			
4.	7			
5.	15			
6.	8			
7.	4			
Total	60			

ADDITIONAL MATERIALS

In addition to this paper you may require a calculator.

INSTRUCTIONS TO CANDIDATES

Use black ink or black ball-point pen. Do not use a gel pen. Do not use correction fluid.

Write your name, centre number and candidate number in the spaces at the top of this page. Answer **all** questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the continuation page at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

A list of equations is printed on page 2. In calculations you should show all your working.

You are reminded that assessment will take into account the quality of written communication (QWC) used in your answers to questions 3 and 6(b).



Equations

power = voltage × current	P = VI
resistance = $\frac{\text{voltage}}{\text{current}}$	$R = \frac{V}{I}$
power = $current^2 \times resistance$	$P = I^2 R$
speed = $\frac{\text{distance}}{\text{time}}$	
acceleration [or deceleration] = $\frac{\text{change in velocity}}{\text{time}}$	$a = \frac{\Delta v}{t}$
acceleration = gradient of a velocity-time graph	
distance travelled = area under a velocity-time graph	
momentum = mass \times velocity	p = mv
resultant force = mass × acceleration	F = ma
$force = \frac{change in momentum}{time}$	$F = \frac{\Delta p}{t}$
work = force × distance	W = Fd
kinetic energy = $\frac{\text{mass} \times \text{speed}^2}{2}$	$KE = \frac{1}{2} mv^2$
change in = mass × gravitational × change potential energy field strength in height	PE = mgh

SI multipliers

Prefix	Multiplier
р	10^{-12}
n	10 ⁻⁹
μ	10^{-6}
m	10 ⁻³

Prefix	Multiplier
k	10 ³
М	10 ⁶
G	10 ⁹
Т	10 ¹²




BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE









© WJEC CBAC Ltd.

4473 020005

<i>(a)</i>	(i)	Use the graph to find the current through the lamp when the voltmeter reading is 6V. [1]	
		Current =A	
	(ii)	Using an equation from page 2, calculate the resistance of the lamp at 6 V. [2]	
		Resistance =Ω	2
	(iii)	Find the current through ammeter A_1 at 6 V. [1]	
		Current = A	
<i>(b)</i>	The	voltage supply in the diagram is increased from 6 V to 12 V.	
	(i)	Compare the resistances of the lamp and wire at 12 V. [1]	4473
	(ii)	Give a reason for your answer. [1]	
(c)	the p	the voltage is increased from 0 V to 12 V, the power of the wire is not always less than power of the lamp." lain why this statement is true. [2]	
			8



It i Th	ises a	radioactive source that emits alpha particles. a particles ionise the air inside the detector causing an electric current.
An	iy smo	ke getting into the detector absorbs the alpha particles and changes the current. nge in current sets off the alarm.
(a)	(i)	What is an alpha particle?
u)		
	(ii)	Explain why the detector would not work if the radioactive source emitted gamma rays only. [2]
	(iii)	Explain why, in normal use, the radioactive source in the detector is not a risk to human health. [2]
(b)		ericium-241 has a half-life of 432 years. Curium-242 has a half-life of 160 days. Both opes are alpha emitters.
	(i)	Explain why Americium-241 is more suitable for use in the smoke detector than Curium-242. [2]
	·····	




(ii) An average smoke detector contains about 0.4 micrograms (µg) of Americium-24I which has an initial activity of 52 000 units. (I) Name the unit of activity. [I] (II) Calculate how long it will take for the activity to drop to 26 000 units. [2] Time =years [1] (III) Calculate the mass of Americium-24I remaining after 864 years. [2] III) [2]					Examin only	er
(11) Calculate how long it will take for the activity to drop to 26 000 units. [2] Time =years (11) Calculate the mass of Americium-241 remaining after 864 years. [2] Mass remaining =µg 12	(ii)	An a whic	average smoke detector contains about 0.4 micrograms (µg) of Americium ch has an initial activity of 52 000 units.	ı - 241		
Time =years (III) Calculate the mass of Americium-241 remaining after 864 years. [2] [12] [12] [12] [12] [12] [12] [12]		(I)	Name the unit of activity.	[1]		
(III) Calculate the mass of Americium-241 remaining after 864 years. [2] Mass remaining = µg 12		(II)	Calculate how long it will take for the activity to drop to 26 000 units.	[2]		
Mass remaining =			Time =	vears		
		(III)	Calculate the mass of Americium-241 remaining after 864 years.	[2]		
			Mass remaining =	µg		1
					12	
						4473 020007
		ĥ				




Speed (mph)	Thinking Distance (m)	Braking Distance (m)	Total Stopping Distance (m)	
60	18	55	73	
70	21	75		
80	24	97.5		

	5 m
resist	
Use an equation from page 2 to calculate the loss of kinetic energy of the car.	[3]
Loss in kinetic energy = Use your answer to part (<i>a</i>) along with an equation from page 2 to find the (me resistive force produced by the sand during the collision.	
Resistive force = Write down the value of the horizontal force that acts on the sand in this collision. Force on the sand =	[1]
	7
	Use an equation from page 2 to calculate the loss of kinetic energy of the car. Loss in kinetic energy = Use your answer to part (a) along with an equation from page 2 to find the (me resistive force produced by the sand during the collision.



1 0



(a)		TE:	
	Explain what the term "isotope" means. [2]		
.			
••••••			
(b)	A fusion reaction can occur when two deuterium nuclei, ${}_{1}^{2}H$, collide at high speed to produce a helium nucleus (He) and release a neutron.		
	Explain the importance of this reaction along with its benefits and the difficulties in achieving it in a controlled manner. Your answer should include a balanced nuclear equation. [6 QWC]	·	
•••••			
·····			
·····			
·····			
·····			
.			
•••••			
		 r	

]Exan
Two filament lamps, X and Y, are connected in series. Lamp Y is brighter than lamp X.	on
Choose and write down an equation from page 2 to explain why lamp Y is brighter than lamp X . [Hint: consider power] [4]	
END OF PAPER	
	J



Question number	Additional page, if required. Write the question numbers in the left-hand margin.	Examiner only
		1
		·
		·
	1	1



BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

