

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

* 8 5 0 9 4 9 6 0 7

CO-ORDINATED SCIENCES

0654/42

Paper 4 Theory (Extended)

February/March 2021

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 120.
- The number of marks for each question or part question is shown in brackets [].
- The Periodic Table is printed in the question paper.

This document has 28 pages. Any blank pages are indicated.

1	(a)	State the balanced equation for photosynthesis.
		[2]
	(b)	A student investigates the effect of temperature on the rate of photosynthesis of an aquatic plant.
		The student counts the number of bubbles produced per minute at different temperatures.
		The number of bubbles produced per minute is an indication of the rate of photosynthesis.
		The results are shown in Table 1.1.

Table 1.1

temperature/°C	number of bubbles produced per minute
0	0
5	6
10	9
15	13
20	16
25	18
30	17
35	9
40	0

(i)	State the temperature that resulted in the fastest rate of photosynthesis.		
		°C	[1]
(ii)	Place ticks (✓) next to all the statements that explain the results between 0 °C	and 10	٥°C.
The kin	etic energy of the particles increases.		
There a	are more frequent collisions between the substrate and enzyme.		
Fewer	substrate molecules fit into the active site of enzymes.		
The ter	nperature affects the pH of the enzymes.		
The nu	mber of bubbles increases as more substrate is produced.		
			[2]

(iii)	Explain the results at 40 °C from Table 1.1.	
		[3]
(iv)	Explain why the light intensity is kept constant during this investigation.	
		[2]
		[Total: 10]

2 (a) Table 2.1 shows some information about the charges and relative masses of three subatomic particles.

Table 2.1

	particle A	particle B	particle C
charge	+1	no charge	-1
relative mass	1	1	1 1840

Use the information in Table 2.1 to state the names of particles A , B and C .	
particle A	
particle B	
particle C	[3]
	[U

(b) Fig. 2.1 shows a diagram of an atom of magnesium.




Fig. 2.1

(i)	State why this atom of magnesium has a proton number (atomic number) of 12 a nucleon number (mass number) of 24.	ınd a
	proton number of 12 because	
	nucleon number of 24 because	
(ii)	State the electronic structure of this atom of magnesium.	[2]
		F4 1

(c)	Magnesium reacts with oxygen to form magnesium oxide, MgO.
	Magnesium oxide is an ionic compound with a melting point of 2852 °C.
	Oxygen reacts with hydrogen to form water, H ₂ O.
	Water is a covalent compound with a melting point of 0 °C.
	Explain the difference in the melting points of magnesium oxide and water in terms of attractive forces.
	[3]
(d)	Magnesium oxide reacts with dilute hydrochloric acid, HC1.
	Magnesium chloride, ${\rm MgC}l_2$, is made.
	The balanced symbol equation for the reaction is shown.
	$MgO + 2HCl \rightarrow MgCl_2 + H_2O$
	In an experiment, 2.0 g of magnesium oxide reacts with excess dilute hydrochloric acid.
	Calculate the maximum mass of magnesium chloride that can be made.
	Show your working.
	[A _r : Cl, 35.5; H, 1; Mg, 24; O, 16]

mass of magnesium chloride = g [2]

[Total: 11]

3 Fig. 3.1 shows a tennis player throwing a ball in the air before the player hits the ball.

Э



Fig. 3.1

- (a) The ball has a mass of 56.25 g and is thrown vertically upwards with a velocity of 8.0 m/s.
 - (i) Calculate the kinetic energy of the ball immediately after it leaves the player's hand.

(b) Fig. 3.2 shows the tennis player hitting the same ball with the racket.



Fig. 3.2

This causes the ball to accelerate at 1600 m/s².

Calculate the force applied to the ball by the racket.

force = N [2]

(c) A student removes one of the nylon strings from the racket to investigate how it deforms when tensile forces are applied.

Fig. 3.3 shows the equipment used.




Fig. 3.3

The student adds masses to the mass hanger and records the extension of the nylon string.

Fig. 3.4 shows the results from this investigation.




Fig. 3.4

(i)	Use Fig.	3.4 to	find the	force	required	to (give an	extension	of	3 mm
•	•	000119.	0. 1 10	IIII G CIIO	.0.00	10941104		givo aii	OMIGIOIOII	٥.	01111111

[Total: 10]

(ii)	State Hooke's law.

(iii)	Describe how the graph in Fig. 3.4 shows that the nylon string does not obey Hooke' law.	S
		• •
		••

4 (a) A person monitored the concentration of glucose in their blood for 12 hours.

The results are shown in Fig. 4.1.




Fig. 4.1

	<u> </u>	
(i)	The concentration of glucose in the blood increases rapidly after a meal is eaten.	
	Suggest the number of meals the person had during the 12 hours.	
		[1]
(ii)	Complete the sentences to explain the results between 2 and 3 hours.	
	The brain detects an increase in the concentration of glucose in the blood.	
	The releases insulin.	
	The insulin stimulates the to convert glucose to	
	and store it.	
	This lowers the concentration of glucose in the blood.	[2]
		[3]
Nar	ne two normones that increase blood glucose concentration.	
1		
2		
		[2]
	· · · · · · · · · · · · · · · · · · ·	ose
		[1]
	(ii) Nar 1 2 Nar con	Suggest the number of meals the person had during the 12 hours. (ii) Complete the sentences to explain the results between 2 and 3 hours. The brain detects an increase in the concentration of glucose in the blood. The

(d)	Glucose is absorbed in the small intestine.		
	(i)	Define the term absorption.	
			[2]
	(ii)	Describe how the small intestine is adapted for efficient absorption.	
			[2]
		[Total:	113

5 Petroleum is a fossil fuel.

Petroleum can be separated into useful fractions by fractional distillation.

Fig. 5.1 shows a fractionating column.




Fig. 5.1

(a)	Explain why it is possible to separate the substances in petroleum by fractional distillation.	
		[1

(b) Table 5.1 shows the uses of some of the fractions.

Complete Table 5.1.

Table 5.1

fraction	use
refinery gas	bottled gas for heating
gasoline	fuel (petrol) in cars
naphtha	
diesel oil	
bitumen	

(c)	Butane is a hydrocarbon found in the refinery gas fraction.	
	Complete Fig. 5.2 to show the structure of a butane molecule.	
	Show all the covalent bonds.	
	H C H	
	Fig. 5.2	[2]
(d)	Methane, CH ₄ , is also a hydrocarbon found in the refinery gas fraction.	
	Methane burns in oxygen to form carbon dioxide and water.	
	Construct the balanced symbol equation for the burning of methane.	
		[2]
(e)	When methane burns, an exothermic reaction takes place.	
	State what is meant by an exothermic reaction.	
		[1]
(f)	The reaction of hydrogen with oxygen to make water is another exothermic reaction.	
	Look at the equation for this reaction. It shows all the atoms and all the bonds.	
H	$-H + H - H + O = O \rightarrow H - H + H - O \rightarrow H$	
	(i) Dut a single around each of the hands which are hydron when the security to	

(i) Put a circle around each of the bonds which are **broken** when the reaction takes place. [1]

(ii) Explain why the reaction of hydrogen with oxygen is exothermic.

Use ideas about bond breaking and bond making.

[Total: 12]

6

(a)	Electricity may be obtained using the sources listed.			
fossil fuels				
	geo	thermal		
	sola	ur		
	tida	l		
	wind	d		
	(i)	State which of the sources of energy is non-renewable.		
		[1	
(ii) State which two of the sources of energy are not dependent on the Sun.		State which two of the sources of energy are not dependent on the Sun.		
		and	1]	
(b)	Mar	ny types of power station use steam to turn a turbine attached to a generator.		
	-	lain, in terms of the forces and distances between molecules and the motion of molecule steam is able to fill its container.	S	
		[3]	

(c) Fig. 6.1 shows a diagram of a simple a.c. generator.




Fig. 6.1

Explain why the generator produces an a.c. output.	
	. [3]

(ii) On the grid provided in Fig. 6.2, sketch a graph of voltage output against time for this generator.

You must show at least one full cycle.




Fig. 6.2

[1]

[Total: 9]

15

BLANK PAGE

7 Fig. 7.1 is a diagram of a cross-section of a leaf.




Fig. 7.1

(a)	Identify the letter that represents the part in Fig. 7.1:		
	whe	ere most photosynthesis takes place	
	whi	ch transports water from roots to leaves	
	whi	ch controls gas exchange	[3]
(b)	Dra	w an X on Fig. 7.1 to identify a spongy mesophyll cell.	[1]
(c)	The	part labelled B in Fig. 7.1 is responsible for translocation.	
	(i)	Name the part labelled B .	
			[1]
	(ii)	Name the two main substances transported by the part labelled B .	
		1	
		2	
			[2]

(d) Table 7.1 compares the processes of transpiration and translocation.

Place ticks (\checkmark) in the boxes to show the correct features of transpiration and translocation.

Table 7.1

	transpiration	translocation
transports substances to regions of storage		
transports water		
movement of substances is in one direction only		
transport is from source to sink		

[2]

[Total: 9]

8 A scientist investigates the reaction between marble chips (calcium carbonate) and dilute hydrochloric acid.

The scientist uses 20 g of marble chips and 40 cm³ of dilute hydrochloric acid.

The temperature of the acid is 25 °C.

Fig. 8.1 shows the apparatus used.




Fig. 8.1

The scientist measures the mass every 20 seconds until the reaction stops.

The scientist calculates the loss in mass.

Fig. 8.2 shows the results.




Fig. 8.2

(a)	(i)	Use Fig. 8.2 to state the loss in mass after 40 seconds.	
		loss in mass = g [1]
	(ii)	Use Fig. 8.2 to state how long it takes for the reaction to stop.	
		time = seconds [1]
(b)	The	scientist does the experiment again.	
		s time the scientist uses dilute hydrochloric acid which is less concentrated than in the experiment.	те
	The	reaction is slower .	
	Exp	lain why reactions are slower when reactants are less concentrated.	
	Exp	lain your answer in terms of collisions between particles.	
		[2]
(c)	Car	bon dioxide gas is made in the reaction.	
	Sta	te the chemical test and its positive result for carbon dioxide gas.	
	test		
	resi	ultr	
(4)	In ti	nis experiment, 0.47 g of carbon dioxide gas is made.	[2]
(u)	III U	ils experiment, 0.47 g or carbon dioxide gas is made.	
	Cal	culate the volume occupied by 0.47 g of carbon dioxide gas.	
	The	molar gas volume at 25 °C is 24 dm ³ .	
	Sho	ow your working.	
	[<i>A</i> _r :	C, 12; O, 16]	

 $volume = \dots dm^3 [3]$

[Total: 9]

9	Visi	ble li	ght is a transverse wave and is part of the electromagnetic spectrum.
	(a)	Stat	te what is meant by a transverse wave.
	/L.\		[1]
	(a)	bloc	9.1 shows a ray of visible light from a torch (flashlight) shining into a rectangular glass ck.
			glass
			air
			Fig. 9.1
		(i)	Complete Fig. 9.1 to show the path the ray takes through and out of the block. [2]
		(ii)	State the term used to describe what happens to the ray of light as it enters the glass block.
			[1]
	((iii)	Explain why this happens to the ray of light.

(c) Fig. 9.2 shows the electrical circuit used in the torch.





Fig. 9.2

When the switch is closed, the current in the lamp is 1.8A and the potential difference across the lamp is 3.0 V.

Calculate the power output of the lamp.



(d) Fig. 9.3 shows two lamps, identical to the torch lamp, connected in parallel.




Fig. 9.3

(i) When the switch is closed, the ammeter reads 2.6A.

State the current in each lamp.

(ii) Calculate the quantity of charge passing through one of the lamps in Fig. 9.3 when it is switched on for 30 seconds.

Give the correct unit for your answer.

[Total: 11]

10 (a) The blood groups of some people are recorded.

Fig. 10.1 shows the results.




Fig. 10.1

(i)	State the total number of people who had their blood group recorded.	
		[1]
(ii)	Describe the evidence in Fig. 10.1 that shows blood group is an example of discontinuous variation.	ous
		[2]
(iii)	Complete this sentence about discontinuous variation.	
	Discontinuous variation is mostly caused by alone.	[1]
(iv)	Name one example of continuous variation in humans.	
		[1]

(b)	Blood	contains	four	main	components.
-----	-------	----------	------	------	-------------

The boxes on the left show the components of blood.

The boxes on the right show the functions of each component.

Draw lines to link each component with its function.

component		function
plasma		antibody production
	-	
platelet		blood clotting
	7	
red blood cell		transport of ions, soluble nutrients and hormones
	7	
white blood cell		transport of oxygen
		[3]
(c) Describe two wa	ays the structure of a r	red blood cell is adapted to its function.
1		
2		[2]
		[Total: 10]

	ium has the electronic structure 2.8.1. Sodium atoms can form sodium ions, Na $^+$. Chlorine atoms can form chloride ions, C l^- .	
(a)	Chlorine atoms can form chloride ions, Cl^- .	
	Describe, in terms of electrons, how a sodium atom forms a sodium ion and a chlorine ato forms a chloride ion.	m
	sodium atom	
	chlorine atom	
		 [1]
(b)	Sodium ions, Na ⁺ , form ionic bonds with chloride ions, C <i>l</i> ⁻ .	
	Describe how ionic bonds form between sodium ions and chloride ions.	
		2]
(c)	Solid aluminium chloride is another ionic compound.	
	Aluminium chloride contains aluminium ions, A l^{3+} , and chloride ions, C l^{-} .	
	Determine the formula of aluminium chloride.	
		[1]

(d) Fig. 11.1 shows the apparatus used to extract aluminium from aluminium ore.




Fig. 11.1

	Describe how aluminium is extracted from aluminium ore.
	Include the starting materials and the essential reaction conditions.
	[3]
(e)	Copper is extracted from copper ore by heating the copper ore with carbon.
	The equation for the reaction is shown.
	2CuO + C \rightarrow 2Cu + CO ₂
	Carbon is the reducing agent in this reaction.
	Define, in terms of electrons, what is meant by the term reducing agent.
	[1]
	[Total: 8]

12 Fig. 12.1 shows a speed-time graph for a train.




Fig. 12.1

(a) Use Fig. 12.1 to calculate the distance travelled by the train in the first 100 s.

(b) Use Fig. 12.1 to calculate the acceleration of the train from 0s to 100s.

acceleration =
$$m/s^2$$
 [2]

(c) Use Fig. 12.1 to describe the motion of the train from 100s to 200s.

(d) Fig. 12.2 shows the forces acting on the train when it is travelling at constant speed.




Fig. 12.2

(i)	State the magnitude of the force P .	
		[1]
(ii)	Calculate the mass of the train.	

The gravitational field strength on Earth, g, is 10 N/kg.

mass =	kg	[1]
The train is made of steel painted dark grey.		
On sunny days, the inside of the train can get very hot.		
Explain why painting the train white would reduce the heating effect.		
		[1]

[Total: 10]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.

© UCLES 2021 0654/42/F/M/21

(e)

The Periodic Table of Elements

		NIII	5 T	helium 4	10	Ne	neon 20	18	Ā	argon 40	36	궃	krypton 84	54	Xe	xenon 131	98	Ru	radon			
		III			6	ш	fluorine 19	17	Cl	chlorine 35.5	35	ğ	bromine 80	53	П	iodine 127	85	¥	astatine -			
		IN			80	0	oxygen 16	16	ഗ	sulfur 32	34	Se	selenium 79	52	<u>e</u>	tellurium 128	84	Ъо	polonium -	116		livermorium —
		^			7	z	nitrogen 14	15	۵	phosphorus 31	33	As	arsenic 75	51	Sb	antimony 122	83	Ξ	bismuth 209			
		//			9	ပ	carbon 12	14	S	silicon 28	32	Ge	germanium 73	20	S	tin 119	82	8 B	lead 207	114	Εl	flerovium -
		≡			5	Δ	boron 11	13	Αl	aluminium 27	31	Ga	gallium 70	49	In	indium 115	81	<i>1</i> L	thallium 204			
											30	Zu	zinc 65	48	В	cadmium 112	80	Нg	mercury 201	112	Ö	copernicium —
SILICITION											29	Cn	copper 64	47	Ag	silver 108	79	Αn	gold 197	111	Rg	roentgenium -
ne renouic lable of Elements	Group										28	ïZ	nickel 59	46	Pd	palladium 106	78	₹	platinum 195	110	Ds	darmstadtium -
ווסמוכ ומ	Gre				_						27	ပိ	cobalt 59	45	R	rhodium 103	11	ŀ	indium 192	109	¥	meitnerium -
ם בו			- ⊐	hydrogen 1							26	Ьe	iron 56	44	Ru	ruthenium 101	9/	SO	osmium 190	108	Ϋ́	hassium -
								_			25	M	manganese 55	43	ပ	technetium -	75	Re	rhenium 186	107	Bh	bohrium —
						pol	ass				24	ပ်	chromium 52	42	Mo	molybdenum 96	74	≯	tungsten 184	106	Sg	seaborgium -
				Key	atomic number	atomic symbo	name relative atomic mass				23	>	vanadium 51	41	g	niobium 93	73	<u>a</u>	tantalum 181	105	Ср	dubnium —
						atc	lei lei				22	j=	titanium 48	40	Zr	zirconium 91	72	Ξ	hafnium 178	104	Ŗ	rutherfordium —
											21	Sc	scandium 45	39	>	yttrium 89	57-71	lanthanoids		89–103	actinoids	
		=			4	Be	beryllium 9	12	Mg	magnesium 24	20	Ca	calcium 40	88	S	strontium 88	26	Ba	barium 137	88	Ra	radium -
		_			8	:=	lithium 7	11	Na	sodium 23	19	×	potassium 39	37	Rb	rubidium 85	55	S	caesium 133	87	ъ	francium -

	22	58	59	09	61	62	63	64	65	99	29	89	69		71
lanthanoids	Гa	Ce	Ą	PN	Pm	Sm	Eu	P9	Д	ò	웃	ш	E		P
	lanthanum 139	cerium 140	praseodymium 141	neodymium 144	promethium -	samarium 150	europium 152	gadolinium 157	terbium 159	dysprosium 163	holmium 165	erbium 167	thulium 169	ytterbium 173	lutetium 175
	88	06	91	92	93	94	92	96	26	86	66	100	101		103
actinoids	Ac	드	Ра	\supset	ď	Pu	Am	Cu	Ř	ŭ	Es	Fm	Md		۲
	actinium	thorium	protactinium	uranium	neptunium	plutonium	americium	curium	berkelium	californium	einsteinium	ferminm	mendelevium		lawrencium
	ı	232	231	238	ı	ı	ı	ı	ı	1	ı	ı	ı	ı	ı

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).