Please check the examination details b	elow before ente	ring your candidate information
Candidate surname		Other names
Ce	entre Number	Candidate Number
Pearson Edexcel		
Level 1/Level 2 GCSE (9–1)		
		10
Thursday 16 M	ay 20	19
Morning (Time: 1 hour 10 minutes)	Paper Re	eference 1SC0/1CH
Combined Science	ce	
Paper 2: Chemistry 1		
Taper Evenement,		
		Higher Tier
		Thigher ther
You must have:		Total Marks
Calculator, ruler		

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions.
- Answer the questions in the spaces provided
 there may be more space than you need.
- Calculators may be used.
- Any diagrams may NOT be accurately drawn, unless otherwise indicated.
- You must show all your working out with your answer clearly identified at the end of your solution.

Information

- The total mark for this paper is 60
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.
- In questions marked with an **asterisk** (*), marks will be awarded for your ability to structure your answer logically showing how the points that you make are related or follow on from each other where appropriate.
- There is a periodic table on the back cover of the paper.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







Answer ALL questions. Write your answers in the spaces provided.

Some questions must be answered with a cross in a box \boxtimes . If you change your mind about an answer, put a line through the box \boxtimes and then mark your new answer with a cross \boxtimes .

1 In Figure 1, the letters **A**, **E**, **G**, **J**, **X** and **Z** show the positions of six elements in the periodic table.

These letters are not the symbols of the atoms of these elements.

1	2						3	4	5	6	7	0
A							E			G		
J												X
				Z								

Figure 1

- (a) Using the letters A, E, G, J, X and Z
 - (i) give the letters of the **two** elements that are non-metals

(1)

(ii) give the letters of **two** elements in period 2

(1)

(iii) give the letter of an element that normally forms an ion with a charge of +1.

(1)

(b) Element **E** has an atomic number of 5. In a sample of **E** there are two isotopes. One isotope has a mass number of 10 and the other isotope has a mass number of 11.

(i) Explain, in terms of subatomic particles, what is meant by the term **isotopes**.

(2)

	/::\	٨Π	atoms of alamont E in this sample contain	
			atoms of element E in this sample contain	(1)
	X		5 protons	
	X	В	5 neutrons	
	X	C	6 protons	
	X	D	6 neutrons	
(c)	Ele	me	nt X has an atomic number of 18.	
	Sta	ite t	he electronic configuration of an atom of element X .	
				(1)
(d)	a c Cal	om Icula	experiment, 3.5 g of element A reacted with 4.0 g of element G to form bound. The ate the empirical formula of this compound. The atomic masses: $\mathbf{A} = 7$, $\mathbf{G} = 16$)	
			ust show your working.	
	101	u III	ast snow your working.	(3)
	••••••			
			empirical formula of this compound =	



2	(a)	Water, acidified with sulfuric acid, is decomposed by electrolysis.
		The water is decomposed to produce hydrogen and oxygen.

(i)	A sample of hydrogen	is mixed with ai	r and ignited.
-----	----------------------	------------------	----------------

State what would happen.

(1)

(ii) Throughout the experiment the volume of hydrogen and the volume of oxygen are measured at two-minute intervals.

The results are shown in Figure 2.

time in minutes	volume of hydrogen in cm³	volume of oxygen in cm³
0	0	0
2	4	2
4	8	4
6	12	6
8	16	8

Figure 2

Describe, using the data in Figure 2, what the results show about the volumes of hydrogen and of oxygen produced in this experiment.

r	á			'n	
		7)	-1	١
	4		_	п	

(b) N	Nolten lead bromide is electrolysed.	
Т	he products of this electrolysis are	(4)
× A	hydrogen and bromine	(1)
⊠ B	hydrogen and oxygen	
⊠ C	lead and bromine	
× D	lead and oxygen	
(c) C	Calcium nitrate and calcium carbonate are both ionic compounds.	
	Calcium nitrate mixed with water behaves as an electrolyte. Calcium carbonate mixed with water does not behave as an electrolyte.	
E	explain, in terms of solubility and movement of ions, this difference in behaviour.	(2)
(d) V	When molten zinc chloride is electrolysed, zinc ions, Zn ²⁺ , form zinc atoms.	
٧	Vrite the half equation for this reaction.	(2)



3	(a)	One way to extract metals from land contaminated with metal compounds is phytoextraction. When plants grow they absorb metal ions through their roots. The plants are harvested, dried and burned forming an ash. The ash contains metal compounds.	
		Plants were grown in a piece of ground contaminated with nickel compounds.	
		(i) 1 kg of the ash from these plants contained 142.0 g of nickel compounds.	
		Calculate the percentage by mass of nickel compounds in the ash.	(3)
		percentage by mass =	=
		(ii) Nickel is extracted from nickel compounds.	
		State an advantage of extracting nickel by phytoextraction rather than from its ore.	
			(1)
	(b)	Some nickel ores contain nickel sulfide.	
		(i) In the first stage of extracting nickel from nickel sulfide, the nickel sulfide, NiS, is heated in air to form nickel oxide, NiO, and sulfur dioxide.	
		Write the balanced equation for this reaction.	(2)

	(ii)		the final stage of the extraction process, a nickel compound is electrolysed produce pure nickel.	
		An	advantage of producing a metal by electrolysis is that	(1)
	X	A	electrolysis uses a large amount of electricity	
	×	В	the metal produced by electrolysis is very pure	
	X	C	electrolysis is a very cheap method of extraction	
	X	D	electrolysis is the only method of extracting unreactive metals	
(c)	The	uids e bo e bo	ferent method of obtaining nickel, the process produces a mixture of the nickel tetracarbonyl and iron pentacarbonyl. biling point of nickel tetracarbonyl is 43 °C. biling point of iron pentacarbonyl is 103 °C. two liquids mix together completely.	
			oe the process used to separate these two liquids.	
				(3)
			(Total for Question 3 = 10 mag	rks)



4 (a) Hydrated copper sulfate, CuSO₄.5H₂O, is a blue solid. Anhydrous copper sulfate, CuSO₄, is a white solid.

Heat energy is needed to convert hydrated copper sulfate to anhydrous copper sulfate. This is a reversible reaction.

$$CuSO_4.5H_2O \rightleftharpoons CuSO_4 + 5H_2O$$

Devise an experiment to show that this is a reversible reaction.

(4)

(b)	Hydrogen reacts with iodine to form hydrogen iodide.
	lodine gas is purple and hydrogen iodide gas is colourless.

$$H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$$

Hydrogen and iodine are placed in a sealed container. The container is left until equilibrium is reached.

The conditions are changed favouring the forward reaction.

Explain what you would see.

(2)

	(Total for Question 4 = 8 mai	rks)
	number of atoms =	
		(2)
(C)	Calculate the number of atoms combined in one mole of copper iodide, CuI_2 . (Avogadro constant = 6.02×10^{23})	
(6)	Calculate the number of stome combined in one male of conner indide Cul	

5 Pieces of zinc react with copper sulfate solution. Zinc sulfate solution is colourless.

$$Zn(s) + CuSO_4(aq) \rightarrow ZnSO_4(aq) + Cu(s)$$

(a) Describe what you would **see** when an excess of zinc is added to copper sulfate solution and the mixture left until the reaction is complete.

(2)

(b) This reaction is described as a redox reaction.

Explain, in terms of electrons, which particles have been oxidised and which particles have been reduced in this reaction.

(4)

(c) The copper sulfate solution used has a concentration of $15.95 \mathrm{g}\mathrm{dm}^{-3}$.	
Calculate the number of moles of copper sulfate, CuSO ₄ , in 50.00 cm ³ of this solution.	
(relative atomic masses: $O = 16$, $S = 32$, $Cu = 63.5$) (3)	
number of moles of copper sulfate =	mol
(d) In another experiment, 0.043 mol of copper sulfate, CuSO ₄ , is used.	
Calculate, to one decimal place, the minimum mass of zinc that must be added to react with all the copper sulfate. (relative atomic mass: $Zn = 65$)	
(relative atomic mass: $Z\Pi = 65$) (2)	
mass =	g
(Total for Question 5 = 11 marks)	



- **6** (a) **X** and **Y** are solutions of two different acids. The concentration of acid in each solution, in mol dm⁻³, is the same. Solution **X** has a pH of 3.40 and solution **Y** has a pH of 4.40.
 - (i) State what could be used to measure these pH values of 3.40 and 4.40.

(1)

(ii) What is the concentration of hydrogen ions in solution **X** compared with that in solution **Y**?

(1)

- A ten times lower
- **B** lower by a factor of 3.30/4.40
- ☑ C higher by a factor of 4.40/3.30
- **D** ten times higher

(b)) Ar ad	n experime ded to 50	ent is planned to record the change in pH as a powdered base is cm³ dilute hydrochloric acid.	
	Th	e method	l suggested is	
		step 1	add dilute hydrochloric acid up to the 50 cm ³ mark on a beaker	
		step 2	add one spatula of the base and stir	
		step 3	measure the pH of the mixture	
		step 4	repeat steps 2 and 3 until the pH stops changing.	
	(i)		w you could change the method so that the amounts of dilute	
		hydroch	loric acid and of the base can be measured more accurately.	(2)
diluta	hyd	Irochloric	acid	
dilute	TIYC	irocritoric	aciu	
base				
	(ii) During the experiment the pH changes from 2 to 10. If phenolphthalein indicator is added at the beginning of the experiment, a colour change occurs as the base is added.State the colour change that occurs.			
		colour a	t start	
		colour a	t end	
	(iii) Explain, the expe	in terms of the particles present, why the pH increases during	
				(2)



*(c) Some properties of four solids, **A**, **B**, **C** and **D**, are shown in Figure 3.

The solids, in no particular order, are copper carbonate, copper oxide, magnesium metal and sodium hydroxide.

	Α	В	С	D
colour of solid	black	silver	white	green
observation when solid is added to water	black solid remains	a few bubbles appear on surface of solid	solid dissolves and forms colourless solution	green solid remains
pH of mixture of solid added to water	7	8	13	7
observation when solid is added to dilute sulfuric acid	on warming, solid disappears to form blue solution	effervescence solid disappears to form colourless solution	solid disappears to form colourless solution	effervescence solid disappears to form blue solution

Figure 3

Identify the solids A , B , C and D , explaining how the information in Figure 3 supports the identification of each solid.	(4)	
	(6)	

4		
\times		
S S		
H H S		
$\times \times \times \times$		
Z		
XXXXXX		
$\times\!\!\!\circ\!\!\times$		
Ž		
0		
Δ		
THIS AREA		
V)		
0		
\times		

\times		
XXXXX		
A H		
×		
4		
88 <u>88</u> 88		
5		
Ž		
(XXXXX)		

(Total for Question 6 = 13 marks)



The Periodic Table of the Elements

0	4 He helium 2	20 Ne neon 10	40 Ar argon 18	84 Kr krypton 36	131 Xe xenon 54	[222] Rn radon 86	fully
_		19 F fluorine 9	35.5 CI chlorine 17	80 Br bromine 35	127 	[210] At astatine 85	orted but not
9		16 O oxygen 8	32 S sulfur 16	79 Se selenium 34	128 Te tellurium 52	[209] Po polonium 84	ve been repo
2		14 N nitrogen 7	31 P phosphorus 15	75 As arsenic 33	122 Sb antimony 51	209 Bi bismuth 83	s 112-116 ha authenticated
4		12 C carbon 6	28 Si silicon 14	73 Ge germanium 32	119 Sn tin 50	207 Pb	mic numbers a
က		11 B boron 5	27 AI aluminium 13	70 Ga gallium 31	115 In indium 49	204 T thallium 81	Elements with atomic numbers 112-116 have been reported but not fully authenticated
	!			65 Zn zinc 30	112 Cd cadmium 48	201 Hg mercury 80	Elem
				63.5 Cu copper 29	108 Ag silver 47	197 Au gold 79	[272] Rg roentgenium
				59 Ni nickel 28	106 Pd palladium 46	195 Pt platinum 78	Ds damstadtium
				59 Co cobalt 27	103 Rh rhodium 45	192 Ir iridium 77	[268] Mt meitnerium 109
	T hydrogen			56 Fe iron 26	101 Ru ruthenium 44	190 Os osmium 76	[277] Hs hassium 108
_				55 Mn manganese 25	[98] Tc technetium 43	186 Re rhenium 75	[264] Bh bohrium 107
		nass ool umber		52 Cr chromium 24	96 Mo molybdenum 42	184 W tungsten 74	[266] Sg seaborgium 106
	Key	relative atomic mass atomic symbol name atomic (proton) number		51 V vanadium 23	93 Nb niobium 41	181 Ta tantalum 73	[262] Db dubnium 105
		relativ ato atomic		48 Ti titanium 22	91 Zr zirconium 40	178 Hf hafinium 72	[261] Rf rutherfordium 104
	'			45 Sc scandium 21	89 Y yttrium 39	139 La * lanthanum 57	[227] Ac* actinium 89
2		9 Be beryllium 4	24 Mg magnesium 12	40 Ca calcium 20	88 Sr strontium 38	137 Ba barium 56	[226] Ra radium 88
_		7 Li Ilthium 3	23 Na sodium 11	39 K potassium 19	85 Rb rubidium 37	133 Cs caesium 55	[223] Fr francium 87

^{*} The lanthanoids (atomic numbers 58-71) and the actinoids (atomic numbers 90-103) have been omitted.

The relative atomic masses of copper and chlorine have not been rounded to the nearest whole number.