

1. Fig. 11 shows the curve with parametric equations

$$x = 2\cos\theta, \quad y = \sin\theta, \quad 0 \leq \theta \leq 2\pi.$$

The point P has parameter $\frac{1}{4}\pi$. The tangent at P to the curve meets the axes at A and B.




Fig.11

- (a) Show that the equation of the line AB is $x + 2y = 2\sqrt{2}$. [6]

- (b) Determine the area of the triangle AOB. [3]

2. In this question you must show detailed reasoning.

A curve has parametric equations

$$x = \cos t - 3t \text{ and } y = 3t - 4 \cos t - \sin 2t, \text{ for } 0 \leq t \leq \pi.$$

Show that the gradient of the curve is always negative.

[7]

3. In this question you must show detailed reasoning.

Fig. 8 shows the curve with parametric equations

$$x = 7 \cos \theta + 2 \cos 2\theta, \quad y = 2 + \sin \theta, \quad (0 \leq \theta \leq 2\pi).$$




Fig. 8

- (a) Find the coordinates of the point on the curve with the greatest y -coordinate. [4]
- (b) Determine the exact y -coordinates of the points where the curve crosses the y -axis. [6]

- 4.

A curve has parametric equations $x = \frac{t}{1+t^3}$, $y = \frac{t^2}{1+t^3}$ where $t \neq -1$.

- (a) In this question you must show detailed reasoning.

Determine the gradient of the curve at the point where $t = 1$. [5]

- (b) Verify that the cartesian equation of the curve is $x^2 + y^3 = xy$. [3]

5. Fig. 9 shows the curve with parametric equations

$$x = \tan \theta, y = 1 + \cos 2\theta, \text{ for } 0 \leq \theta \leq \pi \text{ with } \theta \neq \frac{1}{2}\pi.$$




Fig. 9

- (a) Explain why $\theta \neq \frac{1}{2}\pi$ [1]

- (b) Show that the maximum value of y for points on the curve is 2. [1]

- (c) Show that the cartesian equation of the curve is $y = \frac{2}{1+x^2}$. [3]

- (d) In this question you must show detailed reasoning.

The point P in the first quadrant lies on the curve. Find the coordinates of P given that OP is the normal to the curve at P.

[7]

6. A curve has parametric equations

$$x = 2 \cot \theta, \quad y = 2 \sin^2 \theta,$$

for values of θ for which both x and y are defined.

- (a) For what values of θ , in the interval $0 \leq \theta \leq 2\pi$, is x undefined? [2]

- (b)

Show that the cartesian equation of the curve is $y = \frac{A}{x^2 + B}$ where A and B are positive integers to be determined.

[3]

- (c) Ali says that the curve lies completely above the x -axis. Determine whether Ali is correct. [2]

END OF QUESTION paper

			Total	9	Parametric Equations
2		<p>DR</p> $\frac{dx}{dt} = -\sin t - 3$ $\frac{dy}{dt} = 3 + 4\sin t - 2\cos 2t$ $\frac{dy}{dx} = \frac{3 + 4\sin t - 2\cos 2t}{-\sin t - 3}$ <p>denominator is always negative since $-1 \leq \sin t \leq 1$</p> <p>Substitution of $\cos 2t = 1 - 2\sin^2 t$ in numerator $4\sin^2 t + 4\sin t + 1$ seen in numerator</p> <p>Numerator is $(2\sin t + 1)^2$ so numerator is always positive and denominator is always negative, $\frac{dy}{dx}$ is always negative</p>	<p>B1(AO 3.1a)</p> <p>B1(AO 1.1)</p> <p>M1(AO 2.1)</p> <p>E1(AO 2.4)</p> <p>M1(AO 1.1)</p> <p>A1(AO 1.1)</p> <p>E1(AO 2.4)</p> <p>[7]</p>		
			Total	7	
3	a	<p>DR</p> $\frac{dy}{dx} = 0 \text{ when } \frac{dy}{d\theta} = \cos \theta = 0$ $\theta = \frac{1}{2}\pi, \frac{3}{2}\pi$	<p>M1(AO3.1a)</p> <p>M1(AO1.1)</p>		

A1(AO1.1)

A1(AO3.2a)

M1

A1

M1

A1

[4]

$$x = -2$$

$$y = 3$$

Alternative solution

Maximum y occurs when $\sin \theta = 1$

$$y = 3$$

$$\theta = \frac{1}{2}\pi$$

$$x = -2$$

DR

$$7 \cos \theta + 2 \cos 2\theta = 0 \Rightarrow 4 \cos^2 \theta + 7 \cos \theta - 2 = 0$$

$$(4 \cos \theta - 1)(\cos \theta + 2) = 0$$

b $\cos \theta = \frac{1}{4}$ or -2

Reject $\cos \theta = -2$

$$\sin^2 \theta = 1 - \cos^2 \theta = 1 - \left(\frac{1}{4}\right)^2 = \frac{15}{16}$$

M1(AO1.1a)

M1(AO1.1)

A1(AO1.1)

B1(AO3.2a)

M1(AO3.1a)

A1(AO2.2a)

Use of double angle formula

Method for solving quadratic

May be seen later

$$\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dt}{dx} = \frac{2t - t^4}{1 - 2t^3}$$


$$t = 1 \Rightarrow \frac{dy}{dx} = -1$$

Parametric Equations

earlier M1

Examiner's Comments

This part saw many attempts at the correct process, but there were some mistakes seen in applying the quotient rule generally with confusion over signs. More difficulty was experienced by those who chose to use the product rule but did not appreciate the need to apply the chain rule to the differentiation of $(1 + t^3)^{-1}$.



Afl

Possibly the use of the product rule to prove the quotient rule leads some candidates to assume that that is the method to use to differentiate quotients. Candidates should know the difference and be able to apply both the product rule and the quotient rule accurately as appropriate.

b

$\text{LHS} = \frac{t^3 + t^6}{(1 + t^3)^3}$	oe
--	----

$$= \frac{t^3(1 + t^3)}{(1 + t^3)^3}$$

M1 (AO 1.1a)

M1(AO 1.1)

A1 (AO 2.1)

[3]

AG
Expression for LHS

Factorising

Completion of argument

$$\text{RHS} = \frac{t^3}{(1+t^3)^2} = \text{LHS}$$

Examiner's Comments

This part was generally started well, but many candidates struggled to complete the argument to score fully.

The majority tackled this question by starting with finding an expression for $x^2 + y^2$ in terms of t , but a few chose to use the expression for x or y and make t the subject then continue to eliminate t from the expressions and generally complete the argument satisfactorily.

Exemplar 4

$$x = \frac{t}{1+t^3} \quad y = \frac{t^2}{1+t^3}$$

$$\frac{1}{y} = \frac{1+t^3}{t^2}$$

$$\frac{x}{y} = \frac{t}{1+t^3} \times \frac{1+t^3}{t^2}$$

$$= \frac{t}{t^2} = t^{-1} = \frac{1}{t}$$

$$t = \frac{y}{x}$$

$$x = \frac{y}{x} \frac{1}{1 + \left(\frac{y}{x}\right)^3} = \frac{y}{x} \left(1 + \left(\frac{y}{x}\right)^3\right) = \frac{y}{x}$$

$$x + \frac{xy^3}{x^3} = \frac{y}{x}$$

$$x^4 + xy^3 = x^2y$$

$$\text{divide by } x \quad \cancel{x^3} + y^3 = \cancel{x}y$$

$$x^3 + y^3 = xy$$

This candidate does full proof of the result gaining all 3 marks.

$$\left(\frac{t}{1+t^3}\right)^3 + \left(\frac{t^2}{1+t^3}\right)^3 = \frac{t^3}{(1+t^3)^3} + \frac{t^6}{(1+t^3)^3}$$

$$= \frac{t^6 + t^3}{(1+t^3)^3} = \frac{t^3(t^3+1)}{(1+t^3)^3} = \frac{t^3}{(1+t^3)^2} = xy$$

Because $\frac{t}{1+t^3} \cdot \frac{t^2}{1+t^3} = \frac{t^3}{(1+t^3)^2}$

$$\frac{dx}{dt} = (1+t^3)^{-1} - 3t^3(1+t^3)^{-2}$$

$$\frac{dy}{dt} = 2t(1+t^3)^{-1} - 3t^2(1+t^3)^{-2}$$

~~at~~ $t=1, \frac{dx}{dt} = -\frac{1}{4}$

$$\frac{dy}{dt} = \frac{1}{4}$$

$$\frac{dy}{dx} = \frac{1}{4} \div -\frac{1}{4} = -1$$

This candidate shows appropriate initial working to verify the equation, but misses out on the final mark, as they do not state what they have shown.

Total			8	Parametric Equations					
5	a	x is not defined for this value	E1 (AO 2.4) [1]	<table border="1"> <tr> <td style="width: 20px; height: 20px;"></td> <td style="width: 20px; height: 20px;"></td> </tr> </table>					
	b	cos 2θ has max value 1 when θ = 0 so max y is 2	E1 (AO 2.4) [1]	<table border="1"> <tr> <td style="width: 100px; height: 40px;">Or using $\frac{dy}{d\theta} = 0$</td> <td style="width: 100px; height: 40px;"></td> </tr> <tr> <td style="width: 100px; height: 20px;">correctly</td> <td style="width: 100px; height: 20px;"></td> </tr> </table>		Or using $\frac{dy}{d\theta} = 0$		correctly	
Or using $\frac{dy}{d\theta} = 0$									
correctly									
	c	$y = 2 \cos^2 \theta$ $1 + x^2 = \sec^2 \theta$ So $1 + x^2 = \frac{2}{y} \Rightarrow y = \frac{2}{1 + x^2}$	M1 (AO 2.2a) M1 (AO 1.1a) E1 (AO 1.1) [3]	Relevant use of $\cos 2\theta = 2 \cos^2 \theta - 1$ Relevant use of $\sec 2\theta = 1 + \tan^2 \theta$ AG Successful completion	At any stage At any stage With use of $\sec \theta = \frac{1}{\cos \theta}$				

DR

$$\frac{dy}{dx} = \frac{-4x}{(1+x^2)^2}$$

$$= \frac{y}{x}$$

Gradient OP

$$\frac{-4x}{(1+x^2)^2} \times \frac{y}{x} = -1$$

$$\frac{4x}{(1+x^2)^2} \times \frac{2}{x(1+x^2)} = 1 \Rightarrow (1+x^2)^3 = 8 \Rightarrow 1+x^2 = 2$$

$$x = 1$$

Point P is (1, 1)

M1 (AO
1.1)

A1 (AO 1.1)

B1 (AO
3.1a)M1 (AO
2.2a)M1 (AO
2.1)A1 (AO
2.2a)A1 (AO
3.2a)

[7]

Use of chain rule or quotient rule

Correct derivative

May be implied

Form and solve equation for x Allow ± 1 at this stage

Or, using parameters:

$$\frac{dy}{dx} = \frac{dy}{d\theta} / \frac{dx}{d\theta} = \frac{-2 \sin 2\theta}{\sec^2 \theta}$$

$$\frac{-2 \sin 2\theta (1 + \cos 2\theta)}{\sec^2 \theta \tan \theta} = -1$$

$$\frac{4 \sin \theta \cos^3 \theta}{1} \times \frac{2 \cos^3 \theta}{\sin \theta} = 1$$

$$\cos \theta = (\pm) \frac{1}{\sqrt{2}}$$

Total			12	Parametric Equations
6	a	$0, 2\pi$ π	B1(AO 1.2) B1(AO 1.2) [2]	
	b	<div style="border: 1px solid black; padding: 5px; display: inline-block; margin-bottom: 10px;"> $\operatorname{cosec}^2 \theta = 1 + \cot^2 \theta$, so $\frac{2}{y} = 1 + \left(\frac{x}{2}\right)^2$ </div> $\frac{2}{y} = \frac{4 + x^2}{4}$ $y = \frac{8}{x^2 + 4}$ <p>Alternative solution 1</p> $x^2 = 4 \cot^2 \theta = \frac{4 \cos^2 \theta}{\sin^2 \theta}$ $x^2 = \frac{2(2 - y)}{\frac{1}{2}y}$ $y = \frac{8}{x^2 + 4}$ <p>Alternative solution 2</p> $\frac{A}{x^2 + B} = \frac{A}{4 \cot^2 \theta + B} = 2 \sin^2 \theta$	M1(AO 3.1a) M1(AO 1.1) A1(AO 2.2a) M1 M1 A1 M1 M1	

				Parametric Equations	
		$8 \cos^2 \theta + 2B \sin^2 \theta = A$ <div style="border: 1px solid black; padding: 5px; display: flex; justify-content: space-between; align-items: center;"> <div style="flex: 1;">so $B = 4$ and $A = 8$, giving</div> <div style="flex: 1; text-align: center;"> $y = \frac{8}{x^2 + 4}$ </div> </div>	A1		
	c	<div style="border: 1px solid black; padding: 5px; display: flex; justify-content: space-between; align-items: center;"> <div style="flex: 1; text-align: center;"> $\frac{8}{x^2 + 4} > 0$ </div> <div style="flex: 1;">for all x</div> </div> <p>So Ali is correct [because $y > 0$ for all x]</p>	M1(AO 2.1) A1(AO 2.3)	<div style="border: 1px solid black; padding: 5px; display: flex; justify-content: space-between; align-items: center;"> <div style="flex: 1;">or $2\sin^2 \theta \geq 0$ and when $2\sin^2 \theta = 0$, x is undefined</div> <div style="flex: 1;">If zero scored, SC1 for $x^2 + 4$ is never negative oe</div> </div>	[2]
		Total	7		