

**Section A: Pure Mathematics**

**1** Consider the equations

$$\begin{aligned}ax - y - z &= 3, \\2ax - y - 3z &= 7, \\3ax - y - 5z &= b,\end{aligned}$$

where  $a$  and  $b$  are given constants.

- (i) In the case  $a = 0$ , show that the equations have a solution if and only if  $b = 11$ .
- (ii) In the case  $a \neq 0$  and  $b = 11$  show that the equations have a solution with  $z = \lambda$  for any given number  $\lambda$ .
- (iii) In the case  $a = 2$  and  $b = 11$  find the solution for which  $x^2 + y^2 + z^2$  is least.
- (iv) Find a value for  $a$  for which there is a solution such that  $x > 10^6$  and  $y^2 + z^2 < 1$ .

**2** Write down a value of  $\theta$  in the interval  $\pi/4 < \theta < \pi/2$  that satisfies the equation

$$4 \cos \theta + 2\sqrt{3} \sin \theta = 5.$$

Hence, or otherwise, show that

$$\pi = 3 \arccos(5/\sqrt{28}) + 3 \arctan(\sqrt{3}/2).$$

Show that

$$\pi = 4 \arcsin(7\sqrt{2}/10) - 4 \arctan(3/4).$$

**3** Prove that the cube root of any irrational number is an irrational number.

Let  $u_n = 5^{1/(3^n)}$ . Given that  $\sqrt[3]{5}$  is an irrational number, prove by induction that  $u_n$  is an irrational number for every positive integer  $n$ .

Hence, or otherwise, give an example of an infinite sequence of irrational numbers which converges to a given integer  $m$ .

[An irrational number is a number that cannot be expressed as the ratio of two integers.]

- 4 The line  $y = d$ , where  $d > 0$ , intersects the circle  $x^2 + y^2 = R^2$  at  $G$  and  $H$ . Show that the area of the minor segment  $GH$  is equal to

$$R^2 \arccos\left(\frac{d}{R}\right) - d\sqrt{R^2 - d^2}. \quad (*)$$

In the following cases, the given line intersects the given circle. Determine how, in each case, the expression (\*) should be modified to give the area of the minor segment.

- (i) Line:  $y = c$ ; circle:  $(x - a)^2 + (y - b)^2 = R^2$ .
- (ii) Line:  $y = mx + c$ ; circle:  $x^2 + y^2 = R^2$ .
- (iii) Line:  $y = mx + c$ ; circle:  $(x - a)^2 + (y - b)^2 = R^2$ .
- 5 The position vectors of the points  $A$ ,  $B$  and  $P$  with respect to an origin  $O$  are  $a\mathbf{i}$ ,  $b\mathbf{j}$  and  $l\mathbf{i} + m\mathbf{j} + n\mathbf{k}$ , respectively, where  $a$ ,  $b$ , and  $n$  are all non-zero. The points  $E$ ,  $F$ ,  $G$  and  $H$  are the midpoints of  $OA$ ,  $BP$ ,  $OB$  and  $AP$ , respectively. Show that the lines  $EF$  and  $GH$  intersect.

Let  $D$  be the point with position vector  $d\mathbf{k}$ , where  $d$  is non-zero, and let  $S$  be the point of intersection of  $EF$  and  $GH$ . The point  $T$  is such that the mid-point of  $DT$  is  $S$ . Find the position vector of  $T$  and hence find  $d$  in terms of  $n$  if  $T$  lies in the plane  $OAB$ .

- 6 The function  $f$  is defined by

$$f(x) = |x - 1|,$$

where the domain is  $\mathbf{R}$ , the set of all real numbers. The function  $g_n = f^n$ , with domain  $\mathbf{R}$ , so for example  $g_3(x) = f(f(f(x)))$ . In separate diagrams, sketch graphs of  $g_1$ ,  $g_2$ ,  $g_3$  and  $g_4$ .

The function  $h$  is defined by

$$h(x) = \left| \sin \frac{\pi x}{2} \right|,$$

where the domain is  $\mathbf{R}$ . Show that if  $n$  is even,

$$\int_0^n (h(x) - g_n(x)) \, dx = \frac{2n}{\pi} - \frac{n}{2}.$$

- 7 Show that, if  $n > 0$ , then

$$\int_{e^{1/n}}^{\infty} \frac{\ln x}{x^{n+1}} dx = \frac{2}{n^2 e}.$$

You may assume that  $\frac{\ln x}{x} \rightarrow 0$  as  $x \rightarrow \infty$ .

Explain why, if  $1 < a < b$ , then

$$\int_b^{\infty} \frac{\ln x}{x^{n+1}} dx < \int_a^{\infty} \frac{\ln x}{x^{n+1}} dx.$$

Deduce that

$$\sum_{n=1}^N \frac{1}{n^2} < \frac{e}{2} \int_{e^{1/N}}^{\infty} \left( \frac{1 - x^{-N}}{x^2 - x} \right) \ln x dx,$$

where  $N$  is any integer greater than 1.

- 8 It is given that  $y$  satisfies

$$\frac{dy}{dt} + k \left( \frac{t^2 - 3t + 2}{t + 1} \right) y = 0,$$

where  $k$  is a constant, and  $y = A$  when  $t = 0$ , where  $A$  is a positive constant. Find  $y$  in terms of  $t$ ,  $k$  and  $A$ .

Show that  $y$  has two stationary values whose ratio is  $(3/2)^{6k} e^{-5k/2}$ .

Describe the behaviour of  $y$  as  $t \rightarrow +\infty$  for the case where  $k > 0$  and for the case where  $k < 0$ .

In separate diagrams, sketch the graph of  $y$  for  $t > 0$  for each of these cases.

## Section B: Mechanics

- 9  $AB$  is a uniform rod of weight  $W$ . The point  $C$  on  $AB$  is such that  $AC > CB$ . The rod is in contact with a rough horizontal floor at  $A$  and with a cylinder at  $C$ . The cylinder is fixed to the floor with its axis horizontal. The rod makes an angle  $\alpha$  with the horizontal and lies in a vertical plane perpendicular to the axis of the cylinder. The coefficient of friction between the rod and the floor is  $\tan \lambda_1$  and the coefficient of friction between the rod and the cylinder is  $\tan \lambda_2$ .

Show that if friction is limiting both at  $A$  and at  $C$ , and  $\alpha \neq \lambda_2 - \lambda_1$ , then the frictional force acting on the rod at  $A$  has magnitude

$$\frac{W \sin \lambda_1 \sin(\alpha - \lambda_2)}{\sin(\alpha + \lambda_1 - \lambda_2)}.$$

- 10 A bead  $B$  of mass  $m$  can slide along a rough horizontal wire. A light inextensible string of length  $2\ell$  has one end attached to a fixed point  $A$  of the wire and the other to  $B$ . A particle  $P$  of mass  $3m$  is attached to the mid-point of the string and  $B$  is held at a distance  $\ell$  from  $A$ . The bead is released from rest.

Let  $a_1$  and  $a_2$  be the magnitudes of the horizontal and vertical components of the initial acceleration of  $P$ . Show by considering the motion of  $P$  relative to  $A$ , or otherwise, that  $a_1 = \sqrt{3}a_2$ . Show also that the magnitude of the initial acceleration of  $B$  is  $2a_1$ .

Given that the frictional force opposing the motion of  $B$  is equal to  $(\sqrt{3}/6)R$ , where  $R$  is the normal reaction between  $B$  and the wire, show that the magnitude of the initial acceleration of  $P$  is  $g/18$ .

- 11 A particle  $P_1$  is projected with speed  $V$  at an angle of elevation  $\alpha$  ( $> 45^\circ$ ), from a point in a horizontal plane. Find  $T_1$ , the flight time of  $P_1$ , in terms of  $\alpha, V$  and  $g$ . Show that the time after projection at which the direction of motion of  $P_1$  first makes an angle of  $45^\circ$  with the horizontal is  $\frac{1}{2}(1 - \cot \alpha)T_1$ .

A particle  $P_2$  is projected under the same conditions. When the direction of the motion of  $P_2$  first makes an angle of  $45^\circ$  with the horizontal, the speed of  $P_2$  is instantaneously doubled. If  $T_2$  is the total flight time of  $P_2$ , show that

$$\frac{2T_2}{T_1} = 1 + \cot \alpha + \sqrt{1 + 3 \cot^2 \alpha}.$$

## SECTION C: Probability and Statistics

- 12** The life of a certain species of elementary particles can be described as follows. Each particle has a life time of  $T$  seconds, after which it disintegrates into  $X$  particles of the same species, where  $X$  is a random variable with binomial distribution  $B(2, p)$ . A population of these particles starts with the creation of a single such particle at  $t = 0$ . Let  $X_n$  be the number of particles in existence in the time interval  $nT < t < (n + 1)T$ , where  $n = 1, 2, \dots$

Show that  $P(X_1 = 2 \text{ and } X_2 = 2) = 6p^4q^2$ , where  $q = 1 - p$ . Find the possible values of  $p$  if it is known that  $P(X_1 = 2 | X_2 = 2) = 9/25$ .

Explain briefly why  $E(X_n) = 2pE(X_{n-1})$  and hence determine  $E(X_n)$  in terms of  $p$ . Show that for one of the values of  $p$  found above  $\lim_{n \rightarrow \infty} E(X_n) = 0$  and that for the other  $\lim_{n \rightarrow \infty} E(X_n) = +\infty$ .

- 13** The random variable  $X$  takes the values  $k = 1, 2, 3, \dots$ , and has probability distribution

$$P(X = k) = A \frac{\lambda^k e^{-\lambda}}{k!},$$

where  $\lambda$  is a positive constant. Show that  $A = (1 - e^{-\lambda})^{-1}$ . Find the mean  $\mu$  in terms of  $\lambda$  and show that

$$\text{Var}(X) = \mu(1 - \mu + \lambda).$$

Deduce that  $\lambda < \mu < 1 + \lambda$ .

Use a normal approximation to find the value of  $P(X = \lambda)$  in the case where  $\lambda = 100$ , giving your answer to 2 decimal places.

- 14** The probability of throwing a 6 with a biased die is  $p$ . It is known that  $p$  is equal to one or other of the numbers  $A$  and  $B$  where  $0 < A < B < 1$ . Accordingly the following statistical test of the hypothesis  $H_0 : p = B$  against the alternative hypothesis  $H_1 : p = A$  is performed.

The die is thrown repeatedly until a 6 is obtained. Then if  $X$  is the total number of throws,  $H_0$  is accepted if  $X \leq M$ , where  $M$  is a given positive integer; otherwise  $H_1$  is accepted. Let  $\alpha$  be the probability that  $H_1$  is accepted if  $H_0$  is true, and let  $\beta$  be the probability that  $H_0$  is accepted if  $H_1$  is true.

Show that  $\beta = 1 - \alpha^K$ , where  $K$  is independent of  $M$  and is to be determined in terms of  $A$  and  $B$ . Sketch the graph of  $\beta$  against  $\alpha$ .