

OCR (A) Biology A-level

Topic 5.2: Energy for biological processes


Notes



Photosynthesis

Photosynthesis is a reaction in which **light energy** is used to split apart the strong bonds in water molecules in a process of **photolysis** in order to combine **hydrogen** with **carbon dioxide** to produce a fuel in the form of **glucose**. **Oxygen** is a waste product of this reaction and is released into the atmosphere. The rate of photosynthesis is determined by carbon dioxide concentration, light intensity and well as temperature.

The **chloroplast** is the site of photosynthesis and it is adapted to photosynthesise in the following ways:



Chloroplast diagram

Figure 1 Biology TutorVista

- It contains **stacks of thylakoid membranes called grana** which contain the photosynthetic pigments such as **chlorophyll arranged as photosystems**
- It contains **stroma** which is the fluid surrounding the grana, stroma contains all the **enzymes** required for the light independent stage of photosynthesis

There are two stages of photosynthesis:

- **Light-dependent reaction** in which **electrons are excited** to a higher energy level by the energy trapped by **chlorophyll** molecules in the **thylakoid membranes**. Electrons are then passed down the **electron transport chain** from one electron carrier to the next and this process generates **ATP from ADP and inorganic phosphate** in a process called **photophosphorylation**. **Reduced NADP** is also generated in the light-dependent stage as the electrons are transferred to NADP (NADPH) along with a proton. Both ATP and reduced NADP are then used in the light-independent stage of photosynthesis.
- **Light-independent reaction** also known as the **Calvin cycle** is the final stage of photosynthesis which uses ATP (source of energy) and reduced NADP (reducing power) to produce glucose. Light independent reaction occurs as following:
 - 1) **RuBP** is combined with **carbon dioxide** in a reaction called **carbon fixation catalysed by the enzyme RUBISCO**.
 - 2) **RuBP** is converted into **two glycerate 3-phosphate (GP)** molecules
 - 3) Reduced NADP and ATP are used to convert **GP to triose phosphate (TP)**
 - 4) Some of TP molecules are used to make **glucose** which is then converted to essential organic compounds such as **polysaccharides, lipids, amino acids and nucleic acids**.
 - 5) Remaining TP molecules are used to **reform RuBP with the help of ATP**.



Factors affecting photosynthesis

Photosynthesis is affected by many factors, the factor in lowest supply and therefore limiting the rate of the reaction is known as the **limiting factor**.

Limiting factors include:


- **Light intensity** – if this is in short supply the light dependent reaction will slow therefore there will be lower amounts of ATP and NADPH created. This will then affect the Calvin cycle as these are needed to convert GP to TP. So the level of GP will rise and TP will fall which in turn causes RuBP levels to fall.
- **CO₂ concentration** – if this is in short supply the light independent reaction will slow
- **Temperature** – if this is low Rubisco and other molecules will have lower levels of kinetic energy therefore the enzyme-controlled reactions are affected.

Respiration

Aerobic respiration is splitting of the **respiratory substrate**, to release carbon dioxide as a waste product and reuniting of hydrogen with atmospheric oxygen with the release of a large amount of energy whereas **anaerobic respiration** occurs in the absence of oxygen.

Respiration is a multi-step process with each step controlled and catalysed by a **specific** intracellular enzyme.

Glycolysis is the first process of both aerobic and anaerobic respiration and it occurs in cytoplasm.



In this process a molecule of glucose is **phosphorylated** to produce 2 molecules of **pyruvate**, 2 molecules of NADH and a **net** production of 2 molecules of ATP.




Figure 2 IB guides


The next step of aerobic reaction is **the link reaction** where each pyruvate is converted to **acetyl** which **binds to coenzyme A**. In the process **NAD is reduced to NADH** and **CO₂ is produced** as **pyruvate is decarboxylated**.

Co-enzyme A delivers Acetyl to the **Krebs cycle** where glucose is oxidised and carbon dioxide, ATP, **reduced NAD** and **reduced FAD** are produced. Each glucose molecule



causes the cycle to turn twice so **per glucose we produce 4CO₂, 4NADH, 2FADH and 2ATP molecules in the Krebs cycle.** The ATP molecules produced by the Kreb's cycle are done so by **substrate level phosphorylation**

Oxidative phosphorylation



©1999 Addison Wesley Longman, Inc.

Oxidative phosphorylation is the process in which ATP is synthesised in the **electron transport chain** in mitochondria. This process generates the majority of ATP in aerobic respiration and it occurs as following:

- Reduced coenzymes (NADH and FADH) carry **hydrogen ions** and electrons to the electron transport chain which occurs on the **inner mitochondrial membrane**
- Electrons are carried from one electron carrier to another in a series of **redox reactions: the electron carrier** which passes the electron on is oxidised whereas the electron carrier which receives it is reduced

The energy provided by the electrons to the electron carriers is used to move **Hydrogen ions** across the inner membrane into the **intermembrane space** – as a result of that the concentration of the hydrogen ions in the intermembrane space is high

The inner membrane is impermeable to H⁺

- Hydrogen ions diffuse into the **mitochondrial matrix** down the **electrochemical gradient through the ATPase enzyme.**
- ATP is produced on **stalked particle** using ATP synthase
- Hydrogen atoms are produced from hydrogen ions and electrons. The **hydrogen atoms are then combined with oxygen to produce water**
- Oxygen acts as the **final electron acceptor**



There is a theoretical yield of 38 ATP molecules per glucose molecule but in real life this is rarely achieved due to the inner mitochondrial membrane being 'leaky' to H^+ therefore not all H^+ move through the ATPase. The pyruvate made during glycolysis in cytoplasm needs moving into the matrix by active transport and so this uses ATP.

Respiratory substrates include **carbohydrates, lipids and proteins** which release varying amounts of energy, depending on the number of hydrogens in the structure which are oxidised to water. For instance, the number of hydrogens is greater in fatty acids than carbohydrates.

The **respiratory quotient (RQ)** can be measured to determine which respiratory substrate is being used and to determine if the organism is undergoing anaerobic respiration.

$RQ = \text{carbon dioxide produced} / \text{oxygen consumed}$

Different respiratory substrates have different RQ values e.g. carbohydrates have a value of 1.0, lipids – 0.8 and proteins 0.9.

Anaerobic Respiration

This occurs when the concentration of oxygen is low. ATP production still needs to happen but this can't be done by oxidative phosphorylation due to the lack of oxygen to act as the final electron acceptor.

In order for some ATP to be produced anaerobic respiration allows glycolysis to continue. (Glycolysis has a net production of 2 ATP per glucose molecule).

In mammals pyruvate is converted to **lactate**. **Pyruvate acts as the hydrogen acceptor** to enable **NADH to be reoxidised to NAD** which can then be **used to continue the reactions in glycolysis**.

Lactate can then be converted back to pyruvate in the liver cells when the oxygen levels rise again.

Yeast and **plants** use **alcoholic fermentation** to enable glycolysis to continue. In this process Pyruvate is decarboxylated to **Ethanol** which in turn is reduced to **ethanol** reoxidising NAD in the process. So **ethanol is the hydrogen acceptor**. The first step in this process produces CO_2 and therefore this is an **irreversible** reaction.

