

WJEC Chemistry AS-level

1.2: Basic Ideas about Atoms

Detailed Notes English Specification

This work by PMT Education is licensed under CC BY-NC-ND 4.0







Atomic Structure

The model for atomic structure has **evolved over time** as knowledge and scientific understanding have changed.

Plum Pudding Model

It was initially thought that atoms consisted of a **sphere of positive charge**, with **small negative charges distributed** evenly within it.

Electron Shell Model

It is now known that the atom consists of a **small, dense central nucleus** surrounded by **orbiting electrons** in electron **shells**. This was discovered in the Rutherford scattering experiment in 1911, when alpha particles were deflected by a central, positively charged body.

The nucleus consists of **protons and neutrons** giving it an overall **positive charge**. It contains almost the entire mass of the atom. In a neutral atom, the number of electrons is **equal to** the number of protons due to the relative charges.



/wiki/File:Atom_Diagram.svg) AG Caesar / CC BY-SA 3.0

Radioactivity

Not all nuclei of elements are **stable**. This means the **ratio** of protons and neutrons are **imbalanced**. In order to try and regain this balance, **radiation is emitted**, releasing excess protons, neutrons or both. Throughout these particle interactions, **neutrinos and antineutrinos** can be released in order to **conserve energy** and other **quantum values**.

Alpha Decay

Alpha decay is a type of radioactive decay, during which an atomic nucleus **loses two protons** and **two neutrons**. An alpha particle is equivalent to a helium nucleus with two protons and two neutrons. It reduces the atomic number by two and the mass number by four, making the element **more stable**.

Example:



www.pmt.education





Beta Decay

A beta particle, equivalent to an **electron**, is released, **reducing the proton number** by one. It is considered to have **zero mass**, so the mass number is not affected.

Example:



Beta⁺ Decay

A beta plus particle is the **antiparticle to an electron**, meaning it has the **same mass** but **opposite charge**. Therefore, when released, it increases a nucleus' proton number by one and has no effect on mass number.

Example:



Electron Capture

This particle interaction occurs specifically within the nucleus. A proton interacts with an electron, producing a neutron. Therefore mass number remains the same and proton number is reduced by one.

Example:



Gamma Radiation

Gamma radiation is a type of **electromagnetic radiation** rather than a particle. It has a very **short wavelength** and **high frequency**. Its wave nature means it is able to travel long distances.





Radiation Properties

Alpha, Beta and Gamma radiations vary in penetrating power and can be affected by electric and magnetic fields.

	Range in Air	Ionising Power	Penetrating Strength
Alpha	< 5 cm	very strong	stopped by thin paper
Beta	< 1 metre	medium	stopped by thin metal
Gamma	infinite	weak	stopped by thick lead

Half Life

Radioactivity of a substance decreases over time as it gains a more stable composition. Half life is the time it takes for the radioactivity or the number of nuclei present to halve. This time period remains the same throughout the decay of the substance meaning radiation levels in the future can be easily predicted.

Example:



(https://commons.wikimedia.org/wiki/File:Radioactive_decay.png) Kurt Rosenkrantz / CC BY-SA 3.0





Ionisation Energies

Standard Molar Ionisation Energies

Ionisation energy is defined as:

The minimum energy required to remove one mole of electrons from one mole of atoms in a gaseous state. It is measured in kJmol⁻¹.



Successive Ionisation Energies

Successive ionisation energies occur when **further electrons are removed**. This usually requires **more energy** because, as electrons are removed, the **electrostatic force of attraction** between the positive nucleus and the negative outer electron **increases**. More energy is therefore needed to overcome this attraction so **ionisation energy increases**.

First ionisation energy follows trends within the Periodic Table:

Along a Period - first ionisation energy **increases** due to a decreasing atomic radius and greater electrostatic forces of attraction (due to the increasing number of protons).

Down a Group - first ionisation energy **decreases** due to an increasing atomic radius and increasing shielding which reduces the effect of the electrostatic forces of attraction.

When successive ionisation energies are plotted on a graph, a **sudden large increase** indicates a **change in energy level**. This is because the electron is being removed from an orbital closer to the nucleus so more energy is required to do so.







This large energy increase provides supporting evidence for the atomic orbital theory.

The first ionisation energy of aluminium is **lower than expected** due to **the outer electron in the 3p orbital being at a higher energy than the 3s electrons**. Therefore, it takes less energy than expected to remove the outer electron as we move from s block to p block.

Shapes of Electron Orbitals

Electrons are held in **clouds of negative charge** called **orbitals**. There are different types of orbital: *s*, *p*, *d* and *f*. Each one has a different shape. *Example:*



Electron Transition Spectra

When an electron moves from a higher energy level to a lower energy level, a **photon is released** to help **conserve energy** and other quantum values. Likewise, when it gains energy to move up an energy level, **photons are absorbed** to provide the energy for this.

The specific **wavelengths and energies of photons** that are involved in the interactions can be seen using emission and absorption spectra.

Emission Spectra - This displays lines at the specific frequencies of **emitted** photons from when an electron moves **down an energy level**.





Absorption Spectra - This displays an entire spectrum with **black lines** for the 'missing' frequencies of photons that have been **absorbed** when an electron moves **up** an energy level.

Example:



Energy, Wavelength & Frequency

These three quantities are linked in a series of relationships. Energy is **proportional** to frequency with **Planck's constant** (6.63x10⁻³⁴ m²kgs⁻¹). *Example:*

E = hf where E = energy (J), h = Planck's constant (J s), f = frequency of light absorbed (Hz)

Frequency, and therefore energy, is **inversely proportional** to wavelength, with the **speed of light in a vacuum** as the constant (3x10⁸ ms⁻¹).

Example:



These relationships show that the **higher the frequency** of radiation, the **higher its energy**. Likewise, the **higher its wavelength** the **lower its energy**.

Applying this to the EM spectrum, **gamma** radiation has a high frequency and short wavelength and therefore has the **highest energy**. In comparison, **radio waves** have the longest wavelength and lowest frequency and therefore have the **lowest energy**.

🕟 www.pmt.education





Example:



(https://commons.wikimedia.org/wiki/File:EM_Spectrum3-new.jpg) NASA / CC BY-SA 3.0

www.pmt.education

0

 \odot

▶ Image: PMTEducation

