



GCE

Chemistry A

Advanced GCE

Unit F325: Equilibria, Energetics and Elements

Mark Scheme for June 2011

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2011

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
1 (a)	(The enthalpy change that accompanies) the formation of one mole of a(n ionic) compound ✓ from its gaseous ions ✓ (under standard conditions)	2	<p>IGNORE 'Energy needed' OR 'energy required'</p> <p>ALLOW as alternative for compound: lattice, crystal, substance, solid</p> <p>Note: 1st mark requires 1 mole 2nd mark requires gaseous ions</p> <p>IF candidate response has '1 mole of gaseous ions', award 2nd mark but NOT 1st mark</p> <p>IGNORE: $\text{Mg}^{2+}(\text{g}) + 2\text{Cl}^{-}(\text{g}) \longrightarrow \text{MgCl}_2(\text{s})$ (question asks for words)</p>
(b)	(i) Hydration involves bond forming OR bonds are made ✓	1	<p>ALLOW statement of any type of bond being formed ALLOW (chloride) ions attract water (molecules)</p> <p>ALLOW a response in terms of hydrogen bonds breaking AND bond making</p> <p>DO NOT ALLOW response stating that energy is required DO NOT ALLOW response that refers to ions in H_2O, eg H^+</p>
	(ii)	2	<p>Correct species AND state symbols required for both marks Mark each marking point independently</p> <p>ALLOW response on upper line: $\text{Mg}^{2+}(\text{g}) + 2\text{Cl}^{-}(\text{aq})$ (ie Cl^{-} hydrated before Mg^{2+})</p> <p>ALLOW $\text{MgCl}_2(\text{aq})$</p>

F325

Mark Scheme

June 2011

Question			Answer	Mark	Guidance
1	(b)	(iii)	<p>FIRST, CHECK THE ANSWER ON ANSWER LINE IF answer = $-1921 \text{ (kJ mol}^{-1}\text{)}$ award 2 marks</p> <p>-----</p> $(-2493) + (-154) = (2 \times -363) + \Delta H_{\text{hyd}}(\text{Mg}^{2+}) \checkmark$ $\Delta H_{\text{hyd}}(\text{Mg}^{2+}) = (-2493) + (-154) - (2 \times -363)$ $= -1921 \text{ (kJ mol}^{-1}\text{)} \checkmark$	2	<p>IF there is an alternative answer, check to see if there is any ECF credit possible using working below. See list below for marking of answers from common errors</p> <p>-----</p> <p>ALLOW for 1 mark:</p> <p>-2284 use of Cl^- rather than $2 \times \text{Cl}^-$ (+1921 signs all reversed OR lack of 2 for 363 -1613 sign wrong for 154 (+3065 sign wrong for 2493 -3373 sign wrong for 2×363</p>
	(c)		<p>Magnesium ion OR Mg^{2+} is smaller OR Mg^{2+} has greater charge density \checkmark</p> <p>Mg^{2+} has a stronger attraction to H_2O OR Mg^{2+} has a stronger bonding with H_2O \checkmark</p>	2	<p>ORA: Calcium ion OR Ca^{2+} is larger OR Ca^{2+} has smaller charge density</p> <p>IGNORE idea of close packing of ions IGNORE 'atomic' and 'atoms' and assume that Mg or Ca refer to ions, ie ALLOW Mg has a smaller (atomic) radius</p> <p>ALLOW Mg has a stronger attraction to H_2O ORA: e.g. Ca^{2+} has less attraction to H_2O</p> <p>DO NOT ALLOW Mg atoms have a stronger attraction to H_2O</p> <p>DO NOT ALLOW stronger attraction/bonding between ions Note: Response must refer to attraction/bonding with H_2O or this must be implied from the whole response</p>
			Total	9	

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
2 (a)	<p>Temperature: (Forward) reaction is exothermic OR gives out heat OR reverse reaction is endothermic OR takes in heat ✓</p> <p>Pressure: Right-hand side has fewer number of (gaseous) moles ✓ ORA</p> <p>Equilibrium Lower temperature/cooling AND increasing pressure shifts (equilibrium position) to the right ✓</p>	3	<p>ANNOTATE WITH TICKS AND CROSSES, etc</p> <p>ALLOW K_c increases at lower temperatures</p> <p>3rd mark is for stating that BOTH low temperature and high pressure shift equilibrium to the right (Could be separate statements)</p> <p>Note: ALLOW suitable alternatives for 'to right', e.g.: towards NO_2 OR towards products OR in forward direction OR increases yield of NO_2/products</p> <p>ALLOW 'favours the right', as alternative for 'shifts equilibrium to right'</p> <p>IGNORE responses in terms of rate</p>
(b)	<p>$4\text{NH}_3 + 5\text{O}_2 \longrightarrow 4\text{NO} + 6\text{H}_2\text{O}$ ✓</p> <p>$2\text{NO}_2 + \text{H}_2\text{O} \longrightarrow \text{HNO}_3 + \text{HNO}_2$ ✓</p>	2	<p>ALLOW multiples, e.g. $2\text{NH}_3 + 2\frac{1}{2}\text{O}_2 \longrightarrow 2\text{NO} + 3\text{H}_2\text{O}$</p> <p>ALLOW \rightleftharpoons OR \rightarrow in equations</p>
(c) (i)	<p>$(K_c =) \frac{[\text{NO}_2]^2}{[\text{NO}]^2 [\text{O}_2]}$ ✓</p>	1	<p>Square brackets are essential</p>

F325

Mark Scheme


June 2011

Question	Answer	Mark	Guidance
2 (c) (ii)	<p>FIRST, CHECK THE ANSWER ON ANSWER LINE IF answer = 45 dm³ mol⁻¹, award 5 marks IF answer = 45 with incorrect units, award 4 marks</p> <p>-----</p> <p>Equilibrium moles 0.60 mol NO₂ ✓</p> <p>0.20 mol NO AND 0.40 mol O₂ ✓</p> <p>Equilibrium concentrations (equilibrium moles ÷ 2) [NO₂] = 0.30 mol dm⁻³ AND [NO] = 0.10 mol dm⁻³ AND [O₂] = 0.20 mol dm⁻³ ✓</p> <p>Calculation of K_c and units $K_c = \frac{0.30^2}{0.10^2 \times 0.20} = 45 \text{ ✓ dm}^3 \text{ mol}^{-1} \text{ ✓}$</p>	5	<p>IF there is an alternative answer, check to see if there is any ECF credit possible using working below</p> <p>-----</p> <p>ANNOTATE WITH TICKS AND CROSSES, etc ALLOW ECF throughout</p> <p><i>Alternative route if concs NO and O₂ calculated at start:</i></p> <p>initial concentrations: 0.40 mol dm⁻³ NO AND 0.35 mol dm⁻³ O₂ ✓</p> <p>Equilibrium concentrations: [NO₂] = 0.30 mol dm⁻³ ✓ [NO] = 0.10 mol dm⁻³ AND [O₂] = 0.20 mol dm⁻³ ✓</p> <p>For units, ALLOW mol⁻¹ dm³</p> <p>ALLOW ECF using any incorrect values for concentrations OR moles of NO, O₂ AND NO₂ For ECF, ALLOW 2 significant figures up to calculator value correctly rounded ALLOW ECF from incorrect K_c expression for both calculation and units</p> <p>Common ECFs worth less than 5 marks: 22.5 not ÷2 3 marks + unit mark 1.61 0.6 for NO₂ but 0.8 for NO and 0.7 for O₂ No mark for moles NO and O₂ 3 marks + unit mark 0.804 As above but also no ÷2 No mark for moles NO and O₂ AND ÷2 2 marks + unit mark</p>
	Total	11	

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
3	<p>Evidence of at least two half-lives measured on graph OR within text (would need evidence of two half-lives) ✓</p> <p>Any half-life value stated in range 180–220 s OR constant half-life ✓</p> <p>1st order ✓ Note: This is only correct response for order (ie no ECF). If not stated separately, this mark can be awarded from a rate equation, e.g. $rate = k[Br_2]^1$ OR $rate = k[Br_2]$</p> <p>Evidence of tangent on graph drawn to line at $t = 0$ s ✓ e.g.</p> 	4	<p>ANNOTATE ALL Q3 WITH TICKS AND CROSSES, etc</p> <p>MARK ON GRAPH OR IN TEXT</p> <p>LOOK FOR STATEMENT ON GRAPH OR WITHIN TEXT ALLOW almost constant half-life</p> <p>-----</p> <p>Note: Response may use an alternative approach from half-life for the 1st two marks based on gradients of tangents: 1st mark would be awarded for evidence of two tangents drawn on graph 2nd mark would be awarded for stating that ratio of concentrations = ratio of rates, e.g. gradient of tangent at $0.010 \text{ mol dm}^{-3}$ has twice the value of gradient of tangent at $0.005 \text{ mol dm}^{-3}$</p> <p>-----</p> <p>MARK TANGENTS ON GRAPH ALLOW some leeway but tangent must coincide with part of curve that is 'straight' (ie between $[Br_2] = 0.010$–0.009 and MUST NOT cross the curve</p>

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
3	$\text{rate} = \frac{0.010}{250} = 0.000040 \text{ OR } 4.0 \times 10^{-5} \checkmark$ <p>units: $\text{mol dm}^{-3} \text{ s}^{-1} \checkmark$</p>	2	<p>ALLOW values from 1 SF (0.00004 OR 4×10^{-5}) up to calculator value, correctly rounded</p> <p>ALLOW range $\sim \frac{0.010}{160}$ to $\frac{0.010}{300}$:</p> <p>i.e. ALLOW a calculated gradient in the range $6 \times 10^{-5} - 3 \times 10^{-5}$ from a tangent drawn at $t = 0$</p> <p>IF tangent is drawn on graph at a different time or incorrectly (e.g. crossing curve), then mark rate calculation by ECF using the gradient of the tangent drawn by the candidate (<i>ie</i> not the range above).</p> <p>IF no tangent is drawn ALLOW a value in the range above ONLY</p> <p>Credit only attempts at tangents, not just a random straight line</p> <p>IGNORE a ‘– sign’</p>
	$\text{rate} = k[\text{Br}_2] \text{ OR } k = \frac{\text{rate}}{[\text{Br}_2]} \checkmark$ <p>$k =$ calculated result from $\frac{\text{calculated value for rate}}{0.010} \checkmark$</p> <p>units: $\text{s}^{-1} \checkmark$</p>	3	<p>DO NOT ALLOW $\text{rate} = k[\text{Br}]$, <i>ie</i> Br instead of Br_2</p> <p>DO NOT ALLOW just $k[\text{Br}_2]$, <i>ie</i> ‘rate =’ OR ‘r =’ must be present</p> <p>Calculation of k is from candidate’s calculated initial rate</p> <p>From 0.00004, $k = \frac{0.000040}{0.010} = 4 \times 10^{-3} \text{ s}$</p> <p>Note:</p> <p>IF order with respect to Br_2 has been shown as 2nd order, then mark this part by ECF, e.g. if Br_2 shown to be 2nd order, $\text{rate} = k[\text{Br}_2]^2$</p> <p>$k =$ calculated result from $\frac{\text{calculated value for rate}}{0.010^2}$</p> <p>units: $\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ OR $\text{mol}^{-1} \text{ dm}^3 \text{ s}^{-1}$</p> <p>Note: Units mark must correspond to the candidate’s stated rate equation, NOT an incorrectly rearranged k expression</p>
	Total	9	

F325

Mark Scheme

June 2011

Question			Answer	Mark	Guidance
4	(a)	(i)	proton donor ✓	1	ALLOW H ⁺ donor
		(ii)	(the proportion of) dissociation ✓ Correct equation for any of the four acids: C ₆ H ₅ COOH ⇌ H ⁺ + C ₆ H ₅ COO ⁻ OR CH ₃ COOH ⇌ H ⁺ + CH ₃ COO ⁻ OR CH ₃ COCOOH ⇌ H ⁺ + CH ₃ COCOO ⁻ OR CH ₃ CHOHCOOH ⇌ H ⁺ + CH ₃ CHOHCOO ⁻ ✓	2	ALLOW a weak acid partly dissociates ALLOW a strong acid totally dissociates ALLOW ionisation for dissociation ALLOW the ability to donate a proton Equilibrium sign required ALLOW equilibria involving H ₂ O and H ₃ O ⁺ e.g. C ₆ H ₅ COOH + H ₂ O ⇌ H ₃ O ⁺ + C ₆ H ₅ COO ⁻ , etc DO NOT ALLOW HA ⇌ H ⁺ + A ⁻
		(iii)	weakest: CH ₃ COOH acetic acid C ₆ H ₅ COOH benzoic acid CH ₃ CHOHCOOH lactic acid strongest: CH ₃ COCOOH ✓ pyruvic acid	1	ALLOW correct order using any identifier from the table, <i>ie</i> , common name, systematic name, structural formula OR pK _a value
		(iv)	C ₆ H ₅ COOH ₂ ⁺ + CH ₃ CHOHCOO ⁻ ✓	1	BOTH products AND correct charges required for mark Mark ECF from incorrect order in (iii) See response from (iii) below response to (iv)

F325

Mark Scheme




June 2011

Question	Answer	Mark	Guidance
4 (b) (i)	$2\text{CH}_3\text{COCOOH} + \text{Ca}(\text{OH})_2 \rightarrow (\text{CH}_3\text{COCOO})_2\text{Ca} + 2\text{H}_2\text{O} \checkmark$ <p>Note: pyruvic acid must have been used here and formula of pyruvic acid and pyruvate must be correct</p>	1	All species AND balancing required for the mark ALLOW $(\text{CH}_3\text{COCOO}^-)_2\text{Ca}^{2+}$ ALLOW equation showing $2\text{CH}_3\text{COCOO}^- + \text{Ca}^{2+}$ IF charges shown, charges must balance, e.g. DO NOT ALLOW $(\text{CH}_3\text{COCOO}^-)_2\text{Ca}$ IGNORE state symbols if shown ALLOW multiples ALLOW equilibrium sign
	(ii) $\text{H}^+ + \text{OH}^- \longrightarrow \text{H}_2\text{O}$	1	ALLOW multiples but not same species on both sides ALLOW equilibrium sign IGNORE state symbols if shown ALLOW $\text{H}_3\text{O}^+ + \text{OH}^- \longrightarrow 2\text{H}_2\text{O}$ ALLOW $\text{CH}_3\text{COCOOH} + \text{OH}^- \longrightarrow \text{CH}_3\text{COCOO}^- + \text{H}_2\text{O}$
(c)	<p>FIRST, CHECK THE ANSWER ON ANSWER LINE IF answer = 2.11, award 4 marks</p> <p>-----</p> $K_a = 10^{-\text{p}K_a}$ $= 10^{-2.39} \text{ OR } 0.00407 \checkmark$ $K_a = \frac{[\text{H}^+][\text{CH}_3\text{COCOO}^-]}{[\text{CH}_3\text{COCOOH}]} \text{ (ALLOW use of HA, H}^+ \text{ and A}^-)$ <p>OR $[\text{H}^+] = \sqrt{K_a \times [\text{HA}]}$</p> <p>OR $[\text{H}^+] = \sqrt{0.00407 \times 0.0150} \checkmark$ (subsumes 1st marking point) $[\text{H}^+] = 0.00782 \text{ (mol dm}^{-3}) \checkmark$</p> <p>$\text{pH} = -\log 0.00782 = 2.11 \checkmark$</p>	4	<p>IF there is an alternative answer, check to see if there is any ECF credit possible using working below</p> <p>-----</p> <p>IF ECF, ANNOTATE WITH TICKS AND CROSSES, etc</p> <p>ALLOW 0.0041 to calculator value: 0.004073802</p> <p>IF the $\text{p}K_a$ of a different weak acid has been used use ECF from 2nd marking point</p> <p>ALLOW 0.0078 to calculator value (depending on previous rounding) ALLOW ONLY 2.11 (This is to take into account poor previous rounding)</p> <p>IF candidate has used $0.0150 \text{ mol dm}^{-3}$ (<i>ie</i> assumes strong acid) ALLOW final mark ONLY by ECF for a pH of 1.82</p> <p>IF no square root used, pH = 4.21 3 marks</p>

F325

Mark Scheme

June 2011

Question			Answer	Mark	Guidance
4	(d)	(i)		1	<p>ALLOW correct structural OR displayed OR skeletal formula OR recognisable mixture of formulae</p> <p>DO NOT ALLOW molecular formula but ALLOW (COOH)₂ OR (CO₂H)₂</p>  <p>ALLOW OH  BUT not O-H-C</p>
		(ii)	$\text{C}_2\text{H}_2\text{O}_4 \rightleftharpoons \text{H}^+ + \text{C}_2\text{HO}_4^- \checkmark$ $\text{C}_2\text{HO}_4^- \rightleftharpoons \text{H}^+ + \text{C}_2\text{O}_4^{2-} \checkmark$	2	<p>ALLOW in either order</p> <p>ALLOW arrow instead of equilibrium sign</p> <p>ALLOW molecular formulae for this part</p> <p>ALLOW equilibria involving H₂O and H₃O⁺</p> <p>ALLOW equations using structures</p>

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
4 (e)	<p>Chemicals (1 mark) lactic acid / CH₃CHOHCOOH AND (sodium) lactate / CH₃CHOHCOO⁻ (Na⁺) ✓</p> <p>Concentrations (4 marks)</p> <p>EITHER [H⁺(aq)] = 10^{-3.55} OR 2.8 x 10⁻⁴ OR 2.82 x 10⁻⁴ (mol dm⁻³) ✓ <i>separate marking point</i></p> <p>$K_a = 10^{-3.86}$ OR 1.4 x 10⁻⁴ OR 1.38 x 10⁻⁴ (mol dm⁻³) ✓ <i>separate marking point</i></p> <p>$\frac{[HA]}{[A^-]} = \frac{[H^+]}{K_a}$ OR $\frac{[A^-]}{[HA]} = \frac{K_a}{[H^+]}$ ✓</p> <p>$\frac{[HA]}{[A^-]} = \frac{2.8 \times 10^{-4}}{1.4 \times 10^{-4}}$ OR $\frac{2}{1}$ OR 2 OR $\frac{[A^-]}{[HA]} = \frac{0.5}{1}$ OR 0.5 ✓</p> <p>This marking point subsumes previous marking point ONLY</p> <p>Comment (1 mark) Magic tang/taste could come from other chemicals/substances in the sweet OR The buffer would have the same taste/tang as the magic tang ✓</p>	6	<p>ANNOTATE WITH TICKS AND CROSSES, etc</p> <p>ALLOW any lactate salt ALLOW lactic acid AND NaOH OR lactic acid AND OH⁻</p> <p>-----</p> <p>FOR ALTERNATIVE using Henderson–Hasselbalch equation, SEE PAGE 11</p> <p>-----</p> <p>If another weak acid has been selected and salt has been selected, allow ECF for remainder of question SEE PAGE 12</p> <p>-----</p> <p>ALLOW 2.8 x 10⁻⁴ up to calculator value of 2.81838 x 10⁻⁴ ALLOW 0.00028, etc</p> <p>ALLOW 1.4 x 10⁻⁴ up to calculator value of 1.38038 x 10⁻⁴ ALLOW 0.00014, etc</p> <p>ALLOW use of CH₃CHOHCOOH AND CH₃CHOHCOO⁻(Na⁺) ALLOW use of acid AND salt</p> <p>ALLOW value from $\frac{\text{calculated value of } [H^+]}{\text{calculated value of } K_a}$</p> <p>ALLOW 2SF up to calculator value of 2.041742129 correctly rounded but ALLOW 2 if 2.8 x 10⁻⁴ and 1.4 x 10⁻⁴ used ALLOW 2 mol dm⁻³ HA AND 1 mol dm⁻³ A⁻ OR any concentration ratio of 2(acid) : 1(salt)</p> <p>ALLOW 2SF up to calculator value of 0.489778819 correctly rounded but ALLOW 0.5 if 2.8 x 10⁻⁴ and 1.4 x 10⁻⁴ used</p>

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
	<p>ALTERNATIVE approach for concentrations using Henderson–Hasselbalch equation (4 marks)</p> $\text{pH} = \text{p}K_{\text{a}} + \log \frac{[\text{A}^{-}]}{[\text{HA}]} \quad \text{OR} \quad -\log K_{\text{a}} + \log \frac{[\text{A}^{-}]}{[\text{HA}]} \quad \checkmark$ $\log \frac{[\text{A}^{-}]}{[\text{HA}]} = 3.55 - 3.86 \quad \checkmark \quad (\text{subsumes previous mark})$ $\log \frac{[\text{A}^{-}]}{[\text{HA}]} = -0.31 \quad \checkmark \quad (\text{subsumes previous mark})$ $\frac{[\text{A}^{-}]}{[\text{HA}]} = 10^{-0.31} = \frac{0.490}{1} \quad \text{OR} \quad 0.490 \quad \checkmark$		<p>ALLOW use of $\text{CH}_3\text{CHOHCOOH}$ AND $\text{CH}_3\text{CHOHCOO}^{-}(\text{Na}^{+})$ ALLOW use of acid AND salt ALLOW $\text{pH} = \text{p}K_{\text{a}} - \log \frac{[\text{HA}]}{[\text{A}^{-}]}$ OR $-\log K_{\text{a}} - \log \frac{[\text{HA}]}{[\text{A}^{-}]}$ ALLOW $\log \frac{[\text{HA}]}{[\text{A}^{-}]} = 3.86 - 3.55$ (subsumes previous mark) ALLOW $\log \frac{[\text{HA}]}{[\text{A}^{-}]} = 0.31$ (subsumes previous mark) ALLOW $\frac{[\text{HA}]}{[\text{A}^{-}]} = 10^{0.31} = \frac{2.04}{1}$ OR $\frac{2}{1}$ OR 2 For $\frac{[\text{A}^{-}]}{[\text{HA}]}$, ALLOW 2 SF up to calculator value of 0.48978819 For $\frac{[\text{HA}]}{[\text{A}^{-}]}$, ALLOW 2 SF up to calculator value of 2.041737945 but ALLOW 2 if $10^{-0.31}$ used</p>

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
4 (e)	<p>SUMMARY OF 4(e) MARKING POINTS FOR EACH POSSIBLE ACID CHOSEN FIRST, CHECK THE ANSWER ON ANSWER LINE: IF answer is correct for weak acid chosen, award MP2–MP5 IF there is an alternative answer, check to see if there is any ECF credit possible using working below</p>		
	lactic	pyruvic	acetic
pK_a	3.86	2.39	4.76
MP1	lactic AND lactate OR lactic acid AND OH^-	No mark	No mark
MP2: $[\text{H}^+]$	$10^{-3.55}$ OR 2.82×10^{-4} (calc: 2.81838×10^{-4})		
MP3: K_a	$10^{-3.86}$ OR 1.38×10^{-4} calc: $1.380384265 \times 10^{-4}$	$10^{-2.39}$ OR 4.07×10^{-3} calc: $4.073802778 \times 10^{-3}$	$10^{-4.76}$ OR 1.74×10^{-5} calc: $1.737800829 \times 10^{-5}$
MP4: ratio expression	$\frac{[\text{HA}]}{[\text{A}^-]} = \frac{[\text{H}^+]}{K_a}$ OR $\frac{[\text{A}^-]}{[\text{HA}]} = \frac{K_a}{[\text{H}^+]}$		
MP5: $\frac{[\text{HA}]}{[\text{A}^-]}$	$\frac{2.82 \times 10^{-4}}{1.38 \times 10^{-4}}$ OR 2.04 calc: 2.041737945	$\frac{2.82 \times 10^{-4}}{4.07 \times 10^{-3}}$ OR 0.0693 calc: 0.069183097	$\frac{2.82 \times 10^{-4}}{1.74 \times 10^{-5}}$ OR 16.2 calc: 16.21810097
OR $\frac{[\text{A}^-]}{[\text{HA}]}$	$\frac{1.38 \times 10^{-4}}{2.82 \times 10^{-4}}$ OR 0.489 calc: 0.489778819	$\frac{4.07 \times 10^{-3}}{2.82 \times 10^{-4}}$ OR 14.4 calc: 14.45439771	$\frac{1.74 \times 10^{-5}}{2.82 \times 10^{-4}}$ OR 0.0617 calc: 0.0616595
<p>TAKE CARE: Calc values are completely unrounded and may differ between brands of calculator Use actual candidate values at each stage using rounding to 2 or more SF. MP5: calculated using 3 SF from MP2 and MP3 calc values for MP5 are completely unrounded (using calculator values from MP2 and MP3) Be slightly flexible as candidates may have written down rounded values but carried on with calculator values – This approach is ACCEPTABLE</p>			
Total		20	

F325

Mark Scheme



June 2011

Question		Answer	Mark	Guidance
5	(c) (ii)	Two from points below: 1. fuel OR fuel cells 2. manufacture of margarine OR hydrogenation of alkenes/unsaturated fats 3. manufacture of ammonia OR 'Haber process' ✓ 4. manufacture of HCl/hydrochloric acid 5. reduction of metal ores/metal oxides	1	2 uses for one mark IGNORE hydrogenation of margarine
	(d)	FIRST, CHECK THE ANSWER ON ANSWER LINE IF answer = -109, award first 3 marks for calculation ----- At 298 K, $91.2 = 176 - T\Delta S$ ✓ $\Delta S (= \frac{176 - 91.2}{298}) = 0.285 \text{ (kJ K}^{-1} \text{ mol}^{-1}\text{)}$ OR $\Delta S (= \frac{176000 - 91200}{298}) = 285 \text{ (J K}^{-1} \text{ mol}^{-1}\text{)}$ ✓ <i>subsumes 1st marking point</i> At 1000 K, $\Delta G = 176 - 1000 \times 0.285$ $= -109 \text{ (kJ mol}^{-1}\text{)}$ ✓ Reaction does take place (spontaneously) because $\Delta G < 0$ OR ΔG is -ve ✓ Note: If no value of ΔG , this mark cannot be awarded.	4	IF there is an alternative answer, check to see if there is any ECF credit possible using working below ----- ANNOTATE WITH TICKS AND CROSSES, etc ALLOW 0.285 (3 SF) up to calculator value of 0.284563758 ALLOW 285 (3 SF) up to calculator value of 284.563758 ALLOW -109 up to calculator value correctly rounded, i.e. -108.6, -108.56, etc ALLOW ECF from incorrect ΔS , ie calculated value of ΔG from $\Delta G = 176 - 1000 \times$ calculated value of ΔS Answer and reason BOTH needed for mark ALLOW reaction is feasible for 'reaction does take place' Note: If candidate has a + ΔG value, mark ECF , ie reaction does not take place because $\Delta G > 0$ OR ΔG is +ve
Total			11	

F325

Mark Scheme


June 2011

Question	Answer	Mark	Guidance
6 (a)	<p>Ni $1s^2 2s^2 2p^6 3s^2 3p^6 3d^8 4s^2$ ✓</p> <p>d block: (Ni:) 'd' is highest energy sub-shell/orbital ✓</p> <p>Ni^{2+}: $1s^2 2s^2 2p^6 3s^2 3p^6 3d^8$ ✓</p> <p>Transition element: has an ion with an incomplete/partially-filled d sub-shell/orbital ✓</p> <p>-----</p> <p>A ligand donates an electron pair to Ni^{2+} OR metal ion OR metal ✓</p> <p>A complex ion is an ion bonded to ligand(s)/surrounded by ligands ✓</p> <p>Coordinate bond/dative covalent mentioned at least once in the right context ✓</p>	<p>4</p> <p>3</p>	<p>ANNOTATE WITH TICKS AND CROSSES, etc</p> <p>Note: Examples must be for Ni, not other d block elements</p> <p>ALLOW 4s before 3d, ie $1s^2 2s^2 2p^6 3s^2 3p^6 4s^2 3d^8$</p> <p>ALLOW [Ar]$4s^2 3d^8$ OR [Ar]$3d^8 4s^2$</p> <p>ALLOW upper case D, etc and subscripts, e.g. [Ar]$4S_2 3D_8$</p> <p>DO NOT ALLOW highest energy shell is 'd' OR 'd is the outer sub-shell' (4s as well)</p> <p>ALLOW [Ar]$3d^8$</p> <p>ALLOW electron configurations with $4s^0$</p> <p>ALLOW for example $Ni^{3+} 1s^2 2s^2 2p^6 3s^2 3p^6 3d^7$ OR [Ar]$3d^7$</p> <p>No other Ni ions are acceptable</p> <p>ALLOW lone pair forms a coordinate bond to Ni^{2+} (which will also collect the coordinate bond mark)</p> <p>ALLOW diagram of $[Ni(H_2O)_6]^{2+}$ complex ion for 2nd marking point</p>
(b)	(i)	 <p>3D diagram ✓ 90° bond angle ✓</p>	<p>2</p>  <p>ALLOW dotted line OR unfilled wedge as alternatives for dotted wedge</p> <p>Accept bonds to H_2O (does not need to go to 'O')</p> <p>Accept 90° written by diagram.</p> <p>Charge NOT needed.</p> <p>Square brackets NOT needed</p>

F325

Mark Scheme




June 2011

Question			Answer	Mark	Guidance
6	(b)	(ii)	A: NiCl_4^{2-} ✓ B: $\text{Ni}(\text{OH})_2$ ✓	2	ALLOW $[\text{NiCl}_4]^{2-}$ DO NOT ALLOW $\text{Ni}(\text{Cl}^-)_4^{2-}$ ALLOW $\text{Ni}(\text{OH})_2(\text{H}_2\text{O})_4$ OR $[\text{Ni}(\text{OH})_2(\text{H}_2\text{O})_4]$
		(iii)	C: $[\text{Ni}(\text{NH}_3)_6]^{2+}$ ✓	1	Square brackets essential 2+ charge must be outside square brackets ALLOW $[\text{Ni}(\text{OH})_6]^{4-}$
		(iv)	$[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 6\text{NH}_3 \longrightarrow [\text{Ni}(\text{NH}_3)_6]^{2+} + 6\text{H}_2\text{O}$ ✓ ✓	2	1 mark for each side of equation ALLOW equilibrium sign ALLOW ECF from (iii) for the following: $[\text{Ni}(\text{NH}_3)_4]^{2+}$ (wrong number of NH_3) Any 6 coordinate Ni^{2+} complex with NH_3 and H_2O ligands, e.g. $[\text{Ni}(\text{NH}_3)_4(\text{H}_2\text{O})_2]^{2+}$, $[\text{Ni}(\text{NH}_3)_5(\text{H}_2\text{O})]^{2+}$, etc ALLOW from $[\text{Ni}(\text{OH})_6]^{4-}$, $[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 6\text{OH}^- \longrightarrow [\text{Ni}(\text{OH})_6]^{4-} + 6\text{H}_2\text{O}$ OR $[\text{Ni}(\text{H}_2\text{O})_6]^{2+} + 6\text{NH}_3 \longrightarrow [\text{Ni}(\text{OH})_6]^{4-} + 6\text{NH}_4^+$
	(c)	(i)	$\text{C}_{10}\text{H}_8\text{N}_2$ ✓	1	ALLOW atoms in any order
		(ii)	4 ✓	1	
		(iii)	 One mark for each structure 2nd structure must be correct mirror image of 1st structure	2	Charge and N atom labels NOT needed ALLOW any attempt to show bipy. Bottom line is the diagram on the left. 1 mark for 3D diagram with ligands attached for ONE stereoisomer. Must contain 2 out wedges, 2 in wedges and 2 lines in plane of paper: ALLOW structures with Ni in centre

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
6 (c) (iv)	<p>3 marks available</p> <p>1st mark Correct 4,4'-bipy structure shown separately or within attempted structure with Ni²⁺ ✓</p> <p>2 marks The remaining 2 marks are available for a section of the polymer with repeat unit identified as follows:</p> <p>IF Ni is bonded to 4 H₂O (bond to O) with a bond to N end of two 4,4'-bipy structure</p> <p>OR</p> <p>IF each N of 4,4'-bipy is bonded to a Ni bonded to 4 H₂O (bond to O), award 1 mark ✓</p> <p>IF correct repeat unit is shown, award 2 marks ✓✓</p> 	3	<p>ALLOW aromatic rings</p>  <p>Charge NOT needed. Square brackets NOT needed</p> <p>Bonds around Ni do NOT need to be shown 3D Accept bonds to H₂O (does NOT need to go to 'O')</p> <p>ALLOW the following structure for repeat unit for all 2nd and 3rd marks:</p> 
	Total	21	

F325

Mark Scheme

June 2011

Question		Answer	Mark	Guidance
7	(a)	<p>Definition The e.m.f. (of a half-cell) compared with a standard hydrogen half-cell/standard hydrogen electrode ✓</p> <p>Standard conditions Temperature of 298 K / 25°C AND (solution) concentrations of 1 mol dm⁻³ AND pressure of 101 kPa OR 100 kPa ✓</p>	2	<p>ALLOW voltage OR potential difference OR p.d. OR electrode potential OR reduction potential OR redox potential as alternative for e.m.f. IGNORE S.H.E. (as abbreviation for standard hydrogen electrode)</p> <p>ALLOW 1 atmosphere/1 atm OR 10⁵ Pa OR 1 bar</p>
	(b)	1.25 (V) ✓	1	IGNORE any sign
	(c)	(i)	2	<p>2 marks for correct equation ALLOW NiOOH OR NiO₂H</p> <p>ALLOW ⇌ sign for equation (ie assume reaction goes from left to right) ALLOW 1 mark for correctly balanced equation with e⁻ and/or OH⁻ shown e.g.: Cd + 2NiO(OH) + 2H₂O + 2OH⁻ + 2e⁻ → Cd(OH)₂ + 2Ni(OH)₂ + 2OH⁻ + 2e⁻</p> <p>ALLOW 1 mark for balanced correct reverse equation with OH⁻ AND e⁻ cancelled: Cd(OH)₂ + 2Ni(OH)₂ → Cd + 2NiO(OH) + 2H₂O</p>
		(ii)	2	<p>oxidation: Cd from 0 to +2 ✓ '+' sign not required reduction: Ni from +3 to +2 ✓ '+' sign not required</p> <p>ALLOW Cd⁰ → Cd²⁺ (shows 0 and 2+) ALLOW Ni³⁺ → Ni²⁺ (shows 3+ and 2+) ALLOW ECF from (c)(i) equation written 'wrong way around'.</p>
	(d)	(i)	1	<p>reverse reactions to charging OR Cd(OH)₂ + 2e⁻ → Cd + 2OH⁻ Ni(OH)₂ + OH⁻ → NiO(OH) + H₂O + e⁻ OR reaction that is reverse to reaction given in c(i): Cd(OH)₂ + 2Ni(OH)₂ → Cd + 2NiO(OH) + 2H₂O ✓</p> <p>If half-equations are given, then BOTH equations required</p> <p>ALLOW ⇌ sign for equation (ie assume reaction goes from left to right)</p>

F325

Mark Scheme

June 2011

Question			Answer	Mark	Guidance
7	(d)	(ii)	$4\text{OH}^- \longrightarrow \text{O}_2 + 2\text{H}_2\text{O} + 4\text{e}^- \checkmark$ $2\text{H}_2\text{O} + 2\text{e}^- \longrightarrow \text{H}_2 + 2\text{OH}^- \checkmark$	2	ALLOW multiples; ALLOW \rightleftharpoons sign for each equation Note: These are the only correct responses
Total				10	

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
8	<p>step 1 $\text{Cu} + 4\text{HNO}_3 \longrightarrow \text{Cu}^{2+} + 2\text{NO}_3^- + 2\text{NO}_2 + 2\text{H}_2\text{O}$ OR $\text{Cu} + 2\text{H}^+ + 2\text{HNO}_3 \longrightarrow \text{Cu}^{2+} + 2\text{NO}_2 + 2\text{H}_2\text{O}$ OR $\text{Cu} + 4\text{H}^+ + 2\text{NO}_3^- \longrightarrow \text{Cu}^{2+} + 2\text{NO}_2 + 2\text{H}_2\text{O} \checkmark$</p> <p>step 2 2 equations with 1 mark for each $\text{Cu}^{2+} + \text{CO}_3^{2-} \longrightarrow \text{CuCO}_3 \checkmark$ $2\text{H}^+ + \text{CO}_3^{2-} \longrightarrow \text{H}_2\text{O} + \text{CO}_2 \checkmark$</p> <p>step 4 $2\text{Cu}^{2+} + 4\text{I}^- \longrightarrow 2\text{CuI} + \text{I}_2 \checkmark$</p>	4	<p>ANNOTATE ALL Q8 WITH TICKS AND CROSSES, etc</p> <p>ALLOW multiples throughout IGNORE state symbols throughout</p> <p>ALLOW $\text{Cu}(\text{NO}_3)_2$ for $\text{Cu}^{2+} + 2\text{NO}_3^-$</p> <p>AWARD 2 MARKS for a combined equation: $\text{Cu}^{2+} + 2\text{H}^+ + 2\text{CO}_3^{2-} \longrightarrow \text{CuCO}_3 + \text{H}_2\text{O} + \text{CO}_2 \checkmark\checkmark$</p> <p>DO NOT ALLOW $2\text{H}^+ + \text{CO}_3^{2-} \longrightarrow \text{H}_2\text{CO}_3$</p> <p>ALLOW $2\text{Cu}^{2+} + 4\text{KI} \longrightarrow 2\text{CuI} + \text{I}_2 + 4\text{K}^+$ ALLOW $\text{Cu}^{2+} + \text{I}^- \longrightarrow \text{Cu}^+ + \frac{1}{2}\text{I}_2$</p>

F325

Mark Scheme

June 2011

Question	Answer	Mark	Guidance
8	<p>FIRST, CHECK THE ANSWER ON ANSWER LINE IF answer = 67.6%, award 5 marks. Ignore any attempted equation in step 4</p> <p>IF answer = 33.8% AND IF Cu²⁺/I₂ in step 4 equation shown with 1:1 molar ratio, award 5 marks for ECF</p> <p>-----</p> <p>amount S₂O₃²⁻ used = $0.100 \times \frac{29.8}{1000} = 2.98 \times 10^{-3} \text{ mol } \checkmark$</p> <p>amount I₂ = $1.49 \times 10^{-3} \text{ mol}$ OR amount Cu²⁺ = $2.98 \times 10^{-3} \text{ mol } \checkmark$</p> <p>amount Cu²⁺ in original 250 cm³ = 10 x 2.98×10^{-3} = $2.98 \times 10^{-2} \text{ mol } \checkmark$</p> <p>Mass of Cu/Cu²⁺ in brass = $63.5 \times 2.98 \times 10^{-2} \text{ g}$ = $1.8923 \text{ g } \checkmark$</p> <p>percentage Cu in brass = $\frac{1.8923}{2.80} \times 100$ = $67.6\% \checkmark$</p> <p>MUST be to one decimal place (in the question)</p>	5	<p>IF there is an alternative answer, check to see if there is any ECF credit possible using working below</p> <p>-----</p> <p>Working must be to 3 SF throughout until final % mark BUT ignore trailing zeroes, ie for 0.490 allow 0.49</p> <p>ECF answer above</p> <p>ECF 10 x answer above</p> <p>ECF 63.5 x answer above ALLOW 1.88 g</p> <p>ECF $\frac{\text{answer above}}{2.80} \times 100$ Answer must be to one decimal place</p> <p>ALLOW % Cu = 67.5 % IF mass of Cu has been rounded to 1.89 g in previous step</p> <p>Common ECFs: 6.76% x10 missing 3/5 marks for calculation 2 d.p. MS states 1 d.p.</p> <p>33.8% IF Cu²⁺/I₂ in step 4 equation with 2:1 ratio OR not attempted, response, 4/5 marks for calculation (moles Cu²⁺ incorrect)</p>
	Total	9	

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity



OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553

© OCR 2011