

1 This question is about compounds of Group 3 elements.

(a) Aluminium will combine directly with fluorine.

Write the equation for the reaction between aluminium and fluorine.

..... [1]

(b) Solid aluminium fluoride has a giant ionic lattice structure.

(i) Describe what is meant by the term *ionic lattice*, in terms of the type and arrangement of particles present.

.....  
.....  
.....  
..... [2]

(ii) Draw a 'dot-and-cross' diagram for aluminium fluoride.

Show outer electrons only.

[2]

(c) Solid boron tribromide has a simple molecular lattice structure. The atoms are held together by covalent bonds.

(i) What is meant by the term *covalent bond*?

.....  
..... [1]

(ii) Draw a 'dot-and-cross' diagram to show the bonding in a boron tribromide molecule.

Show outer electrons only.

[1]

(d) State whether the following substances conduct electricity when solid or molten, and explain your answers in terms of the particles involved:

- aluminium
- aluminium fluoride
- boron tribromide.

*In your answer you should use appropriate technical terms, spelled correctly.*

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
..... [5]

(e) Aluminium has 13 successive ionisation energies.

(i) Write the equation for the **third** ionisation energy of aluminium.


Include state symbols.

..... [1]

(ii) On the axes below, add crosses to show the 13 successive ionisation energies of aluminium.

The value for the first ionisation energy has been completed for you.

You do not have to join the crosses.



[2]

[Total: 15]






3 Oxides can have different types of bonding.

(a) H<sub>2</sub>O has hydrogen bonding.

(i) Complete the diagram below to show hydrogen bonding between the H<sub>2</sub>O molecule shown and **one** other H<sub>2</sub>O molecule.

Include relevant dipoles and lone pairs.

Label the hydrogen bond.



[2]

(ii) State and explain **two** anomalous properties of ice caused by hydrogen bonding.

1 .....

.....

.....

2 .....

.....

.....

[4]

(b) Draw a 'dot-and-cross' diagram to show the bonding in CO<sub>2</sub>.

Show outer electrons only.

[1]

(c) Silicon dioxide, SiO<sub>2</sub>, has the same structure and bonding as diamond.

State the structure and bonding in SiO<sub>2</sub>.

.....  
..... [1]

(d) Describe and explain the electrical conductivity of sodium oxide, Na<sub>2</sub>O, and sodium in their solid and molten states.




*In your answer you should use appropriate technical terms, spelled correctly.*

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
..... [5]

[Total: 13]

4 The hydrides of Group 5 elements all exist as gases at room temperature.

(a) Phosphine gas,  $\text{PH}_3$ , can be prepared by adding phosphorus,  $\text{P}_4$ , to warm concentrated aqueous sodium hydroxide as shown in the equation below.



(i) Using oxidation numbers, explain why this is a disproportionation reaction.

.....  
.....  
.....  
.....  
..... [3]

(ii) A chemist reacts 1.86 g of  $\text{P}_4$  with excess  $\text{NaOH}(\text{aq})$ .

Calculate the volume of phosphine gas, in  $\text{cm}^3$ , produced at room temperature and pressure, RTP.

volume of phosphine gas = .....  $\text{cm}^3$  [2]

(b) Phosphine gas burns in air to form an oxide of phosphorus,  $\text{P}_4\text{O}_{10}$ , and water.

Write the equation for this reaction.

..... [1]



(c) Phosphoric acid,  $\text{H}_3\text{PO}_4$ , can be made by reacting  $\text{P}_4\text{O}_{10}$  with water.

Sodium phosphate,  $\text{Na}_3\text{PO}_4$ , is a salt that can be prepared by reacting  $\text{H}_3\text{PO}_4$  with sodium hydroxide,  $\text{NaOH}$ .

A student prepared a solution of  $\text{Na}_3\text{PO}_4$  by reacting  $15.0\text{ cm}^3$  of  $0.100\text{ mol dm}^{-3}$   $\text{H}_3\text{PO}_4$  with  $0.200\text{ mol dm}^{-3}$   $\text{NaOH}$ .


(i) Why is  $\text{Na}_3\text{PO}_4$  described as a salt of  $\text{H}_3\text{PO}_4$ ?

.....  
..... [1]

(ii) Calculate the amount, in moles, of  $\text{H}_3\text{PO}_4$  in  $15.0\text{ cm}^3$  of  $0.100\text{ mol dm}^{-3}$   $\text{H}_3\text{PO}_4$ .

amount = ..... mol [1]

(iii) The equation for the preparation of  $\text{Na}_3\text{PO}_4$  from  $\text{NaOH}$  and  $\text{H}_3\text{PO}_4$  is shown below.



Calculate the volume of  $0.200\text{ mol dm}^{-3}$   $\text{NaOH}$  that reacts exactly with  $15.0\text{ cm}^3$  of  $0.100\text{ mol dm}^{-3}$   $\text{H}_3\text{PO}_4$ .

volume = .....  $\text{cm}^3$  [1]

(d) Ammonia,  $\text{NH}_3$ , is another gaseous Group 5 hydride.

$\text{NH}_3$  and  $\text{PH}_3$  are both simple molecules. The boiling points of  $\text{NH}_3$  and  $\text{PH}_3$  are shown in the table below.

Group 5 hydride	Boiling point / °C
$\text{NH}_3$	-33
$\text{PH}_3$	-88

(i) Complete the table below to show the main intermolecular forces present in  $\text{NH}_3$  and  $\text{PH}_3$ .

Group 5 hydride	Main intermolecular force
$\text{NH}_3$	
$\text{PH}_3$	


[2]

(ii) Suggest why  $\text{PH}_3$  has a lower boiling point than  $\text{NH}_3$ .

.....  
.....  
..... [1]

(e)  $\text{NH}_3$  reacts with molecules of  $\text{BF}_3$  to form  $\text{H}_3\text{NBF}_3$ , shown below.

One of the bonds in  $\text{H}_3\text{NBF}_3$  is a dative covalent bond.



(i) A covalent bond is a shared pair of electrons.

What is a *dative* covalent bond?

.....  
.....  
..... [1]

