

## Mark Scheme - AS 2.8 Instrumental Analysis

1 (a) (i) % H = 14.3 (1)

$$\text{C} : \text{H} = \frac{85.7}{12.0} : \frac{14.3}{1.01} = 7.14 : 14.16 \text{ (1)}$$




empirical formula = CH<sub>2</sub> (1) [3]

(ii) M<sub>r</sub> = 42/ largest fragment has mass 42 (1)

(CH<sub>2</sub> = 14) therefore molecular formula = C<sub>3</sub>H<sub>6</sub> (1) [2]


(iii) CH<sub>3</sub> is present [1]

(b) 1 mark for each [3]



Total [9]

- 2 (a) (i) 1 mark for arrows in first diagram; 1 mark for arrow in second diagram;  
1 mark for all charges



2 max if incorrect isomer given [3]

- (ii) 2-bromopropane formed from a secondary carbocation (1)  
Secondary carbocations are more stable than primary carbocations (1)  
[2]

(b) Empirical formula =  $C_3H_5Br$  (1)

Molecular formula =  $C_3H_5Br$


(must show use of mass spectrum to gain this mark) (1)

Two molecular ion peaks as there are two isotopes of bromine (1)

Peaks at 15 =  $CH_3^+$  and 41 =  $C_3H_5^+$  (1)

$550\text{ cm}^{-1}$  = C-Br       $1630\text{ cm}^{-1}$  = C=C       $3030\text{ cm}^{-1}$  = C-H (1)

Molecule is:







(1) [6]

QWC: legibility of text, accuracy of spelling, punctuation and grammar, clarity of meaning [1]

**Total [12]**


- 3 (a) (i) Mass C =  $1.79 \times 12/44 = 0.488$  (g) [1]
- (ii) Mass O = 0.65 (g) ecf from part (i) [1]
- (iii) C : H : O =  $0.488/12 : 0.061/1 : 0.65/16 = 0.0407 : 0.061 : 0.0406$  (1)  
 = 2:3:2 empirical formula is  $C_2H_3O_2$  (1)  
 No ecf from incorrect ratios [2]
- (iv) Mr of empirical formula = 59 so molecular formula is  $C_4H_6O_4$  so  
 F is acid 2/ molecular formula acid 1 is  $C_5H_8O_2$  so empirical formula is  
 not  $C_2H_3O_2$  molecular formula acid 2 is  $C_4H_6O_4$  so empirical formula is  
 $C_2H_3O_2$  [1]
- (v) Bromine turns from brown/red-brown to colourless for Acid 1 [1]
- (vi) 
$$\begin{array}{ccccccc} & H & H & H & H & & \\ & | & | & | & | & & \\ HO & -C & -C & -C & -C & -OH & \\ & | & | & | & | & & \\ & H & H & H & H & & \end{array}$$
 [1]
- (b) (i) Mr / molecular ion (is 46) [1]
- (ii)  $CH_3$  (present) [1]
- (iii) OH (present) [1]
- (c) Ethene to ethanol: steam (1)  
 $H_3PO_4$  (catalyst) (1)  
 Ethanol to ethene: conc  $H_2SO_4$ /  $Al_2O_3$ / pumice (1)  
 High temperature >  $150^\circ C$  for  $H_2SO_4$   
 >  $300^\circ C$  for  $Al_2O_3$ / pumice (1) [4]

**Total [14]**



[4]

- (b) (i) T neutral and sweet-smelling therefore an ester (1)  
 Infrared spectrum at  $1750\text{ cm}^{-1}$  shows C=O and at  $3000\text{ cm}^{-1}$  shows O-H therefore X is an acid (1)  
 Y is an alcohol, formed from ethanal must be ethanol (1)  
 5 carbons in ester therefore X must be propanoic acid (1)  
 Structure of T is



(Maximum 4 marks) [4]

QWC Legibility of text; accuracy of spelling, punctuation and grammar, clarity of meaning (1)  
 Selection of a form and style of writing appropriate to purpose and to complexity of subject matter (1) [2]

- (ii) I Reagent to form Y is  $\text{NaBH}_4$  /  $\text{LiAlH}_4$  [1]  
 II Sulfuric acid acts as a catalyst / removes water so pushes equilibrium to right [1]

(c)	$\text{CH}_3(\text{CH}_2)$	0.1 to 2.0 ppm triplet (1)	
	$(\text{CH}_2)\text{CH}_2\text{O}$	3.5 to 4.0 ppm quadruplet (1)	
	$\text{CH}_2\text{CO}$	2.5 to 3.0 ppm singlet (1)	
	$\text{CH}_3\text{CO}$	2.0 to 2.5 ppm singlet (1)	[4]

(d)	Isomer P (1)		
	Only P can form hydrogen bonds between molecules (1)		
	Hydrogen bonds are the strongest intermolecular bonds / need more energy to break hydrogen bonds (1)		[3]

QWC The information is organised clearly and coherently, using specialist vocabulary where appropriate [1]

**Total [20]**