



Mark Scheme - 4.2 Aromaticity

1. (a) Any valid ester structure with formula $C_{10}H_{12}O_2$ [1]

Examples:



- (b) (i) Compound X [1]



- (iii) Rotate the plane of polarised light in opposite directions [1]

(c)

Reagent(s)	Observation if the test is positive	Compound(s) that would give a positive result
$I_2 / NaOH$ (aq)	Yellow solid	X
Na_2CO_3 (aq)	Bubbles of colourless gas / effervescence	W
$FeCl_3$ (aq)	Dark purple/blue/green - do not accept 'precipitate'	X, Z

(1 mark for each box) [6]


(d) (i) Heat / Alkaline / Potassium manganate(VII) / then acidify
 (1 mark for Potassium manganate + 1 other point; 2 marks for all) [2]

(ii) I. Addition polymer – One large molecule formed only / Condensation polymer – one large molecule with small molecules (e.g. water) lost. (1)

Addition polymer – one starting material / Condensation – two starting materials


OR Addition polymer – one functional group in each molecule / Condensation polymer – two functional groups in each molecule (1) [2]

II.




(e) (i) NaBH_4 / LiAlH_4 or name(1) Reduction (1) [2]
 - ignore conditions unless LiAlH_4 in water - do not accept 'redox'

(ii)



Accept structures with only one -OH group reacted. [1]

(iii) [1]



[19 marks]

2.

- (a) Benzene is a compound whose molecules contain six carbon atoms bonded in a (hexagonal) ring (1)
All the carbon to carbon bond lengths are equal / intermediate (1)
Each carbon atom is bonded to two other carbon atoms and a hydrogen atom (1) by σ -bonds (1)
All the C – C – C angles are the same / 120° (1)
The remaining p electron of each carbon atom / overlap of p orbitals forms a delocalised cloud of electrons / π -system (1) above and below the plane (1)
Credit can be gained from labelled diagram
[Candidates can gain a maximum of (4) for this part]

This delocalisation increases the **stability** (1) of the molecule and this stability is maintained by benzene undergoing substitution reactions in preference to addition reactions (that would destroy the delocalised system)
The π -cloud is **electron rich** and will be attracted to electron deficient electrophiles (1)
[Candidates can gain (2) for this part]


[6]

QWC Selection of a form and style of writing appropriate to purpose and to complexity of subject matter (1)

Legibility of text; accuracy of spelling, punctuation and grammar; clarity of meaning. (1)

QWC [2]

(b)



catalyst eg AlCl_3 (anhydrous) (1)

[2]


- (c) (i) (There are two environments for the protons), the 3 aromatic protons at $\sim 6.8 \delta$ and the 9 methyl / aliphatic protons at $\sim 2.3 \delta$ (1)
 These give a peak area of 3:9, ie. 1:3 (1)
 These environments are separate / discrete (1) therefore no splitting pattern [3]

- (ii) Dissolve in the minimum volume (1)
 Of hot water (1)
 (Filter hot) (1)
 Cool (1)
 Filter (1)
 Dry (1)

(up to 4 max but candidates must give the first two points in order to gain full credit)

[4]

(iii)



[1]

- (iv) Reagent S is alkaline potassium manganate(VII) (1)


Reagent T is eg hydrochloric acid (1)

[2]

Total [20]

3.

(a) (i) Electrophilic substitution [1]



Formation of Br^+ (1), curly arrows (1), intermediate (1) [3]

(b) (i) The extra stability in the benzene molecule due to electron delocalisation / the difference in energy between the experimental ΔH^\ominus reaction for benzene and the ΔH^\ominus reaction according to the Kekulé structure [1]

(ii) If benzene had 3 double bonds enthalpy change would be $3 \times -120 = -360 \text{ kJ mol}^{-1}$ (1)

Delocalisation energy is difference between -360 and $-208 = 152 \text{ kJ mol}^{-1}$ (1) [2]

(c) Benzene is carcinogenic / toxic [1]

4.

(a) (i) (Concentrated) nitric acid / (concentrated) sulfuric acid / Temperature of 40-80°C
(Any 2 = 1 mark; All 3 = 2 marks)
Electrophilic substitution (1) [3]

(ii) I. Peak area is proportional to amount of substance (1)
Percentage = $(30 / 38) \times 100 = 79\%$ (1)
(Can obtain both marks from correct percentage) [2]

II. 45 = COOH⁺, 46 = NO₂⁺, 122 = C₆H₄NO₂⁺ and 167 = C₇H₅NO₄⁺.
(Any 2 = 1 mark; All 4 = 2 marks) [2]

(iii) I. Lower melting point / melts over a range [1]
II. 1 mark for each point.


- Dissolve in the minimum volume
- Of hot water
- Filter hot
- Allow to cool
- Filter
- Dry residue under suction / in oven below 142°C

Max 4 marks [4]

QWC: legibility of text, accuracy of spelling, punctuation and grammar, clarity of meaning. [1]

(b) (i) Tin and concentrated hydrochloric acid [1]

(ii) Below 10°C (1)




(iii) N=N double bond is chromophore (1)
Compound absorbs blue / green / complementary colours to red / all colours but red (1)
Remaining frequencies are transmitted, giving the red colour seen. (1)
Any 2 out of 3 [2]

(c) Nitrogen has a lone pair (1) which can accept a proton (1) [2]

[20 marks]

5.

- (a) (i) Both molecules have lone pairs on nitrogen (1)
The lone pairs can form (coordinate) bonds with H^+ ions (1) [2]
- (ii) Lone pair on N in phenylamine is delocalised over benzene ring (1) therefore less able to accept H^+ (1) [2]
- (iii) I Arrow in first step (1)
Cation structure in second step (1)
Arrow in second step (1)



- II (fractional) distillation / steam distillation [3]
[1]
- III Sn and conc. HCl (1) followed by NaOH (1) [2]

6.

(a)blue (1)higher (1)higher (1) [3]




accept C₆H₅ in place of the ring accept equations that show the catalyst

(ii) It acts as a halogen carrier / it helps produce the electrophile/CH₃⁺ / increases polarity of the halogenoalkane [1]

(c) There are 6 methyl protons and 4 aromatic protons, hence a ratio of 3:2 (1)
All the methyl protons are equivalent as are all the aromatic protons (1) [2]

(d) (i) Any 2 from NMR / HPLC / GC / refractive index / mass spectra / boiling temperature [2]




(ii) protein / dipeptide / polypeptide [1]

Total [12]

7.

- (a) $\text{CH}_3\text{CH}(\text{CH}_3)\text{CH}_2\text{Cl}$ (1) $\text{AlCl}_3 / \text{FeCl}_3$ (1) Room temperature / in the dark (1) [3]
- (b) (i) 2,4-DNP (1) Orange precipitate (1) [2]
- (ii) Tollen's reagent (1) Silver mirror with C, no reaction with B (1) [2]
- (c) Optical isomerism is where a molecule and its mirror image are different / non-superimposable (1)
Compound C has a chiral centre / 4 different groups attached to one carbon atom (1)



The two isomers rotate the plane of polarised light in opposite directions (1) [4]

QWC: organisation of information clearly and coherently; use of specialist vocabulary where appropriate (1) [1]

- (d) Dilute acid (1) heat (1) hydrolysis (1) [3]
- (e) Acidified potassium dichromate (VI) (1) / heat (1)

One step reactions are generally better as they have a better yield / there is waste in each stage (1)

Two step process may be cheaper / use more sustainable reagents/ may give a better yield in this case / produce less harmful waste materials / potassium dichromate may react with other parts of the molecule as well / may be easier to separate product (1)

Do not credit same idea twice e.g. if 'better yield' gains first mark, a different point is required to gain second mark [4]

QWC: selection of a form and style of writing appropriate to purpose and to complexity of subject matter [1]

Total [20]