



Cambridge International Examinations
Cambridge International General Certificate of Secondary Education

CHEMISTRY

0620/43

Paper 4 Theory (Extended)

May/June 2017

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

© IGCSE is a registered trademark.

This syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **8** printed pages.

Question	Answer	Marks
1(a)(i)	J	1
1(a)(ii)	E	1
1(a)(iii)	D	1
1(a)(iv)	J	1
1(a)(v)	L	1
1(a)(vi)	D	1
1(b)(i)	(atoms with) same number of protons / atomic number / of same element	1
	different number of neutrons / different mass number / different nucleon number	1
1(b)(ii)	E AND G	1
1(b)(iii)	they have the same number of electrons in their outer shell	1

Question	Answer	Marks
2(a)(i)	CH ₂	1
2(a)(ii)	initial colour: orange	1
	final colour: colourless/none	1
2(b)(i)	approximately horizontal line draw to right of and below the reagent line	1
	energy change shown starting level with the reactant energy AND finishing level with the product energy AND having only one (correct) arrow head AND labelled ΔH /energy change	1
2(b)(ii)	(energy required to break bonds =) 3624	1
	(energy given out when bonds made =) 3738	1
	-114 (kJ/mol)	1
2(c)	(attractive) forces between molecules	1
	(forces of attraction) are stronger in iodine	1

Question	Answer	Marks
3(a)	regular arrangement / lattice of positive ions / magnesium ions / Mg^{2+} ions	1
	sea of electrons OR delocalised electrons	1
	attraction between (positive) ions and (delocalised / sea of) electrons	1
3(b)	electrons	1
	move / flow (throughout / through the structure)	1
3(c)	layers (of atoms or ions)	1
	layers / atoms / ions can slide / slip / glide (over each other) (without breaking the metallic bonds)	1
3(d)(i)	magnesium shown as (2, 8) using crosses	1
	sulfide shown as (2, 8, 8), with the two gained electrons in the outer shell of sulfur shown as crosses and all other electrons on sulfur shown as dots	1
	magnesium ion charge as 2^+ AND sulfide charge as 2^-	1
3(d)(ii)	melt / fused	1
	ions can move OR are mobile	1

Question	Answer	Marks
4(a)	petroleum	1
4(b)(i)	<i>saturated</i> : only single bonds OR no double / multiple bonds (between carbon atoms)	1
	<i>hydrocarbon</i> : (compound that) contains carbon and hydrogen	1
	<i>hydrocarbon</i> : and no other elements / only	1
4(b)(ii)	alkane(s)	1
4(b)(iii)	any 2 from: <ul style="list-style-type: none"> • same / similar chemical properties • (same) general formula • (consecutive members) differ by CH₂ • same functional group • common (allow similar) methods of preparation • physical properties vary in predictable manner / show trends / gradually change / example of a physical property variation 	2
4(b)(iv)	CO ₂ and H ₂ O on right-hand side and no other products / reagents	1
	11 (O ₂), 7 (CO ₂), 8 (H ₂ O)	1
4(c)(i)	acid rain	1

Question	Answer	Marks
4(c)(ii)	<i>carbon monoxide</i> : from incomplete combustion (of fuel)	1
	<i>oxides of nitrogen</i> : nitrogen (from the air) reacts with oxygen (from the air)	1
	<i>oxides of nitrogen</i> : at high temperatures (in engine) OR (electrical) spark (in the engine)	1
4(c)(iii)	poisonous / toxic / death	1
4(c)(iv)	any 3 from: <ul style="list-style-type: none"> oxides of nitrogen are reduced / lose oxygen (to form nitrogen) oxides of nitrogen form nitrogen (oxides of nitrogen) react with carbon monoxide gases (adsorb / stick) on the catalyst's surface 	3
4(d)(i)	butane	1
4(d)(ii)	(molecules with) the same molecular formula	1
	different structural formula / different displayed formula	1
4(d)(iii)	UV light / sunlight	1
	H–C \downarrow	1
	any mono to deca chloro-substituted derivative of methyl propane	1

Question	Answer	Marks
5(a)(i)	loss (of electrons)	1
5(a)(ii)	$\text{Ni} \rightarrow \text{Ni}^{2+} + 2\text{e}^{-}$	1
5(a)(iii)	goes down/gets less/decreases/lower/smaller	1
5(b)(i)	beryllium	1
	most negative voltage with any (named) metal OR biggest voltage with cobalt/nickel	1
5(b)(ii)	cobalt AND nickel	1
5(b)(iii)	– sign	1
	2.7	1
5(c)	(set up cell) using magnesium and beryllium (electrodes)	1
	voltage positive if magnesium is metal 2	1
	OR	
	(set up cells) using both magnesium and beryllium with the same metal as the other electrode	1
	larger (magnitude) voltages with magnesium	1
	OR	
	use magnesium with a different metal and compare to a reference value in a table	1
value is more negative than with beryllium, if magnesium is metal 1	1	

Question	Answer	Marks
6(a)(i)	$\text{BaCO}_3 \rightarrow \text{BaO} + \text{CO}_2$	1
6(a)(ii)	anything pH in the range pH 10 to pH 14	1
6(a)(iii)	nitrogen dioxide	1
	oxygen	1
6(b)(i)	$\text{Na}_2\text{CO}_3 + \text{Ba}(\text{NO}_3)_2 \rightarrow \text{BaCO}_3 + 2\text{NaNO}_3$ M1 formula of NaNO_3 M2 equation fully correct	2
6(b)(ii)	filter	1
	wash (the residue) using water	1
	dry the residue between filter papers/in a warm place	1
6(c)(i)	$M_r = 197$	1
	$(9.85 / 197 =) 0.05 \text{ (mol)}$	1
6(c)(ii)	0.05 (mol)	1
6(c)(iii)	$(0.05 \times 24) = 1.2 \text{ (dm}_3\text{)}$	1
6(c)(iv)	moles of HCl at the start = $(250 / 1000 \times 1.00) = 0.25$	1
	moles HCl in excess = $0.25 - (2 \times 0.05) = 0.15 \text{ (mol)}$	1