

Please check the examination details below before entering your candidate information

Candidate surname					Other names									
<b>Pearson Edexcel</b> <b>International</b> <b>Advanced Level</b>					Centre Number					Candidate Number				
					<input type="text"/>					<input type="text"/>				
Sample Assessment Materials for first teaching September 2018														
(Time: 1 hour 30 minutes)							Paper Reference <b>WMA13/01</b>							
<b>Mathematics</b> <b>International Advanced Level</b> <b>Pure Mathematics P3</b>														
<b>You must have:</b> Mathematical Formulae and Statistical Tables, calculator												Total Marks		

**Candidates may use any calculator permitted by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.**

### Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided  
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

### Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 10 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets  
– *use this as a guide as to how much time to spend on each question.*

### Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.
- If you change your mind about an answer, cross it out and put your new answer and any working underneath.

Turn over ►

S59757A

©2018 Pearson Education Ltd.

1/1/1/



Pearson

Answer ALL questions. Write your answers in the spaces provided.

1. Express

$$\frac{6x+4}{9x^2-4} - \frac{2}{3x+1}$$

as a single fraction in its simplest form.

(4)

$$\frac{6x+4}{(3x+2)(3x-2)} - \frac{2}{3x+1} = \frac{2(3x+2)}{(3x+2)(3x-2)} - \frac{2}{3x+1}$$

$$= \frac{2}{3x-2} - \frac{2}{3x+1} = \frac{2(3x+1) - 2(3x-2)}{(3x-2)(3x+1)} = \frac{6x+2-6x+4}{(3x-2)(3x+1)}$$

$$= \frac{6}{(3x-2)(3x+1)}$$

2.

$$f(x) = x^3 + 3x^2 + 4x - 12$$

(a) Show that the equation  $f(x) = 0$  can be written as

$$x = \sqrt{\left(\frac{4(3-x)}{3+x}\right)} \quad x \neq -3 \quad (3)$$

The equation  $x^3 + 3x^2 + 4x - 12 = 0$  has a single root which is between 1 and 2

(b) Use the iteration formula

$$x_{n+1} = \sqrt{\left(\frac{4(3-x_n)}{3+x_n}\right)} \quad n \geq 0$$

with  $x_0 = 1$  to find, to 2 decimal places, the value of  $x_1$ ,  $x_2$  and  $x_3$  (3)

The root of  $f(x) = 0$  is  $\alpha$ .

(c) By choosing a suitable interval, prove that  $\alpha = 1.272$  to 3 decimal places. (2)

$$(a) f(x) = x^3 + 3x^2 + 4x - 12 = 0$$

$$3x^2 = 12 - 4x - x^3$$

$$3x^2 = 4(3-x) - x^3$$

$$3x^2 + x^3 = 4(3-x)$$

$$x^2(3+x) = 4(3-x)$$

$$\frac{x^2(3+x)}{(3+x)} = \frac{4(3-x)}{(3+x)}$$

$$\sqrt{x^2} = \sqrt{\frac{4(3-x)}{(3+x)}}$$

$$x = \sqrt{\left(\frac{4(3-x)}{(3+x)}\right)}$$

$$(b) x_0 = 1$$

$$x_1 = \sqrt{\left(\frac{4(3-1)}{(3+1)}\right)} = \sqrt{2} = 1.41$$

$$x_2 = \sqrt{\left(\frac{4(3-1.41)}{(3+1.41)}\right)} = 1.20$$

$$x_3 = \sqrt{\left(\frac{4(3-1.20)}{(3+1.20)}\right)} = 1.31$$

## Question 2 continued

$$(i) \alpha = 1.2720$$

$$f(1.2715) = (1.2715)^3 + 3(1.2715)^2 + 4(1.2715) - 12 = -8.21 \times 10^{-3}$$

$$f(1.2725) = (1.2725)^3 + 3(1.2725)^2 + 4(1.2725) - 12 = 8.27 \times 10^{-3}$$

There is a change of sign which implies that there is a root in between 1.2715 and 1.2725

$$\frac{1.2715 + 1.2725}{2} = 1.2720$$

2

3.




Figure 1

Figure 1 shows a sketch of part of the graph  $y = f(x)$  where

$$f(x) = 2|3 - x| + 5 \quad x \geq 0$$

(a) Solve the equation

$$f(x) = \frac{1}{2}x + 30 \tag{3}$$

Given that the equation  $f(x) = k$ , where  $k$  is a constant, has two distinct roots,

(b) state the set of possible values for  $k$ . (2)

$2 3-x  + 5 = \frac{1}{2}x + 30$	$-2(3-x) + 5 = \frac{1}{2}x + 30$
$2(3-x) + 5 = \frac{1}{2}x + 30$	$-6 + 2x + 5 = \frac{1}{2}x + 30$
$6 - 2x + 5 = \frac{1}{2}x + 30$	$\frac{3}{2}x = 31$
$\frac{5}{2}x = -19$	$x = 62/3$
$x = -38/5$	

Since  $x \geq 0$ ,  $x = 62/3$

(b)  $f(x) = k$  has two distinct roots (i.e. 2 intersections)

turning point / vertex = (3, 5)

y axis ( $x=0$ )  $2(3-0) + 5$

$$2(3) + 5 = 11 \quad (0, 11)$$

$$5 \leq k \leq 11$$

when  $k > 11$ , root is negative

$\therefore k$  cannot be greater than 11



4. (i) Find

$$\int_5^{13} \frac{1}{(2x-1)} dx$$

writing your answer in its simplest form.

(4)

(ii) Use integration to find the exact value of

$$\int_0^{\pi/2} \sin 2x + \sec \frac{1}{3} x \tan \frac{1}{3} x dx$$

(3)

$$(i) \int_5^{13} \frac{1}{(2x-1)} = \frac{\ln|2x-1|}{2} = \frac{1}{2} \ln|2x-1|$$

$$\left[ \frac{1}{2} \ln|2x-1| \right]_5^{13}$$

$$\frac{1}{2} \ln 25 - \frac{1}{2} \ln 9$$

$$\frac{1}{2} \ln \left( \frac{25}{9} \right) \quad \ln \left( \frac{25}{9} \right)^{1/2} = \ln \left( \frac{5}{3} \right)$$

$$(ii) \int_0^{\pi/2} \sin 2x + \sec \frac{1}{3} x \tan \frac{1}{3} x \quad \sin 2x = \frac{-\cos 2x}{2}$$

$$\sec x \tan x = \sec x$$

$$\left[ -\frac{1}{2} \cos 2x + 3 \sec \frac{1}{3} x \right]_0^{\pi/2}$$

$$\sec \frac{1}{3} x \tan \frac{1}{3} x = \frac{\sec \frac{1}{3} x}{1/3}$$

$$-\frac{1}{2} \cos \left( 2 \times \frac{\pi}{2} \right) + 3 \sec \left( \frac{1}{3} \times \frac{\pi}{2} \right) =$$

$$\left[ -\frac{1}{2} \cos \left( 2 \left( \frac{\pi}{2} \right) \right) + 3 \sec \left( \frac{1}{3} \left( \frac{\pi}{2} \right) \right) \right]$$

$$-\frac{1}{2} (-1) + 3 \left( \frac{2\sqrt{3}}{3} \right)$$

$$- \left[ -\frac{1}{2} \cos (2 \times 0) + 3 \sec \left( \frac{1}{3} \times 0 \right) \right]$$

$$\frac{1}{2} + 2\sqrt{3} = \frac{1 + 4\sqrt{3}}{2}$$

$$= \frac{1 + 4\sqrt{3}}{2} - \frac{5}{2} = -2 + 2\sqrt{3}$$

$$-\frac{1}{2} \cos (2 \times 0) + 3 \sec \left( \frac{1}{3} \times 0 \right)$$

$$-\frac{1}{2} (1) + 3 (1) = \frac{5}{2}$$

$$= 2\sqrt{3} - 2$$

5. Given that

$$y = \frac{5x^2 - 10x + 9}{(x-1)^2} \quad x \neq 1$$

show that  $\frac{dy}{dx} = \frac{k}{(x-1)^3}$ , where  $k$  is a constant to be found.

(6)

$$y = \frac{5x^2 - 10x + 9}{(x-1)^2} = u - v$$

$$du/dx = 10x - 10$$

$$dv/dx \rightarrow \text{let } u = x-1 \quad du/dx = 1$$

$$v = u^2 \quad dy/dv = 2u$$

$$\frac{vdu/dx - u dv/dx}{v^2}$$

$$1 \times 2u = 2u$$

$$2(x-1)$$

$$\frac{(x-1)^2(10x-10) - (5x^2-10x+9)(2(x-1))}{(x-1)^4}$$

$$\frac{(x-1) [(x-1)(10x-10) - 2(5x^2-10x+9)]}{(x-1)^3}$$

$$(x-1)(10x-10) - 2(5x^2-10x+9)$$

$$x(10x-10) - 1(10x-10) - 10x^2 + 20x - 18$$

$$10x^2 - 10x - 10x + 10 - 10x^2 + 20x - 18 = -8$$

$$\frac{dy}{dx} = \frac{-8}{(x-1)^3} \quad k = -8$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


6. The functions  $f$  and  $g$  are defined by

$$f: x \mapsto e^x + 2 \quad x \in \mathbb{R}$$

$$g: x \mapsto \ln x \quad x > 0$$

- (a) State the range of  $f$ . (1)
- (b) Find  $fg(x)$ , giving your answer in its simplest form. (2)
- (c) Find the exact value of  $x$  for which  $f(2x + 3) = 6$  (4)
- (d) Find  $f^{-1}$  stating its domain. (3)
- (e) On the same axes sketch the curves with equation  $y = f(x)$  and  $y = f^{-1}(x)$ , giving the coordinates of all the points where the curves cross the axes. (4)

(a)  $f(x) = e^x + 2$



Range:  $f(x) > 2$

(b)  $fg(x)$

$$g(x) = \ln x$$

$$e^{\ln x} + 2 = x + 2$$

(c)  $f(2x + 3) = 6$

$$e^{2x + 3} + 2 = 6$$

$$e^{2x + 3} = 4$$

$$\ln e^{2x + 3} = \ln 4$$

$$2x + 3 = \ln 4$$

$$2x = \ln 4 - 3$$

$$x = \frac{\ln 4 - 3}{2} = \ln 2 - \frac{3}{2}$$



## Question 6 continued

(d)  $f^{-1}(x)$

$$y = e^x + 2$$

$$x = e^y + 2$$

$$e^y = x - 2$$


domain:  $x > 2$

$$\ln e^y = \ln(x - 2)$$

$$y = \ln(x - 2)$$

$$f^{-1}(x) = \ln(x - 2)$$

(e)  $y = f(x)$  and  $y = f^{-1}(x)$



7. The point  $P$  lies on the curve with equation

$$x = (4y - \sin 2y)^2$$

Given that  $P$  has  $(x, y)$  coordinates  $\left(p, \frac{\pi}{2}\right)$ , where  $p$  is a constant,

(a) find the exact value of  $p$

(1)

The tangent to the curve at  $P$  cuts the  $y$ -axis at the point  $A$ .

(b) Use calculus to find the coordinates of  $A$ .

(6)

$$(a) \quad x = (4y - \sin 2y)^2 \quad \left(p, \frac{\pi}{2}\right)$$

$$x = \left[4\left(\frac{\pi}{2}\right) - \sin\left(2 \cdot \frac{\pi}{2}\right)\right]^2$$

$$= \underline{4\pi^2}$$

$$(b) \quad x = (4y - \sin 2y)^2 \quad \therefore \frac{dx}{dy} = 2(4y - \sin 2y)(4 - 2\cos 2y)$$

$$y = \frac{\pi}{2} \quad \therefore \frac{dx}{dy} = \frac{1}{24\pi}$$

equation of tangent

$$y - \frac{\pi}{2} = \frac{1}{24\pi} (x - 4\pi^2)$$

$$y - \frac{\pi}{2} = \frac{1}{24\pi} (x - 4\pi^2) \quad x = 0 \quad \therefore y = \frac{\pi}{3}$$

8. In a controlled experiment, the number of microbes,  $N$ , present in a culture  $T$  days after the start of the experiment were counted.

$N$  and  $T$  are expected to satisfy a relationship of the form

$$N = aT^b \quad \text{where } a \text{ and } b \text{ are constants}$$

- (a) Show that this relationship can be expressed in the form

$$\log_{10} N = m \log_{10} T + c$$

giving  $m$  and  $c$  in terms of the constants  $a$  and/or  $b$ .

(2)




Figure 2

Figure 2 shows the line of best fit for values of  $\log_{10} N$  plotted against values of  $\log_{10} T$

- (b) Use the information provided to estimate the number of microbes present in the culture 3 days after the start of the experiment.

(4)

- (c) With reference to the model, interpret the value of the constant  $a$ .

(1)



## Question 8 continued

$$(a) N = aT^b$$

$$\log N = \log_{10} aT^b$$

$$\log_{10} N = \log_{10} a + \log_{10} T^b$$

$$\log_{10} N = \log_{10} a + b \log_{10} T$$

$$\log_{10} N = b \log_{10} T + \log_{10} a$$

$$m = b$$

$$c = \log_{10} a$$

$$(b) c = \log_{10} a = 1.84$$

$$10^{1.84} = 69.183 = a$$

$$T =$$

$$(0, 1.84) (1.22, 4.6)$$

$$\frac{4.6 - 1.84}{1.22 - 0} = \frac{2.76}{1.22} = 2.26$$

$$m = 2.26$$

$$N = 69.183 (T)^{2.26}$$

$$N = 69.183 (3)^{2.26} = 828.5 \quad \times$$

(c) "a" is the no. of microbes 1 day <sup>after the</sup> start of the experiment.



9. (a) Prove that

$$\sec 2A + \tan 2A \equiv \frac{\cos A + \sin A}{\cos A - \sin A} \quad A \neq \frac{(2n+1)\pi}{4} \quad n \in \mathbb{Z} \quad (5)$$

(b) Hence solve, for  $0 \leq \theta < 2\pi$

$$\sec 2\theta + \tan 2\theta = \frac{1}{2}$$

Give your answers to 3 decimal places.

(4)

(a)  $\sec 2A + \tan 2A$

$$\frac{1}{\cos 2A} + \frac{\sin 2A}{\cos 2A}$$

$$\begin{aligned} 1 &= \sin^2 A + \cos^2 A & \sin^2 A + \cos^2 A + 2\sin A \cos A \\ \sin 2A &= 2\sin A \cos A & \cos^2 A - \sin^2 A \\ \cos 2A &= \cos^2 A - \sin^2 A & = 1 + 2\sin A \cos A \rightarrow \sin 2A \\ & & \cos^2 A - \sin^2 A \rightarrow \cos 2A \end{aligned}$$

$$\frac{\cos^2 A + \sin^2 A + 2\sin A \cos A}{\cos^2 A - \sin^2 A} = \frac{(\cos^2 A + 2\sin A \cos A + \sin^2 A)}{(\cos A - \sin A)(\cos A + \sin A)} \quad \text{continuation}$$

(b)  $\frac{\cos A + \sin A}{\cos A - \sin A} = \frac{1}{2}$        $2\cos A + 2\sin A = \cos A - \sin A$   
 $\frac{\cos A}{\cos A} = \frac{-3\sin A}{\cos A}$

$$\begin{aligned} -3\tan A &= 1 \\ \tan A &= -\frac{1}{3} \end{aligned} \quad \begin{array}{l} \sqrt{\sin A} \\ \hline \sqrt{\cos A} \end{array}$$

$$\begin{aligned} &= -0.32175 \\ &= 2.8198, 5.9619 \\ &= 2.82, 5.96 \end{aligned}$$

## Question 9 continued

$$(a) \frac{(\cancel{\cos A} + \cancel{\sin A})(\cos A + \sin A)}{(\cancel{\cos A} + \cancel{\sin A})(\cos A - \sin A)} = \frac{\cos A + \sin A}{\cos A - \sin A}$$

10. The amount of an antibiotic in the bloodstream, from a given dose, is modelled by the formula

$$x = De^{-0.2t}$$

where  $x$  is the amount of the antibiotic in the bloodstream in milligrams,  $D$  is the dose given in milligrams and  $t$  is the time in hours after the antibiotic has been given.

A first dose of 15 mg of the antibiotic is given.

- (a) Use the model to find the amount of the antibiotic in the bloodstream 4 hours after the dose is given. Give your answer in mg to 3 decimal places. (2)

A second dose of 15 mg is given 5 hours after the first dose has been given. Using the same model for the second dose,

- (b) show that the total amount of the antibiotic in the bloodstream 2 hours after the second dose is given is 13.754 mg to 3 decimal places. (2)

No more doses of the antibiotic are given. At time  $T$  hours after the second dose is given, the total amount of the antibiotic in the bloodstream is 7.5 mg.

- \* (c) Show that  $T = a \ln\left(b + \frac{b}{e}\right)$ , where  $a$  and  $b$  are integers to be determined. (4)

(a)  $x = De^{-0.2t}$        $t = 4$

$$x = 15e^{-0.2(4)} = 6.739934962 = 6.740 \text{ mg}$$

(b)  $D = 15$

$t = 2$        $15e^{-0.2(2)} = 10.0548$        $15e^{-0.2(7)} = 3.69895$

$$10.0548 + 3.69895 = 13.754 \text{ mg}$$

\* (c)  $7.5 = De^{-0.2t}$       second dose is given 5 hours after the first dose

$$15e^{-0.2T} + 15e^{-0.2(5+T)} = 7.5$$

hours after the second dose is given  $7.5 = 15(e^{-0.2T} + e^{-0.2(T+5)})$

$$\frac{1}{2} = e^{-0.2T} + e^{-0.2(T+5)}$$

$$\frac{1}{2} = e^{-0.2T} + e^{-0.2T-1}$$

$$\frac{1}{2} = e^{-0.2T} + (e^{-0.2T} \times e^{-1})$$

$$\frac{1}{2} = e^{-0.2T}(1 + e^{-1})$$

$$(1 + e^{-1}) \quad (1 + e^{-1})$$

## Question 10 continued

$$e^{-0.2T} = \frac{(1 + e^{-1})}{2} \quad e^{-0.2T} = \frac{1}{2(1 + e^{-1})}$$

$$e^{0.2T} = 2(1 + e^{-1})$$

$$\ln e^{0.2T} = \ln(2 + 2e^{-1})$$

$$\frac{0.2T}{0.2} = \frac{\ln\left(2 + \frac{2}{e}\right)}{0.2} \quad T = 5 \ln\left(2 + \frac{2}{e}\right)$$