

## Exercise 4F

$$\begin{aligned}
 \mathbf{1\ a} \quad \text{LHS} &\equiv \frac{\cos 2A}{\cos A + \sin A} \\
 &\equiv \frac{\cos^2 A - \sin^2 A}{\cos A + \sin A} \\
 &\equiv \frac{(\cos A + \sin A)(\cos A - \sin A)}{\cos A + \sin A} \\
 &\equiv \cos A - \sin A \equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{b} \quad \text{LHS} &\equiv \frac{\sin B}{\sin A} - \frac{\cos B}{\cos A} \\
 &\equiv \frac{\sin B \cos A - \cos B \sin A}{\sin A \cos A} \\
 &\equiv \frac{\sin(B - A)}{\frac{1}{2}(2 \sin A \cos A)} \\
 &\equiv \frac{2 \sin(B - A)}{\sin 2A} \\
 &\equiv 2 \operatorname{cosec} 2A \sin(B - A) \equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{c} \quad \text{LHS} &\equiv \frac{1 - \cos 2\theta}{\sin 2\theta} \\
 &\equiv \frac{1 - (1 - 2 \sin^2 \theta)}{2 \sin \theta \cos \theta} \\
 &\equiv \frac{2 \sin^2 \theta}{2 \sin \theta \cos \theta} \\
 &\equiv \frac{\sin \theta}{\cos \theta} \\
 &\equiv \tan \theta \equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{d} \quad \text{LHS} &\equiv \frac{\sec^2 \theta}{1 - \tan^2 \theta} \\
 &\equiv \frac{1}{\cos^2 \theta (1 - \tan^2 \theta)} \\
 &\equiv \frac{1}{\cos^2 \theta - \sin^2 \theta} \quad \left( \text{as } \tan^2 \theta = \frac{\sin^2 \theta}{\cos^2 \theta} \right) \\
 &\equiv \frac{1}{\cos 2\theta} \\
 &\equiv \sec 2\theta \equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 1 \text{ e} \quad \text{LHS} &\equiv 2(\sin^3 \theta \cos \theta + \cos^3 \theta \sin \theta) \\
 &\equiv 2 \sin \theta \cos \theta (\sin^2 \theta + \cos^2 \theta) \\
 &\equiv \sin 2\theta \quad (\text{since } \sin^2 \theta + \cos^2 \theta \equiv 1) \\
 &\equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{f} \quad \text{LHS} &\equiv \frac{\sin 3\theta}{\sin \theta} - \frac{\cos 3\theta}{\cos \theta} \\
 &\equiv \frac{\sin 3\theta \cos \theta - \cos 3\theta \sin \theta}{\sin \theta \cos \theta} \\
 &\equiv \frac{\sin(3\theta - \theta)}{\frac{1}{2} \sin 2\theta} \\
 &\equiv \frac{\sin 2\theta}{\frac{1}{2} \sin 2\theta} \\
 &\equiv 2 \equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{g} \quad \text{LHS} &\equiv \operatorname{cosec} \theta - 2 \cot 2\theta \cos \theta \\
 &\equiv \operatorname{cosec} \theta - 2 \frac{\cos 2\theta}{\sin 2\theta} \cos \theta \\
 &\equiv \operatorname{cosec} \theta - \frac{2 \cos 2\theta \cos \theta}{2 \sin \theta \cos \theta} \\
 &\equiv \frac{1}{\sin \theta} - \frac{\cos 2\theta}{\sin \theta} \\
 &\equiv \frac{1 - \cos 2\theta}{\sin \theta} \\
 &\equiv \frac{1 - (1 - 2 \sin^2 \theta)}{\sin \theta} \\
 &\equiv \frac{2 \sin^2 \theta}{\sin \theta} \\
 &\equiv 2 \sin \theta \equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{h} \quad \text{LHS} &\equiv \frac{\sec \theta - 1}{\sec \theta + 1} \\
 &\equiv \frac{\frac{1}{\cos \theta} - 1}{\frac{1}{\cos \theta} + 1} \\
 &\equiv \frac{1 - \cos \theta}{1 + \cos \theta} \\
 &\equiv \frac{1 - (1 - 2 \sin^2 \frac{\theta}{2})}{1 + (2 \cos^2 \frac{\theta}{2} - 1)} \\
 &\equiv \frac{2 \sin^2 \frac{\theta}{2}}{2 \cos^2 \frac{\theta}{2}} \\
 &\equiv \tan^2 \frac{\theta}{2} \equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 1 \text{ i } \text{ LHS} &\equiv \tan\left(\frac{\pi}{4} - x\right) \\
 &\equiv \frac{\tan\frac{\pi}{4} - \tan x}{1 + \tan\frac{\pi}{4} \tan x} \\
 &\equiv \frac{1 - \tan x}{1 + \tan x} \\
 &\equiv \frac{1 - \frac{\sin x}{\cos x}}{1 + \frac{\sin x}{\cos x}} \\
 &\equiv \frac{\cos x - \sin x}{\cos x + \sin x} \\
 &\equiv \frac{\cos^2 x + \sin^2 x - 2 \sin x \cos x}{\cos^2 x - \sin^2 x} \quad (\text{multiply 'top and bottom' by } \cos x - \sin x) \\
 &\equiv \frac{1 - \sin 2x}{\cos 2x} \equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 2 \text{ a } \text{ LHS} &\equiv \sin(A + 60^\circ) + \sin(A - 60^\circ) \\
 &\equiv \sin A \cos 60^\circ + \cos A \sin 60^\circ + \sin A \cos 60^\circ - \cos A \sin 60^\circ \\
 &\equiv 2 \sin A \cos 60^\circ \\
 &\equiv \sin A \quad \left(\text{since } \cos 60^\circ = \frac{1}{2}\right) \\
 &\equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{b } \text{ LHS} &\equiv \frac{\cos A}{\sin B} - \frac{\sin A}{\cos B} \\
 &\equiv \frac{\cos A \cos B - \sin A \sin B}{\sin B \cos B} \\
 &\equiv \frac{\cos(A + B)}{\sin B \cos B} \\
 &\equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{c } \text{ LHS} &\equiv \frac{\sin(x + y)}{\cos x \cos y} \\
 &\equiv \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y} \\
 &\equiv \frac{\sin x \cos y}{\cos x \cos y} + \frac{\cos x \sin y}{\cos x \cos y} \\
 &\equiv \frac{\sin x}{\cos x} + \frac{\sin y}{\cos y} \\
 &\equiv \tan x + \tan y \\
 &\equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 2 \text{ d } \text{LHS} &\equiv \frac{\cos(x+y)}{\sin x \sin y} + 1 \\
 &\equiv \frac{\cos(x+y) + \sin x \sin y}{\sin x \sin y} \\
 &\equiv \frac{\cos x \cos y - \sin x \sin y + \sin x \sin y}{\sin x \sin y} \\
 &\equiv \frac{\cos x \cos y}{\sin x \sin y} \\
 &\equiv \cot x \cot y \\
 &\equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{e } \text{LHS} &\equiv \cos\left(\theta + \frac{\pi}{3}\right) + \sqrt{3} \sin \theta \\
 &\equiv \cos \theta \cos \frac{\pi}{3} - \sin \theta \sin \frac{\pi}{3} + \sqrt{3} \sin \theta \\
 &\equiv \frac{1}{2} \cos \theta - \frac{\sqrt{3}}{2} \sin \theta + \sqrt{3} \sin \theta \\
 &\equiv \frac{\sqrt{3}}{2} \sin \theta + \frac{1}{2} \cos \theta \\
 &\equiv \sin \theta \cos \frac{\pi}{6} + \cos \theta \sin \frac{\pi}{6} \quad \left( \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2}, \sin \frac{\pi}{6} = \frac{1}{2} \right) \\
 &\equiv \sin\left(\theta + \frac{\pi}{6}\right) \quad (\sin(A+B)) \\
 &\equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{f } \text{LHS} &\equiv \cot(A+B) \equiv \frac{\cos(A+B)}{\sin(A+B)} \\
 &\equiv \frac{\cos A \cos B - \sin A \sin B}{\sin A \cos B + \cos A \sin B} \\
 &\equiv \frac{\cos A \cos B}{\sin A \cos B} - \frac{\sin A \sin B}{\sin A \sin B} \quad (\text{dividing top and bottom by } \sin A \sin B) \\
 &\equiv \frac{\cot A \cot B - 1}{\cot A + \cot B} \equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 2 \text{ g } \text{LHS} &\equiv \sin^2(45^\circ + \theta) + \sin^2(45^\circ - \theta) \\
 &\equiv (\sin 45^\circ \cos \theta + \cos 45^\circ \sin \theta)^2 + (\sin 45^\circ \cos \theta - \cos 45^\circ \sin \theta)^2 \\
 &\equiv (\sin 45^\circ \cos \theta + \sin 45^\circ \sin \theta)^2 + (\sin 45^\circ \cos \theta - \sin 45^\circ \sin \theta)^2 \quad (\text{as } \sin 45^\circ = \cos 45^\circ) \\
 &\equiv (\sin 45^\circ)^2 \left( (\cos \theta + \sin \theta)^2 + (\cos \theta - \sin \theta)^2 \right) \\
 &\equiv \frac{1}{2} (\cos^2 \theta + 2 \sin \theta \cos \theta + \sin^2 \theta + \cos^2 \theta - 2 \sin \theta \cos \theta + \sin^2 \theta) \\
 &\equiv \frac{1}{2} (2(\sin^2 \theta + \cos^2 \theta)) \\
 &\equiv \frac{1}{2} \times 2 \quad (\sin^2 \theta + \cos^2 \theta \equiv 1) \\
 &\equiv 1 \\
 &\equiv \text{RHS}
 \end{aligned}$$

Alternatively as  $\sin(90^\circ - x^\circ) \equiv \cos x^\circ$ , if  $x = 45^\circ + \theta^\circ$  then  $\sin(45^\circ - \theta^\circ) \equiv \cos(45^\circ + \theta^\circ)$  and original LHS becomes  $\sin^2(45 + \theta)^\circ + \cos^2(45 + \theta)^\circ$ , which = 1

$$\begin{aligned}
 \text{h } \text{LHS} &\equiv \cos(A + B)\cos(A - B) \\
 &\equiv (\cos A \cos B - \sin A \sin B)(\cos A \cos B + \sin A \sin B) \\
 &\equiv \cos^2 A \cos^2 B - \sin^2 A \sin^2 B \\
 &\equiv \cos^2 A(1 - \sin^2 B) - (1 - \cos^2 A)\sin^2 B \\
 &\equiv \cos^2 A - \cos^2 A \sin^2 B - \sin^2 B + \cos^2 A \sin^2 B \\
 &\equiv \cos^2 A - \sin^2 B \\
 &\equiv \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 3 \text{ a } \text{LHS} &\equiv \tan \theta + \cot \theta \\
 &\equiv \frac{\sin \theta}{\cos \theta} + \frac{\cos \theta}{\sin \theta} \\
 &\equiv \frac{\sin^2 \theta + \cos^2 \theta}{\sin \theta \cos \theta} \\
 &\equiv \frac{2}{2 \sin \theta \cos \theta} \quad (\sin^2 \theta + \cos^2 \theta \equiv 1) \\
 &\equiv \frac{2}{\sin 2\theta} \\
 &\equiv 2 \operatorname{cosec} 2\theta \equiv \text{RHS}
 \end{aligned}$$

b Use  $\theta = 75^\circ$


$$\Rightarrow \tan 75^\circ + \cot 75^\circ = 2 \operatorname{cosec} 150^\circ = 2 \times \frac{1}{\sin 150^\circ} = 2 \times \frac{1}{\frac{1}{2}} = 4$$

$$\begin{aligned}
 4 \text{ a } \sin 3\theta &\equiv \sin(2\theta + \theta) \equiv \sin 2\theta \cos \theta + \cos 2\theta \sin \theta \\
 &\equiv (2\sin \theta \cos \theta) \cos \theta + (\cos^2 \theta - \sin^2 \theta) \sin \theta \\
 &\equiv 2\sin \theta \cos^2 \theta + \sin \theta \cos^2 \theta - \sin^3 \theta \\
 &\equiv 3\sin \theta \cos^2 \theta - \sin^3 \theta
 \end{aligned}$$

$$\begin{aligned}
 \text{b } \cos 3\theta &\equiv \cos(2\theta + \theta) \equiv \cos 2\theta \cos \theta - \sin 2\theta \sin \theta \\
 &\equiv (\cos^2 \theta - \sin^2 \theta) \cos \theta - (2\sin \theta \cos \theta) \sin \theta \\
 &\equiv \cos^3 \theta - \sin^2 \theta \cos \theta - 2\sin^2 \theta \cos \theta \\
 &\equiv \cos^3 \theta - 3\sin^2 \theta \cos \theta
 \end{aligned}$$

$$\begin{aligned}
 \text{c } \tan 3\theta &\equiv \frac{\sin 3\theta}{\cos 3\theta} \equiv \frac{3\sin \theta \cos^2 \theta - \sin^3 \theta}{\cos^3 \theta - 3\sin^2 \theta \cos \theta} \\
 &\equiv \frac{3\sin \theta \cos^2 \theta - \sin^3 \theta}{\cos^3 \theta} \\
 &\equiv \frac{\cos \theta}{\cos^3 \theta} \frac{3\sin \theta}{\cos^2 \theta} - \frac{\sin^3 \theta}{\cos^3 \theta} \\
 &\equiv \frac{3\sin \theta}{\cos^2 \theta} - \frac{\sin^3 \theta}{\cos^3 \theta} \\
 &\equiv \frac{3\sin \theta \cos \theta}{\cos^3 \theta} - \frac{\sin^3 \theta}{\cos^3 \theta} \\
 &\equiv \frac{3\sin \theta \cos \theta - \sin^3 \theta}{\cos^3 \theta} \\
 &\equiv \frac{3\tan \theta - \tan^3 \theta}{1 - 3\tan^2 \theta}
 \end{aligned}$$

d Sketch the right-angled triangle containing  $\theta$



This shows  $\tan \theta = 2\sqrt{2}$

$$\text{So } \tan 3\theta = \frac{3(2\sqrt{2}) - (2\sqrt{2})^3}{1 - 3(2\sqrt{2})^2} = \frac{6\sqrt{2} - 16\sqrt{2}}{1 - 24} = \frac{-10\sqrt{2}}{-23} = \frac{10\sqrt{2}}{23}$$

5 a i Using  $\cos 2A \equiv 2\cos^2 A - 1$  with  $A = \frac{x}{2}$

$$\Rightarrow \cos x \equiv 2\cos^2 \frac{x}{2} - 1$$

$$\Rightarrow 2\cos^2 \frac{x}{2} \equiv 1 + \cos x$$

$$\Rightarrow \cos^2 \frac{x}{2} \equiv \frac{1 + \cos x}{2}$$

5 a ii Using  $\cos 2A \equiv 1 - 2\sin^2 A$

$$\Rightarrow \cos x \equiv 1 - 2\sin^2 \frac{x}{2}$$

$$\Rightarrow 2\sin^2 \frac{x}{2} \equiv 1 - \cos x$$

$$\Rightarrow \sin^2 \frac{x}{2} \equiv \frac{1 - \cos x}{2}$$

b i Using (a) (i)  $\cos^2 \frac{\theta}{2} = \frac{1 + \cos \theta}{2} = \frac{1.6}{2} = 0.8 = \frac{4}{5}$

$$\Rightarrow \cos \frac{\theta}{2} = \frac{2}{\sqrt{5}} = \frac{2\sqrt{5}}{5} \quad \left( \text{as } \frac{\theta}{2} \text{ acute} \right)$$

ii Using (a) (ii)  $\sin^2 \frac{\theta}{2} = \frac{1 - \cos \theta}{2} = \frac{0.4}{2} = 0.2 = \frac{1}{5}$

$$\Rightarrow \sin \frac{\theta}{2} = \sqrt{\frac{1}{5}} = \frac{\sqrt{5}}{5}$$

iii  $\tan \frac{\theta}{2} = \frac{\sin \frac{\theta}{2}}{\cos \frac{\theta}{2}} = \frac{\frac{\sqrt{5}}{5}}{\frac{2}{\sqrt{5}}} = \frac{1}{2}$

c Using (a) (i) and squaring

$$\cos^4 \frac{A}{2} \equiv \left( \frac{1 + \cos A}{2} \right)^2 \equiv \frac{1 + 2\cos A + \cos^2 A}{4}$$

but using  $\cos 2A \equiv 2\cos^2 A - 1$  gives

$$\cos^2 A \equiv \frac{1}{2}(1 + \cos 2A)$$

$$\text{So } \cos^4 \frac{A}{2} \equiv \frac{1 + 2\cos A + \frac{1}{2}(1 + \cos 2A)}{4} \equiv \frac{2 + 4\cos A + 1 + \cos 2A}{8} \equiv \frac{3 + 4\cos A + \cos 2A}{8}$$

6 LHS  $\equiv \cos^4 \theta \equiv (\cos^2 \theta)^2 \equiv \left( \frac{1 + \cos 2\theta}{2} \right)^2$

$$\equiv \frac{1}{4}(1 + 2\cos 2\theta + \cos^2 2\theta)$$

$$\equiv \frac{1}{4} + \frac{1}{2}\cos 2\theta + \frac{1}{4}\left( \frac{1 + \cos 4\theta}{2} \right)$$

$$\equiv \frac{1}{4} + \frac{1}{2}\cos 2\theta + \frac{1}{8} + \frac{1}{8}\cos 4\theta$$

$$\equiv \frac{3}{8} + \frac{1}{2}\cos 2\theta + \frac{1}{8}\cos 4\theta \equiv \text{RHS}$$

$$\begin{aligned}
 7 \quad \sin^2(x+y) - \sin^2(x-y) &\equiv [\sin(x+y) + \sin(x-y)][\sin(x+y) - \sin(x-y)] \\
 &\equiv [\sin x \cos y + \cos x \sin y + \sin x \cos y - \cos x \sin y][\sin x \cos y + \cos x \sin y - (\sin x \cos y - \cos x \sin y)] \\
 &\equiv [2 \sin x \cos y][2 \cos x \sin y] \\
 &\equiv [2 \sin x \cos x][2 \cos y \sin y] \\
 &\equiv \sin 2x \sin 2y
 \end{aligned}$$

$$8 \quad \text{Let } \cos 2\theta - \sqrt{3} \sin 2\theta \equiv R \cos(2\theta + \alpha) \equiv R \cos 2\theta \cos \alpha - R \sin 2\theta \sin \alpha$$

$$\text{Compare } \cos 2\theta: R \cos \alpha = 1 \quad (1)$$

$$\text{Compare } \sin 2\theta: R \sin \alpha = \sqrt{3} \quad (2)$$

Divide (2) by (1):

$$\tan \alpha = \sqrt{3} \Rightarrow \alpha = \frac{\pi}{3}$$

Square and add equations:

$$R^2 = 1 + 3 = 4 \Rightarrow R = 2$$

$$\text{So } \cos 2\theta - \sqrt{3} \sin 2\theta \equiv 2 \cos\left(2\theta + \frac{\pi}{3}\right)$$

$$\begin{aligned}
 9 \quad 4 \cos\left(2\theta - \frac{\pi}{6}\right) &\equiv 4 \cos 2\theta \cos \frac{\pi}{6} + 4 \sin 2\theta \sin \frac{\pi}{6} \\
 &\equiv 2\sqrt{3} \cos 2\theta + 2 \sin 2\theta \\
 &\equiv 2\sqrt{3}(1 - 2 \sin^2 \theta) + 4 \sin \theta \cos \theta \\
 &\equiv 2\sqrt{3} - 4\sqrt{3} \sin^2 \theta + 4 \sin \theta \cos \theta
 \end{aligned}$$

$$\begin{aligned}
 10 \text{ a} \quad \text{RHS} &\equiv \sqrt{2} \sin\left(\theta + \frac{\pi}{4}\right) \\
 &\equiv \sqrt{2} \left( \sin \theta \cos \frac{\pi}{4} + \cos \theta \sin \frac{\pi}{4} \right) \\
 &\equiv \sqrt{2} \left( \sin \theta \times \frac{1}{\sqrt{2}} + \cos \theta \times \frac{1}{\sqrt{2}} \right) \\
 &\equiv \sin \theta + \cos \theta \\
 &\equiv \text{LHS}
 \end{aligned}$$

$$\begin{aligned}
 \text{b} \quad \text{RHS} &\equiv 2 \sin\left(2\theta - \frac{\pi}{6}\right) \\
 &\equiv 2 \left( \sin 2\theta \cos \frac{\pi}{6} - \cos 2\theta \sin \frac{\pi}{6} \right) \\
 &\equiv 2 \left( \sin 2\theta \times \frac{\sqrt{3}}{2} - \cos 2\theta \times \frac{1}{2} \right) \\
 &\equiv \sqrt{3} \sin 2\theta - \cos 2\theta \\
 &\equiv \text{LHS}
 \end{aligned}$$



## Challenge

$$1 \text{ a } \cos(A+B) - \cos(A-B) \equiv \cos A \cos B - \sin A \sin B - (\cos A \cos B + \sin A \sin B) \\ \equiv -2 \sin A \sin B$$

$$b \text{ Let } A+B=P \text{ and } A-B=Q$$

Solving simultaneously gives

$$2A = P+Q$$

$$A = \frac{P+Q}{2}$$

and

$$2B = P-Q$$

$$B = \frac{P-Q}{2}$$

Substituting these into the identity from part a gives

$$\cos P - \cos Q \equiv -2 \sin\left(\frac{P+Q}{2}\right) \sin\left(\frac{P-Q}{2}\right)$$

$$c \text{ Rearranging the identity from part a to give } \sin A \sin B \equiv -\frac{1}{2} \cos(A+B) + \frac{1}{2} \cos(A-B)$$

$$3 \sin x \sin 7x \equiv -\frac{3}{2} \cos(x+7x) + \frac{3}{2} \cos(x-7x) \\ \equiv -\frac{3}{2} \cos 8x + \frac{3}{2} \cos(-6x) \\ \equiv -\frac{3}{2} \cos 8x + \frac{3}{2} \cos(6x) \quad (\text{as } \cos(-x) \equiv \cos x) \\ \equiv -\frac{3}{2} (\cos 8x - \cos 6x)$$

$$2 \text{ a } \sin(A+B) + \sin(A-B) \equiv \sin A \cos B + \cos A \sin B + (\sin A \cos B - \cos A \sin B) \\ \equiv 2 \sin A \cos B$$

$$\text{Let } A+B=P \text{ and } A-B=Q$$

Solving simultaneously gives

$$2A = P+Q$$

$$A = \frac{P+Q}{2}$$

and

$$2B = P-Q$$

$$B = \frac{P-Q}{2}$$

Substituting these into the equation for  $\sin(A+B) + \sin(A-B)$  gives

$$\sin P + \sin Q \equiv 2 \sin\left(\frac{P+Q}{2}\right) \cos\left(\frac{P-Q}{2}\right)$$

$$2 \text{ b Let } \frac{11\pi}{24} = \frac{P+Q}{2}, \frac{5\pi}{24} = \frac{P-Q}{2}$$
$$\frac{22\pi}{24} = P+Q, \frac{10\pi}{24} = P-Q$$

Solving simultaneously gives:

$$2P = \frac{32\pi}{24}, P = \frac{16\pi}{24}$$

and

$$2Q = \frac{12\pi}{24}, Q = \frac{6\pi}{24}$$

$$\text{So } 2 \sin \frac{11\pi}{24} \cos \frac{5\pi}{24} = \sin \left( \frac{16\pi}{24} \right) + \sin \left( \frac{6\pi}{24} \right) = \sin \left( \frac{2\pi}{3} \right) + \sin \left( \frac{\pi}{4} \right) = \frac{\sqrt{3}}{2} + \frac{\sqrt{2}}{2} = \frac{\sqrt{3} + \sqrt{2}}{2}$$