

C3

NUMERICAL METHODS

Worksheet B

- 1 For each equation, show that it can be rearranged into the given iterative form. Use this and the given value of x_0 to find x_1 , x_2 and x_3 . Give your value of x_3 correct to 4 decimal places.
- a $9 + 4x - 2x^3 = 0$ $x_{n+1} = \sqrt[3]{2x_n + 4.5}$ $x_0 = 2$
- b $e^x - 8x + 5 = 0$ $x_{n+1} = \ln(8x_n - 5)$ $x_0 = 3$
- c $\tan x - 5x + 13 = 0$ $x_{n+1} = \arctan(5x_n - 13)$ $x_0 = -1.2$
- d $\ln x + \sqrt{x} + 1.4 = 0$ $x_{n+1} = e^{-(\sqrt{x_n} + 1.4)}$ $x_0 = 0.16$
- 2 For each equation, show that it can be rearranged into the given iterative form and state the values of the constants a and b . Use this and the given value of x_0 to find x_1 , x_2 and x_3 . Give your value of x_3 correct to 3 decimal places.
- a $e^{2x-1} - 6x = 0$ $x_{n+1} = a(\ln bx_n + 1)$ $x_0 = 1.7$
- b $\frac{2}{x} + \cos x - 3 = 0$ $x_{n+1} = \frac{a}{b - \cos x_n}$ $x_0 = 0.8$
- c $2x^3 - 6x - 11 = 0$ $x_{n+1} = \sqrt{a + \frac{b}{x_n}}$ $x_0 = 2$
- d $15 \ln(x + 3) - 4x = 0$ $x_{n+1} = e^{ax_n} + b$ $x_0 = -2.5$
- 3 In each case, use the given iteration formula and value of x_0 to find a root of the equation $f(x) = 0$ to the stated degree of accuracy. Justify the accuracy of your answers.
- a $f(x) = 10^x + 3x - 4$ $x_{n+1} = \log_{10}(4 - 3x_n)$ $x_0 = 0.44$ 3 decimal places
- b $f(x) = x^2 + \frac{1}{x-5}$ $x_{n+1} = \sqrt{\frac{x_n^3 + 1}{5}}$ $x_0 = 0.5$ 2 significant figures
- c $f(x) = 30 - 5x + \sin 2x$ $x_{n+1} = 6 + 0.2 \sin 2x_n$ $x_0 = 6$ 3 significant figures
- d $f(x) = e^{4-x} - \ln x$ $x_{n+1} = 4 - \ln(\ln x_n)$ $x_0 = 3.7$ 3 decimal places
- 4 $f(x) = x^5 - 10x^3 + 4$.
- The equation $f(x) = 0$ has a root in the interval $-4 < x < -3$.
- a Use the iteration formula $x_{n+1} = \sqrt[5]{10x_n^3 - 4}$ and the starting value $x_0 = -3.2$ to find the value of this root correct to 2 decimal places.
- The equation $f(x) = 0$ can be rearranged into the iterative form $x_{n+1} = \sqrt[3]{\frac{a}{b-x_n^2}}$.
- b Find the values of the constants a and b in this formula.
- The equation $f(x) = 0$ has another root in the interval $0 < x < 1$.
- c Using the iteration formula with your values from part **b** and the starting value $x_0 = 1$, find the value of this root correct to 3 decimal places.
- 5 $f: x \rightarrow \arcsin 2x - 0.5x - 0.7$, $x \in \mathbb{R}$, $|x| \leq 0.5$
- The equation $f(x) = 0$ can be rearranged into the iterative form $x_{n+1} = a \sin(bx_n + c)$.
- a Find the values of the constants a , b and c in this formula.
- The equation $f(x) = 0$ has a solution in the interval $(0.3, 0.4)$.
- b Using the iterative formula with your values from part **a** and the starting value $x_0 = 0.4$, find this solution correct to 3 decimal places.