

**C3****NUMERICAL METHODS****Answers - Worksheet D**

**1**    **a**  $\frac{dy}{dx} = e^x + 2x$

**b** at A,  $x = 0 \therefore y = -3$ , grad = 1

$$\therefore y = x - 3$$

**c** SP:  $e^x + 2x = 0$

let  $f(x) = e^x + 2x$

$$f(-0.4) = -0.130$$

$$f(-0.3) = 0.141$$

sign change,  $f(x)$  continuous  $\therefore$  root

$\therefore$  x-coord of B in interval  $[-0.4, -0.3]$

**d**  $x_1 = -0.34694$

$$x_2 = -0.35126$$

$$x_3 = -0.35169$$

$$x_4 = -0.35173$$

$\therefore$  x-coord of B = -0.352 (3dp)

**2**    **a**  $f(0) = 0.279$

$$f(5) = -4.10$$

$$f(1) = 0.266$$

$$f(3) = -2.44$$

$$f(2) = -0.853$$

$$\therefore k = 1$$

**b**  $x_0 = 1$

$$x_1 = 1.2684$$

$$x_2 = 1.3106$$

$$x_3 = 1.3106$$

**3**    **a** area of segment =  $\frac{1}{2}r^2\theta - \frac{1}{2}r^2\sin \theta$

$$= \frac{1}{2}r^2(\theta - \sin \theta)$$

$$\therefore \frac{1}{2}r^2\sin \theta = 4 \times \frac{1}{2}r^2(\theta - \sin \theta)$$

$$\sin \theta = 4(\theta - \sin \theta)$$

$$\sin \theta = 4\theta - 4 \sin \theta$$

$$4\theta - 5 \sin \theta = 0$$

**b**  $\theta_1 = 1.11401$

$$\theta_2 = 1.12184$$

$$\theta_3 = 1.12613$$

$$\theta_4 = 1.12844$$

$$\theta_5 = 1.12968$$

$$\therefore \theta = 1.13 \text{ (2dp)}$$

**4**    **a**  $e^{x^2} - x - 3 = 0$

$$e^{x^2} = x + 3$$

$$x^2 = \ln(x + 3)$$

$$x = \sqrt{\ln(x + 3)} \quad \therefore a = 1, b = 3$$

**b** e.g.  $x_0 = 1.5$

$$x_1 = 1.226408$$

$$x_2 = 1.200563$$


$$x_3 = 1.198006$$

$$x_4 = 1.197752$$

$$x_5 = 1.197727$$

$$\therefore \text{solution} = 1.198 \text{ (3dp)}$$

5    a  $y = x^2 - 9$   
 swap  $x = y^2 - 9$   
 $y = \pm\sqrt{x+9}$   
 (domain  $\Rightarrow +$ )  
 $f^{-1}(x) = \sqrt{x+9}, x \in \mathbb{R}, x \geq -9$   
 range:  $f^{-1}(x) \geq 0$




c let  $h(x) = f^{-1}(x) + g(x) = \sqrt{x+9} + x^3$   
 $h(-2) = -5.35$   
 $h(-1) = 1.83$   
 sign change,  $h(x)$  continuous  $\therefore$  root  
 d  $x_1 = -1.41421, x_2 = -1.40174,$   
 $x_3 = -1.40212, x_4 = -1.40211$   
 $\therefore$  root = -1.402 (3dp)

7    a at  $A, x^{\frac{5}{2}} - 3x^{\frac{1}{2}} - 7x = 0$

let  $f(x) = x^{\frac{5}{2}} - 3x^{\frac{1}{2}} - 7x$   
 $f(4) = -2, f(5) = 14.2$   
 sign change,  $f(x)$  continuous  $\therefore$  root  
 $\therefore 4 < \alpha < 5$

b  $\frac{dy}{dx} = \frac{5}{2}x^{\frac{3}{2}} - \frac{3}{2}x^{-\frac{1}{2}} - 7$   
 at  $B, \frac{5}{2}x^{\frac{3}{2}} - \frac{3}{2}x^{-\frac{1}{2}} - 7 = 0$   
 let  $g(x) = \frac{5}{2}x^{\frac{3}{2}} - \frac{3}{2}x^{-\frac{1}{2}} - 7$   
 $g(2) = -0.990, g(3) = 5.12$   
 sign change,  $g(x)$  continuous  $\therefore$  root  
 $\therefore 2 < \beta < 3$

c  $\frac{5}{2}x^{\frac{3}{2}} - \frac{3}{2}x^{-\frac{1}{2}} - 7 = 0$   
 $5x^2 - 3 - 14x^{\frac{1}{2}} = 0$   
 $x^2 = 0.6 + 2.8x^{\frac{1}{2}}$   
 $x > 0 \therefore x = \beta$  is a soln to  $x = \sqrt{0.6 + 2.8x^{\frac{1}{2}}}$   
 d  $x_1 = 2.158144$   
 $x_2 = 2.171031$   
 $x_3 = 2.173853$   
 $x_4 = 2.174470$   
 $x_5 = 2.174604$   
 $\therefore \beta = 2.175$  (4sf)



b  $-(-x^2 - 3x) = \frac{1}{x}$   
 $x^2 + 3x = \frac{1}{x}$   
 $x^3 + 3x^2 = 1$   
 $x^3 + 3x^2 - 1 = 0$   
 c  $x_1 = 0.57735$   
 $x_2 = 0.52871$   
 $x_3 = 0.53234$   
 $x_4 = 0.53207$   
 $\therefore$  x-coord of  $P = 0.532$  (3dp)

8    a  $\frac{dy}{dx} = 3 - \frac{1}{x}$

grad = 2  
 $\therefore$  grad of normal =  $-\frac{1}{2}$   
 $y - 3 = -\frac{1}{2}(x - 1)$   
 $[y = \frac{7}{2} - \frac{1}{2}x]$

b  $3x - \ln x = \frac{7}{2} - \frac{1}{2}x$

$6x - 2\ln x = 7 - x$

$2\ln x - 7x + 7 = 0$

c  $2\ln x = 7x - 7$   
 $\ln x = \frac{7}{2}(x - 1)$   
 $x = e^{\frac{7}{2}(x-1)} \therefore k = \frac{7}{2}$

d  $x_1 = 0.173774$   
 $x_2 = 0.055477$   
 $x_3 = 0.036669$   
 $x_4 = 0.034333$   
 $x_5 = 0.034053$   
 $\therefore$  x-coord of  $Q = 0.034$  (3dp)  
 e let  $f(x) = 2\ln x - 7x + 7$   
 $f(0.0335) = -0.027$   
 $f(0.0345) = 0.025$   
 sign change,  $f(x)$  continuous  $\therefore$  root