

# C3 TRIGONOMETRY

# Worksheet C

- 1** Use the identity  $\sin^2 x + \cos^2 x \equiv 1$  to obtain the identities
- a**  $1 + \tan^2 x \equiv \sec^2 x$
- b**  $1 + \cot^2 x \equiv \operatorname{cosec}^2 x$
- 2** **a** Given that  $\tan A = \frac{1}{3}$ , find the exact value of  $\sec^2 A$ .
- b** Given that  $\operatorname{cosec} B = 1 + \sqrt{3}$ , find the exact value of  $\cot^2 B$ .
- c** Given that  $\sec C = \frac{3}{2}$ , find the possible values of  $\tan C$ , giving your answers in the form  $k\sqrt{5}$ .
- 3** Solve each equation for  $\theta$  in the interval  $0 \leq \theta \leq 2\pi$  giving your answers in terms of  $\pi$ .
- a**  $3 \sec^2 \theta = 4 \tan^2 \theta$
- b**  $\tan^2 \theta - 2 \sec \theta + 1 = 0$
- c**  $\cot^2 \theta - 3 \operatorname{cosec} \theta + 3 = 0$
- d**  $\operatorname{cosec}^2 \theta + \cot^2 \theta = 3$
- e**  $\sec^2 \theta + 2 \tan \theta = 0$
- f**  $\operatorname{cosec}^2 \theta - \sqrt{3} \cot \theta - 1 = 0$
- 4** Solve each equation for  $x$  in the interval  $-180^\circ \leq x \leq 180^\circ$ .  
Give your answers to 1 decimal place where appropriate.
- a**  $\tan^2 x - 2 \sec x - 2 = 0$
- b**  $2 \operatorname{cosec}^2 x + 2 = 9 \cot x$
- c**  $\operatorname{cosec}^2 x + 5 \operatorname{cosec} x + 2 \cot^2 x = 0$
- d**  $3 \tan^2 x - 3 \tan x + \sec^2 x = 2$
- e**  $\tan^2 x + 4 \sec x - 2 = 0$
- f**  $2 \cot^2 x + 3 \operatorname{cosec}^2 x = 4 \cot x + 3$
- 5** Solve each equation for  $x$  in the interval  $0 \leq x \leq 360^\circ$ .
- a**  $\cot^2 2x + \operatorname{cosec} 2x - 1 = 0$
- b**  $8 \sin^2 x + \sec x = 8$
- c**  $3 \operatorname{cosec}^2 x - 4 \sin^2 x = 1$
- d**  $9 \sec^2 x - 8 = \operatorname{cosec}^2 x$
- 6** Prove each of the following identities.
- a**  $\operatorname{cosec}^2 x - \sec^2 x \equiv \cot^2 x - \tan^2 x$
- b**  $(\cot x - 1)^2 \equiv \operatorname{cosec}^2 x - 2 \cot x$
- c**  $(\cos x - 2 \sec x)^2 \equiv \cos^2 x + 4 \tan^2 x$
- d**  $\sec^2 x - \sin^2 x \equiv \tan^2 x + \cos^2 x$
- e**  $(\tan x + \cot x)^2 \equiv \sec^2 x + \operatorname{cosec}^2 x$
- f**  $(\sin x - \sec x)^2 \equiv \sin^2 x + (\tan x - 1)^2$
- g**  $\sec^2 x + \operatorname{cosec}^2 x \equiv \sec^2 x \operatorname{cosec}^2 x$
- h**  $\sec^4 x + \tan^4 x \equiv 2 \sec^2 x \tan^2 x + 1$
- 7** Prove that there are no real values of  $x$  for which  

$$4 \sec^2 x - \sec x + 2 \tan^2 x = 0.$$
- 8** **a** Prove the identity  

$$\operatorname{cosec} x \sec x - \cot x \equiv \tan x.$$
- b** Hence, or otherwise, find the values of  $x$  in the interval  $0 \leq x \leq 360^\circ$  for which  

$$\operatorname{cosec} x \sec x = 3 + \cot x,$$
  
giving your answers to 1 decimal place.