

**Exercise 4B**

**1 a i**  $\sqrt{(4+2x)}$  Write in index form.

$$= (4+2x)^{\frac{1}{2}} \text{ Take out a factor of 4}$$

$$= \left( 4 \left( 1 + \frac{2x}{4} \right) \right)^{\frac{1}{2}} \text{ Remember to put the 4 to the power } \frac{1}{2}$$

$$= 4^{\frac{1}{2}} \left( 1 + \frac{x}{2} \right)^{\frac{1}{2}} \quad 4^{\frac{1}{2}} = 2$$

$$= 2 \left( 1 + \frac{x}{2} \right)^{\frac{1}{2}} \text{ Use the expansion with } n = \frac{1}{2} \text{ and } x = \frac{x}{2}$$

$$= 2 \left( 1 + \left( \frac{1}{2} \right) \left( \frac{x}{2} \right) + \frac{\left( \frac{1}{2} \right) \left( -\frac{1}{2} \right) \left( \frac{x}{2} \right)^2}{2!} + \frac{\left( \frac{1}{2} \right) \left( -\frac{1}{2} \right) \left( -\frac{3}{2} \right) \left( \frac{x}{2} \right)^3}{3!} + \dots \right)$$

$$= 2 \left( 1 + \frac{x}{4} - \frac{x^2}{32} + \frac{x^3}{128} + \dots \right) \text{ Multiply by the 2}$$

$$= 2 + \frac{x}{2} - \frac{x^2}{16} + \frac{x^3}{64} + \dots$$

**ii** Valid if  $\left| \frac{x}{2} \right| < 1 \Rightarrow |x| < 2$

**b i**  $\frac{1}{2+x}$  Write in index form

$$= (2+x)^{-1} \text{ Take out a factor of 2}$$

$$= \left( 2 \left( 1 + \frac{x}{2} \right) \right)^{-1} \text{ Remember to put 2 to the power } -1$$

$$= 2^{-1} \left( 1 + \frac{x}{2} \right)^{-1}, \quad 2^{-1} = \frac{1}{2}. \text{ Use the binomial expansion with } n = -1 \text{ and } x = \frac{x}{2}$$

$$= \frac{1}{2} \left( 1 + (-1) \left( \frac{x}{2} \right) + \frac{(-1)(-2)}{2!} \left( \frac{x}{2} \right)^2 + \frac{(-1)(-2)(-3)}{3!} \left( \frac{x}{2} \right)^3 + \dots \right)$$

$$= \frac{1}{2} \left( 1 - \frac{x}{2} + \frac{x^2}{4} - \frac{x^3}{8} + \dots \right) \text{ Multiply by the } \frac{1}{2}$$

$$= \frac{1}{2} - \frac{x}{4} + \frac{x^2}{8} - \frac{x^3}{16} + \dots$$

**ii** Valid if  $\left| \frac{x}{2} \right| < 1 \Rightarrow |x| < 2$

**1 c i**  $\frac{1}{(4-x)^2}$  Write in index form

$$\begin{aligned}
 &= (4-x)^{-2} \quad \text{Take 4 out as a factor} \\
 &= \left( 4 \left( 1 - \frac{x}{4} \right) \right)^{-2} \\
 &= 4^{-2} \left( 1 - \frac{x}{4} \right)^{-2}, \quad 4^{-2} = \frac{1}{16}. \text{ Use the binomial expansion with } n = -2 \text{ and } x = -\frac{x}{4} \\
 &= \frac{1}{16} \left( 1 + (-2) \left( -\frac{x}{4} \right) + \frac{(-2)(-3)}{2!} \left( -\frac{x}{4} \right)^2 + \frac{(-2)(-3)(-4)}{3!} \left( -\frac{x}{4} \right)^3 + \dots \right) \\
 &= \frac{1}{16} \left( 1 + \frac{x}{2} + \frac{3x^2}{16} + \frac{x^3}{16} + \dots \right) \quad \text{Multiply by } \frac{1}{16} \\
 &= \frac{1}{16} + \frac{x}{32} + \frac{3x^2}{256} + \frac{x^3}{256} + \dots
 \end{aligned}$$

**ii** Valid if  $\left| \frac{x}{4} \right| < 1 \Rightarrow |x| < 4$

**d i**  $\sqrt{9+x}$  Write in index form

$$\begin{aligned}
 &= (9+x)^{\frac{1}{2}} \quad \text{Take 9 out as a factor} \\
 &= \left( 9 \left( 1 + \frac{x}{9} \right) \right)^{\frac{1}{2}} \\
 &= 9^{\frac{1}{2}} \left( 1 + \frac{x}{9} \right)^{\frac{1}{2}}, \quad 9^{\frac{1}{2}} = 3. \text{ Use binomial expansion with } n = \frac{1}{2} \text{ and } x = \frac{x}{9} \\
 &= 3 \left( 1 + \left( \frac{1}{2} \right) \left( \frac{x}{9} \right) + \frac{\left( \frac{1}{2} \right) \left( -\frac{1}{2} \right)}{2!} \left( \frac{x}{9} \right)^2 + \frac{\left( \frac{1}{2} \right) \left( -\frac{1}{2} \right) \left( -\frac{3}{2} \right)}{3!} \left( \frac{x}{9} \right)^3 + \dots \right) \\
 &= 3 \left( 1 + \frac{x}{18} - \frac{x^2}{648} + \frac{x^3}{11664} + \dots \right) \quad \text{Multiply by 3} \\
 &= 3 + \frac{x}{6} - \frac{x^2}{216} + \frac{x^3}{3888} + \dots
 \end{aligned}$$

**ii** Valid for  $\left| \frac{x}{9} \right| < 1 \Rightarrow |x| < 9$

**1 e i**  $\frac{1}{\sqrt{2+x}}$  Write in index form

$$= (2+x)^{-\frac{1}{2}} \text{ Take out a factor of 2}$$

$$= \left( 2 \left( 1 + \frac{x}{2} \right) \right)^{-\frac{1}{2}}$$

$$= 2^{-\frac{1}{2}} \left( 1 + \frac{x}{2} \right)^{-\frac{1}{2}}, \quad 2^{-\frac{1}{2}} = \frac{1}{2^{\frac{1}{2}}} = \frac{1}{\sqrt{2}}. \text{ Use binomial expansion with } n = -\frac{1}{2} \text{ and } x = \frac{x}{2}$$

$$= \frac{1}{\sqrt{2}} \left( 1 + \left( -\frac{1}{2} \right) \left( \frac{x}{2} \right) + \frac{\left( -\frac{1}{2} \right) \left( -\frac{3}{2} \right)}{2!} \left( \frac{x}{2} \right)^2 + \frac{\left( -\frac{1}{2} \right) \left( -\frac{3}{2} \right) \left( -\frac{5}{2} \right)}{3!} \left( \frac{x}{2} \right)^3 + \dots \right)$$

$$= \frac{1}{\sqrt{2}} \left( 1 - \frac{x}{4} + \frac{3x^2}{32} - \frac{5x^3}{128} + \dots \right) \text{ Multiply by } \frac{1}{\sqrt{2}}$$

$$= \frac{1}{\sqrt{2}} - \frac{x}{4\sqrt{2}} + \frac{3x^2}{32\sqrt{2}} - \frac{5x^3}{128\sqrt{2}} + \dots \text{ Rationalise surds}$$

$$= \frac{\sqrt{2}}{2} - \frac{\sqrt{2}x}{8} + \frac{3\sqrt{2}x^2}{64} - \frac{5\sqrt{2}x^3}{256} + \dots$$

**ii** Valid if  $\left| \frac{x}{2} \right| < 1 \Rightarrow |x| < 2$

**f i**  $\frac{5}{3+2x}$  Write in index form

$$= 5(3+2x)^{-1} \text{ Take out a factor of 3}$$

$$= 5 \left( 3 \left( 1 + \frac{2x}{3} \right) \right)^{-1}$$

$$= 5 \times 3^{-1} \left( 1 + \frac{2x}{3} \right)^{-1}, \quad 3^{-1} = \frac{1}{3}. \text{ Use binomial expansion with } n = -1 \text{ and } x = \frac{2x}{3}$$

$$= \frac{5}{3} \left( 1 + (-1) \left( \frac{2x}{3} \right) + \frac{(-1)(-2)}{2!} \left( \frac{2x}{3} \right)^2 + \frac{(-1)(-2)(-3)}{3!} \left( \frac{2x}{3} \right)^3 + \dots \right)$$

$$= \frac{5}{3} \left( 1 - \frac{2x}{3} + \frac{4x^2}{9} - \frac{8x^3}{27} + \dots \right) \text{ Multiply by } \frac{5}{3}$$

$$= \frac{5}{3} - \frac{10x}{9} + \frac{20x^2}{27} - \frac{40x^3}{81} + \dots$$

**ii** Valid if  $\left| \frac{2x}{3} \right| < 1 \Rightarrow |x| < \frac{3}{2}$

**1 g i**

$$\begin{aligned}
 & \frac{1+x}{2+x} = 1 - \frac{1}{2+x} \quad \text{Write } \frac{1}{2+x} \text{ in index form} \\
 &= 1 - (2+x)^{-1} \quad \text{Take out a factor of 2} \\
 &= 1 - \left( 2 \left( 1 + \frac{x}{2} \right) \right)^{-1} \\
 &= 1 - \left( 2^{-1} \left( 1 + \frac{x}{2} \right)^{-1} \right) \quad \text{Expand } \left( 1 + \frac{x}{2} \right)^{-1} \text{ using the binomial expansion} \\
 &= 1 - \left( \frac{1}{2} \left( 1 + (-1) \left( \frac{x}{2} \right) + \frac{(-1)(-2)}{2!} \left( \frac{x}{2} \right)^2 + \frac{(-1)(-2)(-3)}{3!} \left( \frac{x}{2} \right)^3 + \dots \right) \right) \\
 &= 1 - \left( \frac{1}{2} \left( 1 - \frac{x}{2} + \frac{x^2}{4} - \frac{x^3}{8} + \dots \right) \right) \quad \text{Multiply } \left( 1 - \frac{x}{2} + \frac{x^2}{4} - \frac{x^3}{8} + \dots \right) \text{ by } \frac{1}{2} \\
 &= 1 - \left( \frac{1}{2} - \frac{x}{4} + \frac{x^2}{8} - \frac{x^3}{16} + \dots \right) \\
 &= \frac{1}{2} + \frac{1}{4}x - \frac{1}{8}x^2 + \frac{1}{16}x^3 + \dots
 \end{aligned}$$

**ii** Valid for  $\left| \frac{x}{2} \right| < 1 \Rightarrow |x| < 2$

**1 h i**  $\sqrt{\frac{2+x}{1-x}}$

$$= (2+x)^{\frac{1}{2}} (1-x)^{-\frac{1}{2}} \quad \text{Put both in index form}$$

$$= 2^{\frac{1}{2}} \left(1 + \frac{x}{2}\right)^{\frac{1}{2}} (1-x)^{-\frac{1}{2}} \quad \text{Expand both using the binomial expansion}$$

$$= \sqrt{2} \left( 1 + \left(\frac{1}{2}\right) \left(\frac{x}{2}\right) + \frac{\left(\frac{1}{2}\right) \left(-\frac{1}{2}\right)}{2!} \left(\frac{x}{2}\right)^2 + \frac{\left(\frac{1}{2}\right) \left(-\frac{1}{2}\right) \left(-\frac{3}{2}\right)}{3!} \left(\frac{x}{2}\right)^3 + \dots \right)$$

$$\times \left( 1 + \left(-\frac{1}{2}\right) (-x) + \frac{\left(-\frac{1}{2}\right) \left(-\frac{3}{2}\right)}{2!} (-x)^2 + \frac{\left(-\frac{1}{2}\right) \left(-\frac{3}{2}\right) \left(-\frac{5}{2}\right)}{3!} (-x)^3 + \dots \right)$$

$$= \sqrt{2} \left( 1 + \frac{1}{4}x - \frac{1}{32}x^2 + \frac{1}{128}x^3 + \dots \right) \left( 1 + \frac{1}{2}x + \frac{3}{8}x^2 + \frac{5}{16}x^3 + \dots \right) \quad \text{Multiply out}$$

$$= \sqrt{2} \left( 1 \left( 1 + \frac{1}{2}x + \frac{3}{8}x^2 + \frac{15}{16}x^3 + \dots \right) + \frac{1}{4}x \left( 1 + \frac{1}{2}x + \frac{3}{8}x^2 + \frac{5}{16}x^3 + \dots \right) \right. \\ \left. - \frac{1}{32}x^2 \left( 1 + \frac{1}{2}x + \frac{3}{8}x^2 + \frac{5}{16}x^3 + \dots \right) + \frac{1}{128}x^3 \left( 1 + \frac{1}{2}x + \frac{3}{8}x^2 + \frac{5}{16}x^3 + \dots \right) + \dots \right)$$

$$= \sqrt{2} \left( 1 + \frac{1}{2}x + \frac{3}{8}x^2 + \frac{5}{16}x^3 + \frac{1}{4}x + \frac{1}{8}x^2 \right. \\ \left. + \frac{3}{32}x^3 - \frac{1}{32}x^2 - \frac{1}{64}x^3 + \frac{1}{128}x^3 + \dots \right)$$

Collect like terms

$$= \sqrt{2} \left( 1 + \frac{3}{4}x + \frac{15}{32}x^2 + \frac{51}{128}x^3 + \dots \right) \quad \text{Multiply by } \sqrt{2}$$

$$= \sqrt{2} + \frac{3\sqrt{2}}{4}x + \frac{15\sqrt{2}}{32}x^2 + \frac{51\sqrt{2}}{128}x^3 + \dots$$

**ii** Valid if  $\left|\frac{x}{2}\right| < 1$  and  $|x| < 1 \Rightarrow |x| < 1$  for both to be valid.

$$\begin{aligned}
 \mathbf{2} \quad (5+4x)^{-2} &= \left(5\left(1+\frac{4}{5}x\right)\right)^{-2} = 5^{-2}\left(1+\frac{4}{5}x\right)^{-2} = \frac{1}{25}\left(1+\frac{4}{5}x\right)^{-2} \\
 &= \frac{1}{25}\left(1+(-2)\left(\frac{4}{5}x\right) + \frac{(-2)(-3)}{2!}\left(\frac{4}{5}x\right)^2 + \frac{(-2)(-3)(-4)}{3!}\left(\frac{4}{5}x\right)^3 + \dots\right) \\
 &= \frac{1}{25}\left(1+(-2)\left(\frac{4}{5}x\right) + \frac{(-2)(-3)}{2}\frac{16}{25}x^2 + \frac{(-2)(-3)(-4)}{6}\frac{64}{125}x^3 + \dots\right) \\
 &= \frac{1}{25}\left(1-\frac{8}{5}x + \frac{48}{25}x^2 - \frac{256}{125}x^3 + \dots\right) \\
 &= \frac{1}{25} - \frac{8}{125}x + \frac{48}{625}x^2 - \frac{256}{3125}x^3 + \dots
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{3} \quad \mathbf{a} \quad \sqrt{(4-x)} &= (4-x)^{\frac{1}{2}} \\
 &= \left[4\left(1-\frac{x}{4}\right)\right]^{\frac{1}{2}} \\
 &= 4^{\frac{1}{2}}\left(1-\frac{x}{4}\right)^{\frac{1}{2}} \\
 &= 2\left[1+\left(\frac{1}{2}\right)\left(-\frac{x}{4}\right) + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)}{2!}\left(-\frac{x}{4}\right)^2 + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{3!}\left(-\frac{x}{4}\right)^3 + \dots\right] \\
 &= 2\left(1-\frac{x}{8}-\frac{x^2}{128}-\frac{x^3}{1024}+\dots\right) \\
 &= 2-\frac{x}{4}-\frac{x^2}{64}-\frac{x^3}{512}+\dots
 \end{aligned}$$

Valid for  $\left|-\frac{x}{4}\right| < 1 \Rightarrow |x| < 4$

$$\mathbf{b} \quad m\left(\frac{1}{9}\right) = \sqrt{4-\frac{1}{9}} = \sqrt{\frac{35}{9}} = \frac{\sqrt{35}}{3}$$

**3 c**  $m(x) \approx 2 - \frac{1}{4}x - \frac{1}{64}x^2$

$$m\left(\frac{1}{9}\right) = \frac{\sqrt{35}}{3}$$

$$\sqrt{35} = 3m\left(\frac{1}{9}\right)$$

$$\approx 3\left(2 - \frac{1}{4}\left(\frac{1}{9}\right) - \frac{1}{64}\left(\frac{1}{9}\right)^2\right)$$

$$\approx 3\left(2 - \frac{1}{36} - \frac{1}{5184}\right)$$

$$\approx 5.916087963$$

$$\sqrt{35} = 5.916079783$$

$$\text{Percentage error} = \frac{5.916087963 - 5.916079783}{5.916079783} \times 100 = 0.000138\%$$

**4 a**  $\frac{1}{\sqrt{a+bx}} = (a+bx)^{-\frac{1}{2}} = \left(a\left(1+\frac{b}{a}x\right)\right)^{-\frac{1}{2}} = a^{-\frac{1}{2}}\left(1+\frac{b}{a}x\right)^{-\frac{1}{2}} = \frac{1}{a^{\frac{1}{2}}}\left(1+\frac{b}{a}x\right)^{-\frac{1}{2}}$

$$= \frac{1}{a^{\frac{1}{2}}} \left( 1 + \left( -\frac{1}{2} \right) \left( \frac{b}{a}x \right) + \frac{\left( -\frac{1}{2} \right) \left( -\frac{1}{2}-1 \right)}{2!} \left( \frac{b}{a}x \right)^2 + \dots \right)$$

$$= \frac{1}{a^{\frac{1}{2}}} \left( 1 + \left( -\frac{1}{2} \right) \left( \frac{b}{a}x \right) + \frac{\left( -\frac{1}{2} \right) \left( -\frac{3}{2} \right)}{2} \frac{b^2}{a^2} x^2 + \dots \right)$$

$$= \frac{1}{a^{\frac{1}{2}}} \left( 1 - \frac{b}{2a}x + \frac{3b^2}{8a^2}x^2 + \dots \right)$$

$$= \frac{1}{a^{\frac{1}{2}}} - \frac{b}{2a^{\frac{3}{2}}}x + \frac{3b^2}{8a^{\frac{5}{2}}}x^2 + \dots = 3 + \frac{1}{3}x + \frac{1}{18}x^2 + \dots$$

Equating coefficients gives  $\frac{1}{a^{\frac{1}{2}}} = 3$ , so  $a = \frac{1}{9}$

and  $-\frac{b}{2\left(\frac{1}{9}\right)^{\frac{3}{2}}} = \frac{1}{3}$

$$-\frac{b}{\frac{2}{27}} = \frac{1}{3}$$

$$b = -\frac{2}{81}$$

$$a = \frac{1}{9}, b = -\frac{2}{81}$$

**4 b**  $x^3$  term of  $3\left(1 - \frac{2}{9}x\right)^{-\frac{1}{2}} = 3 \left( \frac{\left(-\frac{1}{2}\right)\left(-\frac{1}{2}-1\right)\left(-\frac{1}{2}-2\right)}{3!} \left(-\frac{2}{9}x\right)^3 \right)$

$$= -3 \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)\left(-\frac{5}{2}\right)}{6} \frac{8}{729} x^3$$

$$= \frac{5}{486} x^3$$

Coefficient =  $\frac{5}{486}$

**5**  $\frac{3+2x-x^2}{4-x} \equiv (3+2x-x^2)(4-x)^{-1}$  Write  $\frac{1}{4-x}$  as  $(4-x)^{-1}$

$$= (3+2x-x^2) \left( 4 \left(1 - \frac{x}{4}\right) \right)^{-1} \quad \text{Take out a factor of 4}$$

$$= (3+2x-x^2) \frac{1}{4} \left(1 - \frac{x}{4}\right)^{-1} \quad \text{Expand } \left(1 - \frac{x}{4}\right)^{-1} \text{ using the binomial expansion}$$

$$= (3+2x-x^2) \frac{1}{4} \left( 1 + (-1) \left( -\frac{x}{4} \right) + \frac{(-1)(-2)}{2!} \left( -\frac{x}{4} \right)^2 + \dots \right) \quad \text{Ignore terms higher than } x^2$$

$$= (3+2x-x^2) \frac{1}{4} \left( 1 + \frac{x}{4} + \frac{x^2}{16} + \dots \right) \quad \text{Multiply expansion by } \frac{1}{4}$$

$$= (3+2x-x^2) \left( \frac{1}{4} + \frac{x}{16} + \frac{x^2}{64} + \dots \right) \quad \text{Multiply result by } (3+2x-x^2)$$

$$= 3 \left( \frac{1}{4} + \frac{x}{16} + \frac{x^2}{64} + \dots \right) + 2x \left( \frac{1}{4} + \frac{x}{16} + \frac{x^2}{64} + \dots \right) - x^2 \left( \frac{1}{4} + \frac{x}{16} + \frac{x^2}{64} + \dots \right)$$

$$= \frac{3}{4} + \frac{3}{16}x + \frac{3}{64}x^2 + \frac{1}{2}x + \frac{1}{8}x^2 - \frac{1}{4}x^2 + \dots \quad \text{Ignore any terms bigger than } x^2$$

$$= \frac{3}{4} + \frac{11}{16}x - \frac{5}{64}x^2$$

Expansion is valid if  $\left| \frac{-x}{4} \right| < 1 \Rightarrow |x| < 4$

## Pure Mathematics 4 Solution Bank

**6 a**

$$\begin{aligned} \frac{1}{\sqrt{5+2x}} &= (5+2x)^{-\frac{1}{2}} = 5^{-\frac{1}{2}} \left(1 + \frac{2}{5}x\right)^{-\frac{1}{2}} = \frac{1}{\sqrt{5}} \left(1 + \frac{2}{5}x\right)^{-\frac{1}{2}} \\ &= \frac{1}{\sqrt{5}} \left(1 + \left(-\frac{1}{2}\right) \left(\frac{2}{5}x\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{1}{2}-1\right)}{2!} \left(\frac{2}{5}x\right)^2 + \dots\right) \\ &= \frac{1}{\sqrt{5}} \left(1 + \left(-\frac{1}{2}\right) \left(\frac{2}{5}x\right) + \frac{\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{2} \frac{4}{25}x^2 + \dots\right) \\ &= \frac{1}{\sqrt{5}} \left(1 - \frac{1}{5}x + \frac{3}{50}x^2 + \dots\right) \\ &= \frac{1}{\sqrt{5}} - \frac{1}{5\sqrt{5}}x + \frac{3}{50\sqrt{5}}x^2 + \dots \end{aligned}$$

**b**

$$\begin{aligned} \frac{2x-1}{\sqrt{5+2x}} &= \frac{2x-1}{\sqrt{5}} \left(1 + \frac{2}{5}x\right)^{-\frac{1}{2}} \\ &= (2x-1) \left(\frac{1}{\sqrt{5}} - \frac{1}{5\sqrt{5}}x + \frac{3}{50\sqrt{5}}x^2 + \dots\right) \\ &= \frac{2}{\sqrt{5}}x - \frac{2}{5\sqrt{5}}x^2 - \frac{1}{\sqrt{5}} + \frac{1}{5\sqrt{5}}x - \frac{3}{50\sqrt{5}}x^2 + \dots \\ &= -\frac{1}{\sqrt{5}} + \frac{11}{5\sqrt{5}}x - \frac{23}{50\sqrt{5}}x^2 + \dots \end{aligned}$$

**7 a**

$$\begin{aligned} (16-3x)^{\frac{1}{4}} &= \left(16 \left(1 - \frac{3}{16}x\right)\right)^{\frac{1}{4}} = 2 \left(1 - \frac{3}{16}x\right)^{\frac{1}{4}} \\ &= 2 \left(1 + \left(\frac{1}{4}\right) \left(-\frac{3}{16}x\right) + \frac{\left(\frac{1}{4}\right)\left(\frac{1}{4}-1\right)}{2!} \left(-\frac{3}{16}x\right)^2 + \dots\right) \\ &= 2 \left(1 + \left(\frac{1}{4}\right) \left(-\frac{3}{16}x\right) + \frac{\left(\frac{1}{4}\right)\left(-\frac{3}{4}\right)}{2} \frac{9}{256}x^2 + \dots\right) \\ &= 2 \left(1 - \frac{3}{64}x - \frac{27}{8192}x^2 + \dots\right) \\ &= 2 - \frac{3}{32}x - \frac{27}{4096}x^2 + \dots \end{aligned}$$

**7 b** Let  $x = 0.1$

$$\begin{aligned}\sqrt[4]{15.7} &\approx 2 - \frac{3}{32}(0.1) - \frac{27}{4096}(0.1)^2 \\ &= 1.991\end{aligned}$$

$$\begin{aligned}\mathbf{8 a} \quad \frac{3}{4-2x} &= 3(4-2x)^{-1} = 3(4(1-\frac{1}{2}x))^{-1} = \frac{3}{4}(1-\frac{1}{2}x)^{-1} \\ &= \frac{3}{4} \left( 1 + (-1)(-\frac{1}{2}x) + \frac{(-1)(-2)}{2!}(-\frac{1}{2}x)^2 + \dots \right) \\ &= \frac{3}{4} \left( 1 - \left( -\frac{1}{2}x \right) + \frac{1}{4}x^2 + \dots \right) \\ &= \frac{3}{4} + \frac{3}{8}x + \frac{3}{16}x^2 + \dots\end{aligned}$$

$$\begin{aligned}\frac{2}{3+5x} &= 2(3+5x)^{-1} = 2(3(1+\frac{5}{3}x))^{-1} = \frac{2}{3}(1+\frac{5}{3}x)^{-1} \\ &= \frac{2}{3} \left( 1 + (-1)(\frac{5}{3}x) + \frac{(-1)(-2)}{2!}(\frac{5}{3}x)^2 + \dots \right) \\ &= \frac{2}{3} \left( 1 - \frac{5}{3}x + \frac{25}{9}x^2 + \dots \right) \\ &= \frac{2}{3} - \frac{10}{9}x + \frac{50}{27}x^2 + \dots\end{aligned}$$

$$\begin{aligned}\frac{3}{4-2x} - \frac{2}{3+5x} &= \frac{3}{4} + \frac{3}{8}x + \frac{3}{16}x^2 + \dots - \left( \frac{2}{3} - \frac{10}{9}x + \frac{50}{27}x^2 + \dots \right) \\ &= \frac{1}{12} + \frac{107}{72}x - \frac{719}{432}x^2 + \dots\end{aligned}$$

$$\mathbf{b} \quad g(0.01) = \frac{3}{4-2(0.01)} - \frac{2}{3+5(0.01)} = 0.0980311$$

**c** Using the series expansion:

$$g(0.01) \approx \frac{1}{12} + \frac{107}{72}(0.01) - \frac{719}{432}(0.01)^2 = 0.098028009$$

$$\text{Percentage error} = \frac{0.0980311 - 0.098028009}{0.098028009} \times 100 = 0.0032\%$$