

1. (a) Show that $\sum_{r=1}^n (r+1)(r+5) = \frac{1}{6}n(n+7)(2n+7)$.

(4)

(b) Hence calculate the value of $\sum_{r=10}^{40} (r+1)(r+5)$

(2)

(Total 6 marks)

1. (a) Expand brackets and attempt to use appropriate formulae. M1
- $$\Sigma r^2 + 6r + 5 = \frac{n}{6}(n+1)(2n+1) + 6\frac{n}{2}(n+1) + 5n$$
- A1
- $$= \frac{n}{6}[2n^2 + 3n + 1 + 18n + 18 + 30]$$
- M1
- $$= \frac{n}{6}[2n^2 + 21n + 49] = \frac{n}{6}(n+7)(2n+7) (*)$$
- A1 4
- (b) Use $S(40) - S(9) = \frac{40}{9} \times 47 \times 87 - \frac{9}{6} \times 16 \times 25$ M1
- $$= 26660$$
- A1 2

[6]

1. (a) On the whole, candidates were able to expand $(r+1)(r+5)$ accurately and were able to substitute correctly for Σr^2 , Σr and to deal with $\Sigma 5$ correctly – the provision of the answer helped many to check the accuracy of their subsequent expansions, collection of terms and factorisation! A small group of candidates attempted Mathematical Induction, but rarely correctly, most being daunted by the algebra involved.
- (b) For those not working out $S(40) - S(9)$, the most common mistake was to use $S(40) - S(10)$, although some returned to using $(r+1)(r+5)$ with $r = 40$, and 9 or 10; some just calculated $S(40)$, totally ignoring the starting value of r .