

Mark Scheme (Results)

January 2013

GCE Further Pure Mathematics FP1 (6667/01)

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.

Their contact details can be found on this link: www.edexcel.com/teachingservices.

You can also use our online Ask the Expert service at www.edexcel.com/ask. You will need an Edexcel username and password to access this service.

Pearson: helping people progress, everywhere

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

January 2013

Publications Code UA034509

All the material in this publication is copyright

© Pearson Education Ltd 2013

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Unless indicated in the mark scheme a correct answer with no working should gain full marks for that part of the question.

EDEXCEL GCE MATHEMATICS

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
 - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

In some instances, the mark distributions (e.g. M1, B1 and A1) printed on the candidate's response may differ from the final mark scheme.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \checkmark will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper
 - \square The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but incorrect answers should never be awarded A marks.
 5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.

6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.
8. The maximum mark allocation for each question/part question(item) is set out in the marking grid and you should allocate a score of '0' or '1' for each mark, or "trait", as shown:

	0	1
aM		•
aA	•	
bM1		•
bA1	•	
bB	•	
bM2		•
bA2		•


Jan 2013
Further Pure Mathematics FP1 6667
Mark Scheme

Question Number	Scheme	Marks
1.	$\sum_{r=1}^n 3(4r^2 - 4r + 1) = 12 \sum_{r=1}^n r^2 - 12 \sum_{r=1}^n r + \sum_{r=1}^n 3$ $= \frac{12}{6} n(n+1)(2n+1) - \frac{12}{2} n(n+1), \quad +3n$ $= n[2(n+1)(2n+1) - 6(n+1) + 3]$ $= n[4n^2 - 1] = n(2n+1)(2n-1)$	<p>M1</p> <p>A1, B1</p> <p>M1</p> <p>A1 cso</p> <p style="text-align: right;">[5]</p>
Notes:	<p>Induction is not acceptable here</p> <p>First M for expanding given expression to give a 3 term quadratic and attempt to substitute.</p> <p>First A for first two terms correct or equivalent.</p> <p>B for +3n appearing</p> <p>Second M for factorising by n</p> <p>Final A for completely correct solution</p>	

Question Number	Scheme	Marks
2.	<p>(a) $\frac{50}{3+4i} = \frac{50(3-4i)}{(3+4i)(3-4i)} = \frac{50(3-4i)}{25} = 6-8i$</p> <p>(b) $z^2 = (6-8i)^2 = 36 - 64 - 96i = -28 - 96i$</p> <p>(c) $z = \sqrt{6^2 + (-8)^2} = 10$</p> <p>(d) $\tan \alpha = \frac{-96}{-28}$</p> <p>so $\alpha = -106.3^\circ$ or 253.7°</p>	<p>M1 A1cao (2)</p> <p>M1 A1 (2)</p> <p>M1 A1ft (2)</p> <p>M1 A1cao (2) [8]</p>
	<p>Alternatives</p> <p>(c) $z = \frac{50}{ 3+4i } = 10$</p> <p>(d) $\arg(3+4i) = 53.13..$ so $\arg\left(\frac{50}{3+4i}\right)^2 = -2 \times 53.13... = -106.3$</p>	<p>M1 A1</p> <p>M1 A1</p>
Notes:	<p>(a) M for $\times \frac{3-4i}{3-4i}$ (accept use of $-3+4i$) and attempt to expand using $i^2=-1$, A for $6-8i$ only</p> <p>(b) M for attempting to expand their z^2 using $i^2=-1$, A for $-28-96i$ only. If using original z then must attempt to multiply top and bottom by conjugate and use $i^2=-1$.</p> <p>(c) M for $\sqrt{a^2+b^2}$, A for 'their 10'</p> <p>(d) M for use of \tan or \tan^{-1} and values from their z^2 either way up ignoring signs. Radians score A0.</p>	

Question Number	Scheme	Marks
3.	<p>(a) $f'(x) = x^{-\frac{1}{2}} - \frac{1}{2}x^{-\frac{3}{2}}$</p> <p>(b) $f(5) = -0.0807$ $f'(5) = 0.4025$</p> $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} = 5 - \frac{-0.0807}{0.4025}$ $= 5.2(0)$	<p>M1 A1 (2)</p> <p>B1 M1 M1 A1 (4) [6]</p>
Notes	<p>The B and M marks are implied by a correct answer only with no working or by $\frac{5}{9}(10\sqrt{5} - 13)$</p> <p>(a) M for at least one of $\pm ax^{-\frac{1}{2}}$ or $\pm bx^{-\frac{3}{2}}$, A for correct (equivalent) answer only</p> <p>(b) B for awrt -0.0807, first M for attempting their $f'(5)$, M for correct formula and attempt to substitute, A for awrt 5.20, but accept 5.2</p>	

Question Number	Scheme	Marks
4.	<p>(a) $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$</p> <p>(b) $\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$</p> <p>(c) R = QP</p> <p>(d) $R = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$</p> <p>(e) Reflection in the y axis</p>	<p>B1 (1)</p> <p>B1 (1)</p> <p>B1 (1)</p> <p>M1 A1 cao (2)</p> <p>B1 B1 (2) [7]</p>
Notes	<p>(a) and (b) Signs must be clear for B marks.</p> <p>(c) Accept QP or their 2x2 matrices in the correct order only for B1.</p> <p>(d) M for their QP where answer involves ± 1 and 0 in a 2x2 matrix, A for correct answer only.</p> <p>(e) First B for Reflection, Second B for 'y axis' or 'x=0'. Must be single transformation. Ignore any superfluous information.</p>	

Question Number	Scheme	Marks
5.	<p>(a) $4x^2 + 9 = 0 \Rightarrow x = ki, \quad x = \pm \frac{3}{2}i$ or equivalent</p> <p>Solving 3-term quadratic by formula or completion of the square</p> $x = \frac{6 \pm \sqrt{36 - 136}}{2} \text{ or } (x - 3)^2 - 9 + 34 = 0$ $= 3 + 5i \text{ and } 3 - 5i$ <p>(b)</p> 	<p>M1, A1</p> <p>M1</p> <p>A1 A1ft</p> <p>(5)</p> <p>B1ft</p> <p>B1ft</p> <p>(2)</p> <p>[7]</p>
Notes	<p>(a) Final A follow through conjugate of their first root.</p> <p>(b) First B award only for first pair imaginary, Second B award only if second pair complex. Complex numbers labelled, scales or coordinates or vectors required for B marks.</p>	

Question Number	Scheme	Marks
6.	<p>(a) Determinant: $2 - 3a = 0$ and solve for $a =$ So $a = \frac{2}{3}$ or equivalent</p> <p>(b) Determinant: $(1 \times 2) - (3 \times -1) = 5$ (Δ) $Y^{-1} = \frac{1}{5} \begin{pmatrix} 2 & 1 \\ -3 & 1 \end{pmatrix} = \begin{bmatrix} 0.4 & 0.2 \\ -0.6 & 0.2 \end{bmatrix}$</p> <p>(c) $\frac{1}{5} \begin{pmatrix} 2 & 1 \\ -3 & 1 \end{pmatrix} \begin{pmatrix} 1-\lambda \\ 7\lambda-2 \end{pmatrix} = \frac{1}{5} \begin{pmatrix} 2-2\lambda+7\lambda-2 \\ -3+3\lambda+7\lambda-2 \end{pmatrix} = \begin{pmatrix} \lambda \\ 2\lambda-1 \end{pmatrix}$</p>	<p>M1 A1 (2)</p> <p>M1A1 (2)</p> <p>M1depM1A1 A1 (4) [8]</p>
	<p>Alternative method for (c) $\begin{pmatrix} 1 & -1 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1-\lambda \\ 7\lambda-2 \end{pmatrix}$ so $x - y = 1 - \lambda$ and $3x + 2y = 7\lambda - 2$</p> <p>Solve to give $x = \lambda$ and $y = 2\lambda - 1$</p>	<p>M1M1 A1A1</p>
Notes	<p>(b) M for $\frac{1}{\text{their det}} \begin{pmatrix} 2 & 1 \\ -3 & 1 \end{pmatrix}$</p> <p>(c) First M for their $Y^{-1}B$ in correct order with B written as a 2×1 matrix, second M dependent on first for attempt at multiplying their matrices resulting in a 2×1 matrix, first A for λ, second A for $2\lambda - 1$</p> <p>Alternative for (c) First M to obtain two linear equations in x, y, λ Second M for attempting to solve for x or y in terms of λ</p>	

Question Number	Scheme	Marks
7.	<p>(a) $y = \frac{25}{x}$ so $\frac{dy}{dx} = -25x^{-2}$</p> $\frac{dy}{dx} = -\frac{25}{(5p)^2} = -\frac{1}{p^2}$ $y - \frac{5}{p} = -\frac{1}{p^2}(x - 5p) \Rightarrow p^2y + x = 10p \quad (*)$ <p>(b) $q^2y + x = 10q$ only</p> <p>(c) $(p^2 - q^2)y = 10(p - q)$ so $y = \frac{10(p - q)}{(p^2 - q^2)} = \frac{10}{p + q}$</p> $x = 10p - p^2 \frac{10}{p + q} = \frac{10pq}{p + q}$ <p>(d) Line PQ has gradient $\frac{\frac{5}{p} - \frac{5}{q}}{5p - 5q} \left(= -\frac{1}{pq} \right)$</p> <p>$ON$ has gradient $\frac{\frac{10}{p+q}}{\frac{10pq}{p+q}} \left(= \frac{1}{pq} \right)$ or $\frac{-1}{-1} (= pq)$ could be as unsimplified</p> <p>equivalents seen anywhere</p> <p>As these lines are perpendicular $\frac{1}{pq} \times -\frac{1}{pq} = -1$ so $p^2q^2 = 1$</p> <p>OR for ON $y - y_1 = m(x - x_1)$ with gradient (equivalent to) pq and sub in points O AND N to give $p^2q^2 = 1$</p> <p>OR for PQ $y - y_1 = m(x - x_1)$ with gradient (equivalent to) $-pq$ and sub in points P AND Q to give $p^2q^2 = 1$. NB $-pq$ used as gradient of PQ implies first M1A1</p>	<p>M1</p> <p>A1</p> <p>M1 A1</p> <p>(4)</p> <p>B1</p> <p>(1)</p> <p>M1 A1cso</p> <p>M1 A1 cso</p> <p>(4)</p> <p>M1 A1</p> <p>B1</p> <p>M1 A1</p> <p>(5)</p> <p>[14]</p>

Question Number	Scheme	Marks
	<p><u>Alternatives for first M1 A1 in part (a)</u></p> $x \frac{dy}{dx} + y = 0 \Rightarrow \frac{dy}{dx} = -\frac{y}{x}$ <p>So at P gradient = $\frac{-\frac{5}{p}}{5p} = -\frac{1}{p^2}$</p> <p>Or $x = 5t, y = \frac{5}{t} \Rightarrow \frac{dx}{dt} = 5, \frac{dy}{dt} = -\frac{5}{t^2}$ so $\frac{dy}{dx} =$</p> $\frac{-\frac{5}{t^2}}{5} = -\frac{1}{t^2} \text{ so at } P \text{ gradient} = -\frac{1}{p^2}$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>
Notes	<p>(a) First M for attempt at explicit, implicit or parametric differentiation not using p or q as an initial parameter, first A for $-\frac{1}{p^2}$ or equivalent. Quoting gradient award first M0A0. Second M for using $y - y_1 = m(x - x_1)$ and attempt to substitute or $y = mx + c$ and attempt to find c; gradient in terms of p only and using $\left(5p, \frac{5}{p}\right)$, second A for correct solution only.</p> <p>(c) First M for eliminating x and reaching $y = f(p, q)$, second M for eliminating y and reaching $x = f(p, q)$, both As for given answers. Minimum amount of working given in the main scheme above for 4/4, but do not award accuracy if any errors are made.</p> <p>(d) First M for use of $\frac{y_2 - y_1}{x_2 - x_1}$ and substituting, first A for $-\frac{1}{pq}$ or unsimplified equivalent.</p> <p>Second M for their product of gradients = -1 (or equating equivalent gradients of ON or equating equivalent gradients of PQ), second A for correct answer only.</p>	

Question Number	Scheme	Marks
8.	<p>(a) If $n = 1$, $\sum_{r=1}^n r(r+3) = 1 \times 4 = 4$ and $\frac{1}{3}n(n+1)(n+5) = \frac{1}{3} \times 1 \times 2 \times 6 = 4$,</p> <p>(so true for $n = 1$. Assume true for $n = k$)</p> $\text{So } \sum_{r=1}^{k+1} r(r+3) = \frac{1}{3}k(k+1)(k+5) + (k+1)(k+4)$ $= \frac{1}{3}(k+1)[k(k+5) + 3(k+4)] = \frac{1}{3}(k+1)[k^2 + 8k + 12]$ $= \frac{1}{3}(k+1)(k+2)(k+6) \text{ which implies is true for } n = k + 1$ <p>As result is true for $n = 1$ this implies true for all positive integers and so result is true by induction</p> <p>(b) $u_1 = 1^2(1-1) + 1 = 1$</p> <p>(so true for $n = 1$. Assume true for $n = k$)</p> $u_{k+1} = k^2(k-1) + 1 + k(3k+1)$ $= k(k^2 - k + 3k + 1) + 1 = k(k+1)^2 + 1 \text{ which implies is true for } n = k + 1$ <p>As result is true for $n = 1$ this implies true for all positive integers and so result is true by induction</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>dA1</p> <p>dM1A1cso</p> <p>(6)</p> <p>B1</p> <p>M1, A1</p> <p>M1A1cso</p> <p>(5)</p> <p>[11]</p>
Notes	<p>(a) First B for LHS=4 and RHS =4</p> <p>First M for attempt to use $\sum_1^k r(r+3) + u_{k+1}$</p> <p>First A for $\frac{1}{3}(k+1)$, $\frac{1}{3}(k+2)$ or $\frac{1}{3}(k+6)$ as a factor before the final line</p> <p>Second A dependent on first for $\frac{1}{3}(k+1)(k+2)(k+6)$ with no errors seen</p> <p>Second M dependent on first M and for any 3 of 'true for $n=1$' 'assume true for $n=k$' 'true for $n=k+1$', 'true for all n' (or 'true for all positive integers') seen anywhere</p> <p>Third A for correct solution only with all statements and no errors</p>	

	<p>(b) First B for both some working and 1.</p> <p>First M for $u_{k+1} = u_k + k(3k + 1)$ and attempt to substitute for u_k</p> <p>First A for $k(k + 1)^2 + 1$ with some correct intermediate working and no errors seen</p> <p>Second M dependent on first M and for any 3 of 'true for $n=1$' 'assume true for $n=k$' 'true for $n=k+1$', 'true for all n' (or 'true for all positive integers') seen anywhere</p> <p>Second A for correct solution only with all statements and no errors</p>	
--	---	--

Question Number	Scheme	Marks
<p>9.</p>	<p>(a) $y = 6x^{\frac{1}{2}}$ so $\frac{dy}{dx} = 3x^{-\frac{1}{2}}$</p> <p>Gradient when $x = 4$ is $\frac{3}{2}$ and gradient of normal is $-\frac{2}{3}$</p> <p>So equation of normal is $(y-12) = -\frac{2}{3}(x-4)$ (or $3y + 2x = 44$)</p> <p>(b) S is at point $(9,0)$ N is at $(22,0)$, found by substituting $y=0$ into their part (a) Both B marks can be implied or on diagram. So area is $\frac{1}{2} \times 12 \times (22-9) = 78$</p> <p>Alternatives: First M1 for $ky \frac{dy}{dx} = 36$ or for $x = 9t^2, y = 18t \rightarrow \frac{dx}{dt} = 18t, \frac{dy}{dt} = 18 \rightarrow \frac{dy}{dx} = \frac{1}{t}$</p>	<p>M1</p> <p>M1 A1</p> <p>M1 A1</p> <p>(5)</p> <p>B1 B1ft</p> <p>M1 A1 cao</p> <p>(4) [9]</p>
<p>Notes</p>	<p>(a) First M for $\frac{dy}{dx} = ax^{-\frac{1}{2}}$,</p> <p>Second M for substituting $x=4$ (or $y=12$ or $t=2/3$ if alternative used) into their gradient and applying negative reciprocal.</p> <p>First A for $-\frac{2}{3}$</p> <p>Third M for $y - y_1 = m(x - x_1)$ or $y = mx + c$ and attempt to substitute a changed gradient AND $(4,12)$</p> <p>Second A for $3y + 2x = 44$ or any equivalent equation</p> <p>(b) M for Area = $\frac{1}{2}$ base x height and attempt to substitute including their numerical '(22-9)' or equivalent complete method to find area of triangle PSN.</p>	

Further copies of this publication are available from
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467
Fax 01623 450481
Email publication.orders@edexcel.com
Order Code UA034509 January 2013

For more information on Edexcel qualifications, please visit our website
www.edexcel.com

Pearson Education Limited. Registered company number 872828
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual




Llywodraeth Cynulliad Cymru
Welsh Assembly Government

