

1. (a) $y' = 3\sin 2x + 6x \cos 2x$ M1

$$y'' = 12 \cos 2x - 12x \sin 2x$$
 A1

Substituting $12 \cos 2x - 12x \sin 2x + 12x \sin 2x = k \cos 2x$ M1

$$k = 12$$
 A1 4

(b) General solution is $y = A \cos 2x + B \sin 2x + 3x \sin 2x$ B1

$$(0, 2) \Rightarrow A = 2$$
 B1

$$\left(\frac{\pi}{4}, \frac{\pi}{2}\right) \Rightarrow \frac{\pi}{2} = B + \frac{3\pi}{4} \Rightarrow B = -\frac{\pi}{4}$$
 M1

$$y = 2 \cos 2x - \frac{\pi}{4} \sin 2x + 3x \sin 2x$$
 Needs $y = \dots$ A1 4

[8]

2. (a) $(2r+1)^3 = 8r^3 + 12r^2 + 6r + 1$

$$(2r-1)^3 = 8r^3 - 12r^2 + 6r - 1$$

$$(2r+1)^3 - (2r-1)^3 = 24r^2 + 2 \quad (A = 24, B = 2)$$
 M1 A1 2

Accept $r = 0 \Rightarrow B = 2$ and $r = 1 \Rightarrow A + B = 26 \Rightarrow A = 24$

M1 for both

(b) $\cancel{3^3} - 1^3 = 24 \times 1^2 + 2$

$$\cancel{5^3} - \cancel{3^3} = 24 \times 2^2 + 2$$
 M

$$(2n+1)^3 - \cancel{(2n-1)^3} = 24 \times n^2 + 2$$

$$(2n+1)^3 - 1^3 = 24 \sum_{r=1}^n r^2 + \underline{2n}$$
 ft their B M1 A1 A1ft

$$\sum_{r=1}^n r^2 = \frac{8n^3 + 12n^2 + 4n}{24}$$
 M1

$$= \frac{1}{6}n(2n^2 + 3n + 1) = \frac{1}{6}n(n+1)(2n+1)$$
 cso A1 5

$$(c) \quad \sum_{r=1}^{40} (3r-1)^2 = \sum_{r=1}^{40} (9r^2 - 6r + 1) \quad \text{M1}$$

$$= 9 \times \frac{1}{6} \times 40 \times 41 \times 81 - 6 \times \frac{1}{2} \times 40 \times 41 + 40 \quad \text{M1}$$


$$= 194380 \quad \text{A1} \quad 3 \\ \boxed{[10]}$$

3. (a) $2x^2 + x - 6 = 6 - 3x$ M1

Leading to $x^2 + 2x - 6 = 0$
 $(x+1)^2 = 7 \Rightarrow x = -1 \pm \sqrt{7}$ surds required
 $-2x^2 - x + 6 = 6 - 3x$ M1 A1
 M1

Leading to $2x^2 - 2x = 0, \Rightarrow x = 0, 1$ A1 A1 6

(b) Accept if parts (a) and (b) done in reverse order



Curved shape B1
 Line B1
 At least 3 intersections B1 3

(c) Using all 4 CVs and getting all into inequalities M1

$$x > \sqrt{7} - 1, x < -\sqrt{7} - 1 \quad \text{both} \quad \text{A1ft}$$

ft their greatest positive and their least negative CVs

$$0 < x < 1 \quad \text{A1} \quad 3 \\ \boxed{[12]}$$

4. (a) $\int \frac{2}{120-t} dt = -2 \ln(120-t)$ B1
- $$e^{-2\ln(120-t)} = (120-t)^{-2}$$
- M1 A1
- $$\frac{1}{(120-t)^2} \frac{ds}{dt} + \frac{2S}{(120-t)^3} = \frac{1}{4(120-t)^2}$$
- $$\frac{d}{dt} \left(\frac{S}{(120-t)^2} \right) = \frac{1}{4(120-t)^2} \text{ or integral equivalent}$$
- M1
- $$\frac{S}{(120-t)^2} = \frac{1}{4(120-t)} (+C)$$
- M1 A1
- $$(0, 6) \Rightarrow 6 = 30 + 120^2 C \Rightarrow C = -\frac{1}{600}$$
- M1
- $$S = \frac{120-t}{4} - \frac{(120-t)^2}{600} \quad \text{accept } C = \text{awrt } -0.0017$$
- A1 8
- (b) $\frac{dS}{dt} = -\frac{1}{4} + \frac{2(120-t)}{600}$ M1
- $$\frac{dS}{dt} = 0 \Rightarrow t = 45$$
- M1 A1
- substituting $S = 9\frac{3}{8}$ (kg)
- A1 4
[12]

Alternative forms for S are

$$S = 6 + \frac{3t}{20} - \frac{t^2}{600} = \frac{(t+30)(120-t)}{600}$$

$$= \frac{3600 + 90t - t^2}{600} = \frac{5625 - (t-45)^2}{600}$$

Alternative for part (b)

S can be found without finding t

$$\text{Using } \frac{dS}{dt} = 0 \text{ in the original differential equation } \frac{2S}{120-t} = \frac{1}{4} \quad \text{M1}$$

Substituting for t into the answer to part (a)

$$S = 2S - \frac{64S^2}{600} \quad \text{M1 A1}$$

$$\text{Solving to } S = 9\frac{3}{8} \text{ (kg)} \quad \text{A1} \quad 4$$

5. (a) $f(x) = \cos 2x, \quad f\left(\frac{\pi}{4}\right) = 0$

$$f'(x) = -2 \sin 2x, \quad f'\left(\frac{\pi}{4}\right) = -2 \quad \text{M1}$$

$$f''(x) = -4 \cos 2x, \quad f''\left(\frac{\pi}{4}\right) = 0$$

$$f'''(x) = 8 \sin 2x, \quad f'''\left(\frac{\pi}{4}\right) = 8 \quad \text{A1}$$

$$f^{(iv)}(x) = 16 \cos 2x, \quad f^{(iv)}\left(\frac{\pi}{4}\right) = 0$$

$$f^{(v)}(x) = 32 \sin 2x, \quad f^{(v)}\left(\frac{\pi}{4}\right) = -32 \quad \text{A1}$$

$$\cos 2x = f\left(\frac{\pi}{4}\right) + f'\left(\frac{\pi}{4}\right)(x - \frac{\pi}{4}) + \frac{f''\left(\frac{\pi}{4}\right)}{2}(x - \frac{\pi}{4})^2 + \frac{f'\left(\frac{\pi}{4}\right)}{3!}(x - \frac{\pi}{4})^3 + \dots \quad \text{M1}$$

Three terms are sufficient to establish method

$$\cos 2x = -2(x - \frac{\pi}{4}) + \frac{4}{3}(x - \frac{\pi}{4})^3 - \frac{4}{15}(x - \frac{\pi}{4})^5 + \dots \quad \text{A1} \quad 5$$

(b) Substitute $x = 1$ $(1 - \frac{\pi}{4} \approx 0.21460)$ B1

$$\cos 2 = -2(x - \frac{\pi}{4}) + \frac{4}{3}(x - \frac{\pi}{4})^3 - \frac{4}{15}(x - \frac{\pi}{4})^5 + \dots$$

$$\approx -0.416147 \quad \text{cao} \quad \text{M1 A1} \quad 3$$

[8]

6. (a) In this solution $\cos \theta = c$ and $\sin \theta = s$

$$\cos 5\theta + i \sin 5\theta = (c + is)^5$$

M1

$$(= c^5 + 5c^4 is + 10c^3 (is)^2 + 10c^2 (is)^3 + 5c (is)^4 + (is)^5)$$

$$\sin 5\theta = 5c^4 s - 10c^2 s^3 + s^5$$

M1 A1

$$= 5c^4 s - 10c^2(1 - c^2)s + (1 - c^2)^2 s \quad s^2 = 1 - c^2$$

M1

$$= s(16c^4 - 12c^2 + 1)$$

A1 5

(b) $\sin \theta(16\cos^4 \theta - 12\cos^2 \theta + 1) + 2\cos^2 \theta \sin \theta = 0$

M1

$$\sin \theta = 0 \Rightarrow \theta = 0$$

B1

$$16c^4 - 10c^2 + 1 = (8c^2 - 1)(2c^2 - 1) = 0$$

M1

$$c = \pm \frac{1}{2\sqrt{2}}, \quad c = \pm \frac{1}{\sqrt{2}} \quad \text{any two}$$

A1

$$\theta \approx 1.21, 1.93; \quad \theta = \frac{\pi}{4}, \frac{3\pi}{4} \quad \text{any two}$$

A1

all four

A1 6

*accept awrt 0.79, 1.21, 1.93, 2.36**Ignore any solutions out of range.*

[11]

7. (a) $\left(\frac{dx}{dt} \right)_0 = 0.4 \approx \frac{x_{0.1} - 0}{0.1} \Rightarrow x_{0.1} \approx 0.04$

B1

$$\left(\frac{d^2x}{dt^2} \right)_{0.1} = 3 \sin x_{0.1} \approx \frac{x_{0.2} - 2x_{0.1} + 0}{0.01}$$

M1

Must have their $x_{0.1}$

$$x_{0.2} \approx 0.0788 \quad \text{awrt}$$

A1

$$\left(\frac{d^2x}{dt^2} \right)_{0.2} = 3 \sin x_{0.2} \approx \frac{x_{0.3} - 2x_{0.2} + x_{0.1}}{0.01}$$

M1

Must have their $x_{0.1}, x_{0.2}$

$$x_{0.3} \approx 0.115 \quad \text{awrt}$$

A1 5

(b) $f'(t) = -3\sin x , \quad f''(0) = 0$

$$f'''(t) = -3\cos x \frac{dx}{dt}, \quad f'''(0) = -3 \times 0.4 = -1.2 \quad \text{M1 A1}$$

$$f(t) = f(0) + f'(0)t + \frac{t^2}{2}f''(0) + \frac{t^3}{3!}f'''(0) + \dots$$

$$= 0.4t - 0.2t^3$$

M1 A1 4

- (c) Substituting $t = 0.3$ into their answer to (b) and evaluating
 $f(0.3) \approx 0.1146$ cao

M1
A1 2
[11]

8. (a) Let $z = x + iy$

$$(x - 6)^2 + (y + 3)^2 = 9[(x + 2)^2 + (y - 1)^2]$$

M1

$$\text{Leading to } 8x^2 + 8y^2 + 48x - 24y = 0$$

M1 A1

This is a circle; the coefficients of x^2 and y^2 are the same and there is no xy term.

Allow equivalent arguments and find their $f, (x, y)$ if appropriate.

A1ft

$$(x^2 + 6x + y^2 - 3y = 0)$$

$$\text{Leading to } (x + 3)^2 + (y - \frac{3}{2})^2 = \frac{45}{4}$$

M1

$$\text{Centre: } (-3, \frac{3}{2})$$

A1

$$\text{Radius: } \frac{3}{2}\sqrt{5} \quad \text{or equivalent}$$

A1 7

Alternative

Accept the following argument:-

The locus of P is a Circle of Apollonius, which is a circle with diameter XY, where the points X and Y cut (6, -3) and (-2, 1) internally and externally in the ratio 3 : 1.

M1 A1

$$X: (0, 0) Y: (-6, 3)$$


M1 A1

$$\text{Centre: } (-3, \frac{3}{2})$$

M1 A1

$$\text{Radius: } \frac{3}{2}\sqrt{5} \text{ or equivalent}$$

A1 7



Circle	B1
centre in correct quadrant	B1 ft
through origin	B1
Line cuts -ve x and +ve y axes	B1
intersects with circle on axes and all correct	B1 5

(c)	Shading inside circle and above line with all correct	B1 B1 2
<i>Having 3 instead of 9 in first equation gains maximum of M1 M1 A0 A1 ft M1 A0 A0 B1 B1 B0 B1 B0 B1 B0 8/14</i>		

[14]