

4727 Further Pure Mathematics 3

1	METHOD 1 line segment between l_1 and $l_2 = \pm[4, -3, -9]$ $\mathbf{n} = [1, -1, 2] \times [2, 3, 4] = (\pm)[-2, 0, 1]$ distance = $\frac{ [4, -3, -9] \cdot [-2, 0, 1] }{(\sqrt{2^2 + 0^2 + 1^2})} = \frac{17}{(\sqrt{5})}$ $\neq 0$, so skew	B1 M1* A1 M1 (*dep) A1 5	For correct vector For finding vector product of direction vectors For using numerator of distance formula For correct scalar product and correct conclusion
	METHOD 2 lines would intersect where $\begin{cases} 1 + s = -3 + 2t \\ -2 - s = 1 + 3t \\ -4 + 2s = 5 + 4t \end{cases} \Rightarrow \begin{cases} s - 2t = -4 \\ s + 3t = -3 \\ 2s - 4t = 9 \end{cases}$ \Rightarrow contradiction, so skew	B1 M1* A1 M1 (*dep) A1	For correct parametric form for either line For 3 equations using 2 different parameters For attempting to solve to show (in)consistency For correct conclusion
5			
2 (i)	$(a + b\sqrt{5})(c + d\sqrt{5})$ $= ac + 5bd + (bc + ad)\sqrt{5} \in H$	M1 A1 2	For using product of 2 distinct elements For correct expression
(ii)	$(e =) 1 \text{ OR } 1 + 0\sqrt{5}$	B1 1	For correct identity
(iii)	<i>EITHER</i> $\frac{1}{a + b\sqrt{5}} \times \frac{a - b\sqrt{5}}{a - b\sqrt{5}}$ <i>OR</i> $(a + b\sqrt{5})(c + d\sqrt{5}) = 1 \Rightarrow \begin{cases} ac + 5bd = 1 \\ bc + ad = 0 \end{cases}$ inverse = $\frac{a}{a^2 - 5b^2} - \frac{b}{a^2 - 5b^2}\sqrt{5}$	M1 A1 2	For correct inverse as $(a + b\sqrt{5})^{-1}$ and multiplying top and bottom by $a - b\sqrt{5}$ <i>OR</i> for using definition and equating parts For correct inverse. Allow as a single fraction
(iv)	5 is prime <i>OR</i> $\sqrt{5} \notin \mathbb{Q}$	B1 1	For a correct property (or equivalent)
6			
3	Integrating factor = $e^{\int 2dx} = e^{2x}$ $\Rightarrow \frac{d}{dx}(ye^{2x}) = e^{-x}$ $\Rightarrow ye^{2x} = -e^{-x} + c$ $(0, 1) \Rightarrow c = 2$ $\Rightarrow y = -e^{-3x} + 2e^{-2x}$	B1 M1 A1 M1 A1√ A1 6	For correct IF For $\frac{d}{dx}(y \cdot \text{their IF}) = e^{-3x}$. their IF For correct integration both sides For substituting (0, 1) into their GS and solving for c For correct c f.t. from their GS For correct solution
6			
4 (i)	$(z =) 2, -2, 2i, -2i$	M1 A1 2	For at least 2 roots of the form $k\{1, i\}$ AEF For correct values

(ii)	$\frac{w}{1-w} = 2, -2, 2i, -2i$	M1	For $\frac{w}{1-w} =$ any one solution from (i)
	$w = \frac{z}{1+z}$	M1	For attempting to solve for w , using any solution or in general
	$w = \frac{2}{3}, 2$	B1	For any one of the 4 solutions
	$w = \frac{4}{5} \pm \frac{2}{5}i$	A1	For both real solutions
		A1	For both complex solutions
		5	SR Allow B1√ and one A1√ from $k \neq 2$

7

5 (i)	$\mathbf{AB} = k \left[\frac{2}{3}\sqrt{3}, 0, -\frac{2}{3}\sqrt{6} \right],$	B1	For any one edge vector of ΔABC
	$\mathbf{BC} = k \left[-\sqrt{3}, 1, 0 \right], \mathbf{CA} = k \left[\frac{1}{3}\sqrt{3}, -1, \frac{2}{3}\sqrt{6} \right]$	B1	For any other edge vector of ΔABC
	$\mathbf{n} = k_1 \left[\frac{2}{3}\sqrt{6}, \frac{2}{3}\sqrt{18}, \frac{2}{3}\sqrt{3} \right] = k_2 \left[1, \sqrt{3}, \frac{1}{2}\sqrt{2} \right]$	M1	For attempting to find vector product of any two edges
	substitute A, B or $C \Rightarrow x + \sqrt{3}y + \frac{1}{2}\sqrt{2}z = \frac{2}{3}\sqrt{3}$	M1	For substituting A, B or C into $\mathbf{r} \cdot \mathbf{n}$
		A1	5 For correct equation AG
			SR For verification only allow M1, then A1 for 2 points and A1 for the third point

(ii)	Symmetry in plane OAB or Oxz or $y=0$	B1*	For quoting symmetry or reflection
		B1	For correct plane
		(*dep)2	Allow “in y coordinates” or “in y axis”
		SR	For symmetry implied by reference to opposite signs in y coordinates of C and D , award B1 only

(iii)	$\cos \theta = \frac{\left[\left[1, \sqrt{3}, \frac{1}{2}\sqrt{2} \right] \cdot \left[1, -\sqrt{3}, \frac{1}{2}\sqrt{2} \right] \right]}{\sqrt{1+3+\frac{1}{2}}\sqrt{1+3+\frac{1}{2}}}$	M1	For using scalar product of normal vectors
	$= \frac{\left 1-3+\frac{1}{2} \right }{\frac{9}{2}} = \frac{\frac{3}{2}}{\frac{9}{2}} = \frac{1}{3}$	A1	For correct scalar product
		M1	For product of both moduli in denominator
		A1	4 For correct answer. Allow $-\frac{1}{3}$

11

6 (i)	$(m^2 + 16 = 0 \Rightarrow) m = \pm 4i$	M1	For attempt to solve correct auxiliary equation (may be implied by correct CF)
	CF = $A \cos 4x + B \sin 4x$	A1	2 For correct CF
			(A Etrig but not $Ae^{4ix} + Be^{-4ix}$ only)

(ii)	$\frac{dy}{dx} = p \sin 4x + 4px \cos 4x$	M1	For differentiating PI twice, using product rule
		A1	For correct $\frac{dy}{dx}$
	$\frac{d^2y}{dx^2} = 8p \cos 4x - 16px \sin 4x$	A1√	For unsimplified $\frac{d^2y}{dx^2}$. f.t. from $\frac{dy}{dx}$
	$\Rightarrow 8p \cos 4x = 8 \cos 4x$	M1	For substituting into DE
	$\Rightarrow p = 1$	A1	For correct p
	$\Rightarrow (y =) A \cos 4x + B \sin 4x + x \sin 4x$	B1√	6 For using GS = CF + PI, with 2 arbitrary constants in CF and none in PI

(iii)	$(0, 2) \Rightarrow A = 2$	B1√	For correct A. f.t. from their GS
	$\frac{dy}{dx} = -4A \sin 4x + 4B \cos 4x + \sin 4x + 4x \cos 4x$	M1	For differentiating their GS
	$x = 0, \frac{dy}{dx} = 0 \Rightarrow B = 0$	M1	For substituting values for x and $\frac{dy}{dx}$ to find B
	$\Rightarrow y = 2 \cos 4x + x \sin 4x$	A1 4	For stating correct solution CAO including $y =$
12			
7 (i)	$\cos 6\theta = 0 \Rightarrow 6\theta = k \times \frac{1}{2}\pi$	M1	For multiples of $\frac{1}{2}\pi$ seen or implied
	$\Rightarrow \theta = \frac{1}{12}\pi \{1, 3, 5, 7, 9, 11\}$	A1	A1 for any 3 correct
		A1 3	A1 for the rest, and no extras in $0 < \theta < \pi$
(ii)	METHOD 1		
	$\text{Re}(c + is)^6 = \cos 6\theta = c^6 - 15c^4s^2 + 15c^2s^4 - s^6$	M1	For expanding $(c + is)^6$ at least 4 terms and 2 binomial coefficients needed
	$\cos 6\theta = c^6 - 15c^4(1 - c^2) + 15c^2(1 - c^2)^2 - (1 - c^2)^3$	A1	For 4 correct terms
	$\Rightarrow \cos 6\theta = 32c^6 - 48c^4 + 18c^2 - 1$	M1	For using $s^2 = 1 - c^2$
	$\Rightarrow \cos 6\theta = (2c^2 - 1)(16c^4 - 16c^2 + 1)$	A1	For correct expression for $\cos 6\theta$
		A1 5	For correct result AG (may be written down from correct $\cos 6\theta$)
	METHOD 2		
	$\text{Re}(c + is)^3 = \cos 3\theta = \cos^3 \theta - 3\cos \theta \sin^2 \theta$	M1	For expanding $(c + is)^3$ at least 2 terms and 1 binomial coefficient needed
	$\Rightarrow \cos 6\theta = \cos 2\theta (\cos^2 2\theta - 3\sin^2 2\theta)$	A1	For 2 correct terms
	$\Rightarrow \cos 6\theta = (2\cos^2 \theta - 1) \left(4(2\cos^2 \theta - 1)^2 - 3 \right)$	M1	For replacing θ by 2θ
	$\Rightarrow \cos 6\theta = (2c^2 - 1)(16c^4 - 16c^2 + 1)$	A1	For correct expression in $\cos \theta$ (unsimplified)
		A1	For correct result AG
(iii)	METHOD 1		
	$\cos 6\theta = 0$	M1	For putting $\cos 6\theta = 0$
	$\Rightarrow 6$ roots of $\cos 6\theta = 0$ satisfy	A1	For association of roots with quartic and quadratic
	$16c^4 - 16c^2 + 1 = 0$ and $2c^2 - 1 = 0$	B1	For correct association of roots with quadratic
	But $\theta = \frac{1}{4}\pi, \frac{3}{4}\pi$ satisfy $2c^2 - 1 = 0$	M1	For using product of 4 roots
	<i>EITHER</i> Product of 4 roots <i>OR</i> $c = \pm \frac{1}{2}\sqrt{2 \pm \sqrt{3}}$	OR	For solving quartic
	$\Rightarrow \cos \frac{1}{12}\pi \cos \frac{5}{12}\pi \cos \frac{7}{12}\pi \cos \frac{11}{12}\pi = \frac{1}{16}$	A1 5	For correct value (may follow A0 and B0)

METHOD 2

$\cos 6\theta = 0$	M1	For putting $\cos 6\theta = 0$
$\Rightarrow 6$ roots of $\cos 6\theta = 0$ satisfy	A1	For association of roots with sextic
$32c^6 - 48c^4 + 18c^2 - 1 = 0$		
Product of 6 roots \Rightarrow	M1	For using product of 6 roots
$\cos \frac{1}{12}\pi \cdot \frac{1}{\sqrt{2}} \cdot \cos \frac{5}{12}\pi \cos \frac{7}{12}\pi \cdot \frac{-1}{\sqrt{2}} \cdot \cos \frac{11}{12}\pi = -\frac{1}{32}$	B1	For using $\cos\left\{\frac{3}{12}\pi, \frac{9}{12}\pi\right\} = \left\{\frac{1}{\sqrt{2}}, \frac{-1}{\sqrt{2}}\right\}$
$\cos \frac{1}{12}\pi \cos \frac{5}{12}\pi \cos \frac{7}{12}\pi \cos \frac{11}{12}\pi = \frac{1}{16}$	A1	For correct value

13

8 (i)

$g(x) = \frac{1}{2-2 \cdot \frac{1}{2-2x}} = \frac{2-2x}{2-4x} = \frac{1-x}{1-2x}$	M1	For use of $f f(x)$
	A1	For correct expression AG

$gg(x) = \frac{1 - \frac{1-x}{1-2x}}{1 - 2 \cdot \frac{1-x}{1-2x}} = \frac{-x}{-1} = x$	M1	For use of $gg(x)$
	A1 4	For correct expression AG

(ii) Order of $f = 4$	B1	For correct order
order of $g = 2$	B1 2	For correct order

(iii) METHOD 1

$y = \frac{1}{2-2x} \Rightarrow x = \frac{2y-1}{2y}$	M1	For attempt to find inverse
$\Rightarrow f^{-1}(x) = h(x) = \frac{2x-1}{2x}$ OR $1 - \frac{1}{2x}$	A1 2	For correct expression

METHOD 2

$f^{-1} = f^3 = f g$ or $g f$	M1	For use of $f g(x)$ or $g f(x)$
$f g(x) = h(x) = \frac{1}{2-2\left(\frac{1-x}{1-2x}\right)} = \frac{1-2x}{-2x}$	A1	For correct expression

(iv)

<table border="1" style="display: inline-table; border-collapse: collapse;"> <tr><td>e</td><td>f</td><td>g</td><td>h</td></tr> <tr><td>e</td><td>f</td><td>g</td><td>h</td></tr> <tr><td>f</td><td>f</td><td>g</td><td>h</td></tr> <tr><td>g</td><td>g</td><td>h</td><td>e</td></tr> <tr><td>h</td><td>h</td><td>e</td><td>f</td></tr> </table>	e	f	g	h	e	f	g	h	f	f	g	h	g	g	h	e	h	h	e	f	M1	For correct row 1 and column 1
e	f	g	h																			
e	f	g	h																			
f	f	g	h																			
g	g	h	e																			
h	h	e	f																			
	A1	For e, f, g, h in a latin square																				
	A1	For correct diagonal e - g - e - g																				
	A1 4	For correct table																				

12