

Write your name here

Surname

Other names

Pearson Edexcel
International
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--

Mechanics M2

Advanced/Advanced Subsidiary

Wednesday 2 November 2016 – Morning
Time: 1 hour 30 minutes

Paper Reference

WME02/01**You must have:**

Mathematical Formulae and Statistical Tables (Blue)

Total Marks

--

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$, and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P48103A

©2016 Pearson Education Ltd.

1/1/1/

**PEARSON**

Leave
blank

3. A particle P of mass 4 kg is projected with speed 6 ms^{-1} up a line of greatest slope of a fixed rough inclined plane. The plane is inclined at angle α to the horizontal, where $\sin\alpha = \frac{1}{7}$. The particle is projected from the point A on the plane and comes to instantaneous rest at the point B on the plane, where $AB = 10\text{ m}$.

(a) Show that the work done against friction as P moves from A to B is 16 joules. (4)

After coming to instantaneous rest at B , the particle slides back down the plane.

(b) Use the work-energy principle to find the speed of P at the instant it returns to A . (3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave blank

Question 3 continued

Lined writing area for Question 3 continued.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave blank

4. At time t seconds ($t \geq 0$), a particle P has position vector \mathbf{r} metres with respect to a fixed origin O , where

$$\mathbf{r} = \left(t^3 - \frac{9}{2}t^2 - 24t \right) \mathbf{i} + \left(-t^3 + 3t^2 + 12t \right) \mathbf{j}$$

At time T seconds, P is moving in a direction parallel to the vector $-\mathbf{i} - \mathbf{j}$

Find

- (a) the value of T , (5)

- (b) the magnitude of the acceleration of P at the instant when $t = T$. (5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



5.




Figure 1

A uniform rod AB of length 8 m and weight W newtons rests in equilibrium against a rough horizontal peg P . The end A is on rough horizontal ground. The friction is limiting at both A and P . The distance AP is 5 m, as shown in Figure 1. The rod rests at angle θ to the horizontal, where $\tan \theta = \frac{4}{3}$. The rod is in a vertical plane which is perpendicular to P . The coefficient of friction between the rod and P is $\frac{1}{4}$ and the coefficient of friction between the rod and the ground is μ .

(a) Show that the magnitude of the normal reaction between the rod and P is $0.48W$ newtons. (3)

(b) Find the value of μ . (7)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 5 continued

Handwritten student answer for Question 5, consisting of approximately 28 lines of text.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 5 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Lined writing area for the answer.

Q5

(Total 10 marks)



6.




Figure 2

The uniform lamina L shown shaded in Figure 2 is formed by removing two circular discs, C_1 and C_2 , from a circular disc with centre O and radius $8a$. Disc C_1 has centre A and radius a . Disc C_2 has centre B and radius $2a$. The diameters PR and QS are perpendicular. The midpoint of PO is A and the midpoint of OR is B .

- (a) Show that the centre of mass of L is $\frac{484}{59}a$ from R . (5)

The mass of L is M . A particle of mass kM is attached to L at S . The lamina with the attached particle is suspended from R and hangs freely in equilibrium with the diameter PR at an angle of $\arctan\left(\frac{1}{4}\right)$ to the downward vertical through R .

- (b) Find the value of k . (5)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

7. [In this question, the unit vectors \mathbf{i} and \mathbf{j} are in a vertical plane, \mathbf{i} being horizontal and \mathbf{j} being vertically upwards. Position vectors are given relative to a fixed origin O .]

At time $t = 0$ seconds, the particle P is projected from O with velocity $(3\mathbf{i} + \lambda\mathbf{j}) \text{ ms}^{-1}$, where λ is a positive constant. The particle moves freely under gravity. As P passes through the fixed point A it has velocity $(3\mathbf{i} - 4\mathbf{j}) \text{ ms}^{-1}$. The kinetic energy of P at the instant it passes through A is half the initial kinetic energy of P .

Find the position vector of A , giving the components to 2 significant figures.

(10)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave
blank

Question 7 continued

Lined writing area consisting of approximately 30 horizontal lines.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Leave blank

8. Particles A, B and C, of masses $4m$, km and $2m$ respectively, lie at rest in a straight line on a smooth horizontal surface with B between A and C. Particle A is projected towards particle B with speed $3u$ and collides directly with B. The coefficient of restitution between each pair of particles is $\frac{2}{3}$

Find

- (a) the speed of A immediately after the collision with B, giving your answer in terms of u and k , (6)

- (b) the range of values of k for which A and B will both be moving in the same direction immediately after they collide. (2)

After the collision between A and B, particle B collides directly with C.
Given that $k = 4$,

- (c) show that there will not be a second collision between A and B. (6)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Leave blank

Question 8 continued

Lined area for writing the answer to Question 8 continued.



Leave blank

Question 8 continued

Lined writing area for the answer to Question 8.

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



