Examiner's use only

Team Leader's use only

Question

1

Leave

Centre No.					Pape	er Refer	ence			Surname	Initial(s)
Candidate No.			6	6	8	1	/	0	1	Signature	

Paper Reference(s)

6681/01

Edexcel GCE

Mechanics M5

Advanced/Advanced Subsidiary

Monday 28 June 2010 – Afternoon

Time: 1 hour 30 minutes

Materials	requ	uired	for	examinatio
Mathemati	cal I	Formi	1120	(Pink)

Items included with question papers

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 6 questions in this question paper. The total mark for this paper is 75.

There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates


You must ensure that your answers to parts of questions are clearly labelled. You should show sufficient working to make your methods clear to the Examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.

 $\overset{\text{Printer's Log. No.}}{N35394A}$

W850/R6681/57570 4/5/5/3





Turn over

Total



At time $t = 0$, the position vector of a particle P is $-3\mathbf{j}$ m. At time t see	conds, the position
vector of P is \mathbf{r} metres and the velocity of P is \mathbf{v} m s ⁻¹ . Given that	, 1
$\mathbf{v} - 2\mathbf{r} = 4\mathbf{e}^t \mathbf{j} ,$	
find the time when P passes through the origin.	
	(7)



Leave blank

2.




Figure 1

A uniform circular disc has mass 4m, centre O and radius 4a. The line POQ is a diameter of the disc. A circular hole of radius 2a is made in the disc with the centre of the hole at the point R on PQ where QR = 5a, as shown in Figure 1.

The resulting lamina is free to rotate about a fixed smooth horizontal axis L which passes through Q and is perpendicular to the plane of the lamina.

(a) Show that the moment of inertia of the lamina about L is $69ma^2$.

(7)

The lamina is hanging at rest with P vertically below Q when it is given an angular velocity Ω . Given that the lamina turns through an angle $\frac{2\pi}{3}$ before it first comes to instantaneous rest,

(b) find Ω in terms	or α	and	a
----------------------------	-------------	-----	---

(6)



 . •		•	·
	Leav	Je.)

blank

- 3. A uniform lamina ABC of mass m is in the shape of an isosceles triangle with AB = AC = 5a and BC = 8a.
 - (a) Show, using integration, that the moment of inertia of the lamina about an axis through A, parallel to BC, is $\frac{9}{2}ma^2$.

(6)

The foot of the perpendicular from A to BC is D. The lamina is free to rotate in a vertical plane about a fixed smooth horizontal axis which passes through D and is perpendicular to the plane of the lamina. The lamina is released from rest when DA makes an angle α with the downward vertical. It is given that the moment of inertia of the lamina about an axis through A, perpendicular to BC and in the plane of the lamina, is $\frac{8}{3}ma^2$.

(b) Find the angular acceleration of the lamina when DA makes an angle θ with the downward vertical.

(8)

Given that α is small,

(c) find an approximate value for the period of oscillation of the lamina about the vertical.

(2)



estion 3 continued		



Leave

4.	Two forces $\mathbf{F}_1 = (\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$ N and $\mathbf{F}_2 = (3\mathbf{i} + \mathbf{j} + 2\mathbf{k})$ N act on a rigid body.
	The force \mathbf{F}_1 acts through the point with position vector $(2\mathbf{i} + \mathbf{k})$ m and the force \mathbf{F}_2 acts
	through the point with position vector $(\mathbf{j} + 2\mathbf{k})$ m.
	(a) If the two forces are equivalent to a single force D , find

(a) If the two forces are equivalent to a single force $\, R \,$, find

(1)	К,	
		(2)

(ii) a vector equation of the line of action of $\bf R$, in the form $\bf r=a+\lambda b$.

(b) If the two forces are equivalent to a single force acting through the point with position vector $(\mathbf{i}+2\mathbf{j}+\mathbf{k})$ m together with a couple of moment \mathbf{G} , find the magnitude of \mathbf{G} .

(5)



Leave

5. A raindrop falls vertically under gravity through a cloud. In a model of the motion the raindrop is assumed to be spherical at all times and the cloud is assumed to consist of stationary water particles. At time t = 0, the raindrop is at rest and has radius a. As the raindrop falls, water particles from the cloud condense onto it and the radius of the raindrop is assumed to increase at a constant rate λ . A time t the speed of the raindrop is v.

(a) Show that

$$\frac{\mathrm{d}v}{\mathrm{d}t} + \frac{3\lambda v}{(\lambda t + a)} = g.$$

(8)

(b) Find the speed of the raindrop when its radius is 3a.

(7)

estion 5 continued	



		Leave
6.	A uniform circular disc has mass m , centre O and radius $2a$. It is free to rotate about a fixed smooth horizontal axis L which lies in the same plane as the disc and which is tangential to the disc at the point A . The disc is hanging at rest in equilibrium with O vertically below A when it is struck at O by a particle of mass m . Immediately before the impact the particle is moving perpendicular to the plane of the disc with speed $3\sqrt{(ag)}$. The particle adheres to the disc at O .	blank
	(a) Find the angular speed of the disc immediately after the impact. (5)	
	(b) Find the magnitude of the force exerted on the disc by the axis immediately after the impact.	
	(6)	



estion 6 continued	

