



## Cambridge International AS & A Level

CANDIDATE  
NAME

--


CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 9 7 8 2 9 8 4 7 8 9 \*



**MATHEMATICS**

**9709/43**

Paper 4 Mechanics

**May/June 2021**

**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity ( $g$ ) is needed, use  $10 \text{ m s}^{-2}$ .

### INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **12** pages.

- 1 Particles  $P$  of mass  $0.4\text{ kg}$  and  $Q$  of mass  $0.5\text{ kg}$  are free to move on a smooth horizontal plane.  $P$  and  $Q$  are moving directly towards each other with speeds  $2.5\text{ m s}^{-1}$  and  $1.5\text{ m s}^{-1}$  respectively. After  $P$  and  $Q$  collide, the speed of  $Q$  is twice the speed of  $P$ .

Find the two possible values of the speed of  $P$  after the collision. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

## 3

- 2 A cyclist is travelling along a straight horizontal road. She is working at a constant rate of 150 W. At an instant when her speed is  $4 \text{ m s}^{-1}$ , her acceleration is  $0.25 \text{ m s}^{-2}$ . The resistance to motion is 20 N.
- (a) Find the total mass of the cyclist and her bicycle. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The cyclist comes to a straight hill inclined at an angle  $\theta$  above the horizontal. She ascends the hill at constant speed  $3 \text{ m s}^{-1}$ . She continues to work at the same rate as before and the resistance force is unchanged.

- (b) Find the value of  $\theta$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

3



Four coplanar forces act at a point. The magnitudes of the forces are 20 N, 30 N, 40 N and  $F$  N. The directions of the forces are as shown in the diagram, where  $\sin \alpha^\circ = 0.28$  and  $\sin \beta^\circ = 0.6$ .

Given that the forces are in equilibrium, find  $F$  and  $\theta$ . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 4 A particle is projected vertically upwards with speed  $u \text{ m s}^{-1}$  from a point on horizontal ground. After 2 seconds, the height of the particle above the ground is 24 m.

(a) Show that  $u = 22$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) The height of the particle above the ground is more than  $h$  m for a period of 3.6 s.

Find  $h$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

6

**5** A car of mass 1400 kg is towing a trailer of mass 500 kg down a straight hill inclined at an angle of  $5^\circ$  to the horizontal. The car and trailer are connected by a light rigid tow-bar. At the top of the hill the speed of the car and trailer is  $20 \text{ m s}^{-1}$  and at the bottom of the hill their speed is  $30 \text{ m s}^{-1}$ .

**(a)** It is given that as the car and trailer descend the hill, the engine of the car does 150 000 J of work, and there are no resistance forces.

Find the length of the hill.

[5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) It is given instead that there is a resistance force of 100 N on the trailer, the length of the hill is 200 m, and the acceleration of the car and trailer is constant.

Find the tension in the tow-bar between the car and trailer. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

## 8

- 6 A particle moves in a straight line and passes through the point  $A$  at time  $t = 0$ . The velocity of the particle at time  $t$  s after leaving  $A$  is  $v$  m s<sup>-1</sup>, where

$$v = 2t^2 - 5t + 3.$$

- (a) Find the times at which the particle is instantaneously at rest. Hence or otherwise find the minimum velocity of the particle. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Sketch the velocity-time graph for the first 3 seconds of motion. [3]



9

- (c) Find the distance travelled between the two times when the particle is instantaneously at rest. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

7



A particle  $P$  of mass  $0.3 \text{ kg}$  rests on a rough plane inclined at an angle  $\theta$  to the horizontal, where  $\sin \theta = \frac{7}{25}$ . A horizontal force of magnitude  $4 \text{ N}$ , acting in the vertical plane containing a line of greatest slope of the plane, is applied to  $P$  (see diagram). The particle is on the point of sliding up the plane.

- (a) Show that the coefficient of friction between the particle and the plane is  $\frac{3}{4}$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The force acting horizontally is replaced by a force of magnitude  $4 \text{ N}$  acting up the plane parallel to a line of greatest slope.

- (b) Find the acceleration of  $P$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Starting with  $P$  at rest, the force of 4 N parallel to the plane acts for 3 seconds and is then removed.

Find the total distance travelled until  $P$  comes to instantaneous rest. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**Additional Page**

If you use the following lined page to complete the answer(s) to any question(s), the question number(s) must be clearly shown.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.