



Mark Scheme (Results)

October 2020

Pearson Edexcel IAL Mathematics (WMA14)

Pure Mathematics P4

Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at www.edexcel.com or www.btec.co.uk. Alternatively, you can get in touch with us using the details on our contact us page at www.edexcel.com/contactus.

Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: www.pearson.com/uk

October 2020

Publications Code WMA14_01_2010_MS

All the material in this publication is copyright

© Pearson Education Ltd 2020

General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Pearson Mathematics mark schemes use the following types of marks:
 - **M** marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - **B** marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
 - ft – follow through
 - the symbol \surd or ft will be used for correct ft
 - cao – correct answer only
 - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
 - isw – ignore subsequent working
 - awrt – answers which round to
 - SC: special case
 - oe – or equivalent (and appropriate)
 - d... or dep – dependent
 - indep – independent
 - dp decimal places
 - sf significant figures
 - * The answer is printed on the paper or ag- answer given
 - \square or d... The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected. If you are using the annotation facility on ePEN, indicate this action by 'MR' in the body of the script.
6. If a candidate makes more than one attempt at any question:
 - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
 - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.

General Principles for Core Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

Method mark for solving 3 term quadratic:

1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$, where $|pq| = |c|$, leading to $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$, where $|pq| = |c|$ and $|mn| = |a|$, leading to $x = \dots$

2. Formula

Attempt to use correct formula (with values for a , b and c).

3. Completing the square

Solving $x^2 + bx + c = 0$: $(x \pm \frac{b}{2})^2 \pm q \pm c$, $q \neq 0$, leading to $x = \dots$

Method marks for differentiation and integration:

1. Differentiation

Power of at least one term decreased by 1. ($x^n \rightarrow x^{n-1}$)

2. Integration

Power of at least one term increased by 1. ($x^n \rightarrow x^{n+1}$)

Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are small mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

Exact answers

Examiners' reports have emphasised that where, for example, an exact answer is asked for, or working with surds is clearly required, marks will normally be lost if the candidate resorts to using rounded decimals.

Answers without working

The rubric says that these may not gain full credit. Individual mark schemes will give details of what happens in particular cases. General policy is that if it could be done "in your head", detailed working would not be required. Most candidates do show working, but there are occasional awkward cases and if the mark scheme does not cover this, please contact your team leader for advice

Question Number	Scheme	Marks
1	<p>Assume that there exists a number m such that when m^3 is even, m is odd</p> <p>If m is odd then $m = 2p+1$ (where p is an integer) and $m^3 = (2p+1)^3 = \dots$</p> $= 8p^3 + 12p^2 + 6p + 1$ <p>$2 \times (4p^3 + 6p^2 + 3p) + 1$ is odd and hence we have a contradiction so if n^3 is even, then n is even.</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>(4)</p> <p>(4 marks)</p>

B1: For setting up the contradiction.

Eg Assume that there exists a number m such that when m^3 is even, m is odd

Condone a contra-positive statement here

"Assume that there exists a number m such that when m^3 is even, m is not even"

As a minimum accept "assume if m^3 is even then m is odd."

Condone the other way around "assume if n is odd then n^3 is even"

M1: Attempts to cube an odd number. Accept an attempt at $(2p+1)^3$, $(2p-1)^3$

Look for $(2p+1)^3 = \dots p^3 \dots$

A1: $(2p+1)^3 = 8p^3 + 12p^2 + 6p + 1$ or simplified equivalent such as $2 \times (4p^3 + 6p^2 + 3p) + 1$.

For $(2p-1)^3 = 8p^3 - 12p^2 + 6p - 1$ or equivalent such as $2 \times (4p^3 - 6p^2 + 3p - 1) + 1$

A1: For a fully correct proof. Requires correct calculations with reason and conclusion

E.g. 1 Correct calculations $(2p+1)^3 = 8p^3 + 12p^2 + 6p + 1 =$

Reason (even + 1) = odd

Conclusion "hence we have a contradiction, so if n^3 is even, then n is even."

E.g. 2 Correct calculations $(2p+1)^3 = 8p^3 + 12p^2 + 6p + 1$

Reason $= 2 \times (4p^3 + 6p^2 + 3p) + 1 =$ odd

Conclusion "this is contradiction, so proven."

E.g. 3 Correct calculations $(2p-1)^3 = 8p^3 - 12p^2 + 6p - 1$

Reason $= 8p^3 - 12p^2 + 6p$ is even so $8p^3 - 12p^2 + 6p - 1$ is odd

Conclusion: So if n^3 is even then n must be even

Note that B0 M1 A1 A1 is possible

Question Number	Scheme	Marks
<p>2(a)</p>	$4^{-\frac{1}{2}} \text{ or } \frac{1}{4^{\frac{1}{2}}} \text{ or } \frac{1}{2}$ $(4-5x)^{-\frac{1}{2}} = \frac{1}{2} \left(1 - \frac{5x}{4}\right)^{-\frac{1}{2}}$ $= \dots \left[1 + \left(-\frac{1}{2}\right) \times \left(-\frac{5x}{4}\right) + \frac{\left(-\frac{1}{2}\right) \times \left(-\frac{3}{2}\right)}{2!} \times \left(-\frac{5x}{4}\right)^2 + \dots \right]$ $= \frac{1}{2} + \frac{5}{16}x + \frac{75}{256}x^2 + \dots$ $\frac{2+kx}{(2-3x)^3} = (2+kx) \left(\frac{1}{2} + \frac{5}{16}x + \frac{75}{256}x^2 + \dots \right)$ <p>(b) Compares x terms leading to $k = \dots$ E.g. $\frac{10}{16} + \frac{k}{2} = \frac{3}{10} \Rightarrow k = -\frac{13}{20}$</p> <p>(c) Compares x^2 terms leading to $m = \dots$ E.g. $m = \frac{75}{128} + \frac{5}{16} \times -\frac{13}{20} \Rightarrow m = \frac{49}{128}$</p>	<p>B1</p> <p>M1A1</p> <p>A1</p> <p>(4)</p> <p>M1 A1</p> <p>(2)</p> <p>M1 A1</p> <p>(2)</p> <p>(8 marks)</p>
<p>Alt (a)</p>	$4^{-\frac{1}{2}} \text{ or } \frac{1}{4^{\frac{1}{2}}}$ $(4-5x)^{-\frac{1}{2}} = 4^{-\frac{1}{2}} + \left(-\frac{1}{2}\right) 4^{-\frac{3}{2}} (-5x) + \frac{-\frac{1}{2} \times -\frac{3}{2}}{2} 4^{-\frac{5}{2}} (-5x)^2$ $= \frac{1}{2} + \frac{5}{16}x + \frac{75}{256}x^2 + \dots$	<p>B1</p> <p>M1A1</p> <p>A1</p> <p>(4)</p>

(a)

B1: For taking out a factor of $4^{-\frac{1}{2}}$ or $\frac{1}{2}$

For a direct expansion look for $4^{\frac{1}{2}} + \dots$ or equivalent.

M1: For the form of the binomial expansion $(1+ax)^{-\frac{1}{2}}$ where $a \neq 1$ or -5

To score M1 it is sufficient to see either term two or term three. Allow a slip on the sign.

So allow for either $\left(-\frac{1}{2}\right)(\pm ax)$ or $\frac{\left(-\frac{1}{2}\right) \times \left(-\frac{3}{2}\right)}{2}(\pm ax)^2$

In the alternative version look for $\left(-\frac{1}{2}\right)4^{-\frac{3}{2}}(-5x)$ or $\frac{-\frac{1}{2} \times -\frac{3}{2}}{2}4^{-\frac{5}{2}}(-5x)^2$ condoning sign slips

A1: Any (unsimplified) but correct form of the binomial expansion for $\left(1-\frac{5x}{4}\right)^{-\frac{1}{2}}$

Ignore the factor preceding the bracket for this mark

Score for $1 + \left(-\frac{1}{2}\right) \times \left(-\frac{5x}{4}\right) + \frac{\left(-\frac{1}{2}\right) \times \left(-\frac{3}{2}\right)}{2!} \times \left(-\frac{5x}{4}\right)^2$ o.e.

In the alternative version look for $(4-5x)^{-\frac{1}{2}} = 4^{-\frac{1}{2}} + \left(-\frac{1}{2}\right)4^{-\frac{3}{2}}(-5x) + \frac{-\frac{1}{2} \times -\frac{3}{2}}{2}4^{-\frac{5}{2}}(-5x)^2$

A1: cao $\frac{1}{2} + \frac{5}{16}x + \frac{75}{256}x^2 + \dots$ This must be simplified

$$(2+kx)(P+Qx+Rx^2+\dots) = 1 + \frac{3}{10}x + mx^2$$

(b)

M1: For a correct equation in k formed by comparing the x terms. It must lead to a value for k

Follow through on their expansion. So look for $Pk + 2Q = \frac{3}{10} \Rightarrow k = \dots$ Condone slips, i.e copying errors.

Condone $Pkx + 2Qx = \frac{3}{10}x$ as long as it leads to a value for k

A1: $k = -\frac{13}{20}$

(c)

M1: Correctly compares the x^2 terms, following through on their expansion and their value for k leading to a value for m . Condone slips, i.e copying errors.

Look for $Qk + 2R = m$ Condone $Qkx^2 + 2Rx^2 = mx^2$ as long as it leads to a value for m

A1: $m = \frac{49}{128}$ oe Condone sight of $\frac{49}{128}x^2$ as evidence for a correct value for m .

Question Number	Scheme	Marks
3	Calculates the upper limit as $2\ln 2$ or $\ln 4$ $\int (e^{0.5x} - 2)^2 dx = \int (e^x - 4e^{0.5x} + 4) dx = e^x - 8e^{0.5x} + 4x$ $\text{Volume} = \pi \left[e^x - 8e^{0.5x} + 4x \right]_0^{2\ln 2} = \dots$ $\text{Volume} = 8\pi \ln 2 - 5\pi$	B1 M1 A1 A1 dM1 A1 (6 marks)

B1: Calculates the upper limit as $2\ln 2$ or $\ln 4$ only. Do not accept 1.386, $\frac{\ln 2}{0.5}$ or anything else.

Recovery from $\frac{\ln 2}{0.5}$ is allowed so if the candidate progresses to $8\pi \ln 2 - 5\pi$ do not withhold this mark.

M1: Attempts to square $(e^{0.5x} - 2)$ **with** $\int e^{kx} dx \rightarrow e^{kx}$ seen at least once. Ignore any factor of π

This may be awarded when candidates fails to square $e^{0.5x}$ correctly

So for example $\int (e^{0.5x})^2 - 4e^{0.5x} + 4 dx \rightarrow \dots \pm \dots e^{0.5x}$ is sufficient

A1: Two terms correct $e^x - 8e^{0.5x} + 4x$ Ignore any factor of π

A1: All three terms correct $e^x - 8e^{0.5x} + 4x$ Ignore any factor of π

dM1: A full and complete method of finding the volume of the solid generated.

Look for an attempt to find $\pi \int_0^a (e^{0.5x} - 2)^2 dx$ with a being $2\ln 2$, $\ln 4$, 1.38.. or $\frac{\ln 2}{2}$

It is dependent upon the previous M so some correct integration must be seen as well as use of both limits. There must be some evidence for the 0. Those terms cannot just "disappear" or be set to 0.

A1: cao $8\pi \ln 2 - 5\pi$ or $-5\pi + 8\pi \ln 2$ Condone $\pi(8\ln 2 - 5)$

(Note that a requirement of the question is to write in the form $a \ln 2 + b$)

Question Number	Scheme	Marks
4(a)	A and B are where $t^3 - 4t = 0 \Rightarrow t(t^2 - 4) = 0 \Rightarrow t = 2$ or -2 Substitutes $t = 2, x = 2 \times 4 - 6 \times 2 = -4$ Hence $A = (-4, 0)$ When $t = -2, x = 2 \times 4 - 6 \times -2 = 20, (y = 0)$ Hence $B = (20, 0)$ *	M1 A1 B1* (3)
(b)	$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{3t^2 - 4}{4t - 6}$ Sub $t = -2$ into $\frac{dy}{dx} = \frac{3t^2 - 4}{4t - 6} \Rightarrow \text{gradient} = \left(-\frac{4}{7}\right)$ Uses their $\left(-\frac{4}{7}\right)$ and $(20, 0)$ to produce eqn of tangent $\Rightarrow 7y + 4x - 80 = 0$ *	M1A1 M1 M1 A1* (5)
(c)	Substitutes $x = 2t^2 - 6t, y = t^3 - 4t$, into $7y + 4x - 80 = 0$ $\Rightarrow 7(t^3 - 4t) + 4(2t^2 - 6t) - 80 = 0$ $\Rightarrow 7t^3 + 8t^2 - 52t - 80 = 0$ $\Rightarrow (t + 2)^2(7t - 20) = 0$ $t = \frac{20}{7} \Rightarrow x = \dots$ $x = -\frac{40}{49}$	M1 A1 dM1 A1 (4) (12 marks)

(a)

M1: Sets $t^3 - 4t = 0 \Rightarrow t = \dots$ to reach $t = 2$ or -2 A1: Substitutes $t = 2 \Rightarrow x = 2 \times 4 - 6 \times 2 = -4$ and states coordinates of $A = (-4, 0)$ Accept $t = 2 \Rightarrow A = (-4, 0)$ provided the M1 has been awardedB1*: Show that the coordinates of $B = (20, 0)$

Possible ways of this are

1) Solves $t^3 - 4t = 0 \Rightarrow t = -2$ and substitutes into $2t^2 - 6t \Rightarrow x = 20 \Rightarrow B = (20, 0)$.This is a show that and there must be some evidence for the "20", not just sight of $t = -2$ Look for $2 \times (-2)^2 - 6 \times (-2) \Rightarrow x = 20$ or $8 + 12 \Rightarrow x = 20$

2) Alternatively sets $2t^2 - 6t = 20 \Rightarrow t = -2$ and substitutes into $y = t^3 - 4t \rightarrow y = 0 \Rightarrow B = (20, 0)$

This is a show that and there must be some evidence for the "0", not just sight of $t = -2$

(b)

M1: Attempts to differentiate $x(t)$ and $y(t)$ and calculates $\frac{dy}{dx}$ by using $\frac{dy/dt}{dx/dt}$ in part (b)

A1: $\frac{dy}{dx} = \frac{3t^2 - 4}{4t - 6}$

M1: Sub 'their' $t = -2$ into their $\frac{dy}{dx}$ to find the gradient of the tangent at B .

M1: Uses $(20, 0)$ and their $\frac{dy}{dx}$ to find an equation of the tangent.

If they use $y = mx + c$ they need to proceed as far as $c = \dots$

A1*: Achieves given answer. $cs0 \Rightarrow 7y + 4x - 80 = 0$ The $= 0$ must be seen

ISW after sight of this

(c) **The demand here is to use algebra which is satisfied by the setting up of the cubic equation.**

M1: Attempts to substitute $x = 2t^2 - 6t$, $y = t^3 - 4t$ into $7y + 4x - 80 = 0$

A1: Correct simplified equation in t : $7t^3 + 8t^2 - 52t - 80 = 0$ The $= 0$ may be implied by further work

dM1: Uses a correct method to find x

- Using a calculator this can be scored for finding a value of $t \neq -2$ and substituting this in $x(t)$. You may have to check this on a calculator if their equation is incorrect. An accuracy of 2 sf is required for non exact solutions. Note that the equation may be in different forms under this method so $7t^3 + 8t^2 - 52t = 80 \Rightarrow t = \frac{20}{7}$ is fine

Note: $7(t^3 - 4t) + 4(2t^2 - 6t) - 80 = 0 \Rightarrow t = \frac{20}{7} \Rightarrow x = -\frac{40}{49}$ would score M1 A0 dM1 A0

- Via factorisation look for factors where the ends work, E.g. " $7t^3 + 8t^2 - 52t - 80$ " = $(Pt + a)(Qt + b)(Rt + c)$ with $PQR = "7"$ and $abc = "\pm 80"$ followed by substitution of a value of $t \neq -2$ in $x(t)$

A1: CSO $x = -\frac{40}{49}$ Do not accept decimals here but remember to isw following a correct answer.

Ignore any references to the y coordinate. Penalise extra solutions, incorrect factorisation etc on this mark.

Question Number	Scheme	Marks
5(a)	$\int \frac{\ln x}{x^2} dx = \int x^{-2} \ln x dx = -x^{-1} \ln x + \int x^{-1} \times \frac{1}{x} dx$ $= -x^{-1} \ln x - x^{-1} (+c)$	M1 dM1 A1 (3)
(b)	$\frac{3+2x+\ln x}{x^2} = 3x^{-2} + 2x^{-1} + x^{-2} \ln x \text{ oe}$ $\text{Area} = \left[-\frac{3}{x} + 2 \ln x - \frac{\ln x}{x} - \frac{1}{x} \right]_2^4$ $= \left(-\frac{3}{4} + 2 \ln 4 - \frac{\ln 4}{4} - \frac{1}{4} \right) - \left(-\frac{3}{2} + 2 \ln 2 - \frac{\ln 2}{2} - \frac{1}{2} \right) = 1 + 2 \ln 2$	B1 M1 M1 A1 (4) (7 marks)

(a)

M1: Attempts by parts to reach a form $\int x^{-2} \ln x dx = \pm ax^{-1} \ln x \pm b \int x^{-1} \times \frac{1}{x} dx$ where $a, b \neq 0$

If a formula is stated it must be correct.

dM1: Integrates again to reach a form $\pm ax^{-1} \ln x \pm bx^{-1}$

A1: $-x^{-1} \ln x - x^{-1} (+c)$ o.e such as $-\frac{\ln x}{x} - \frac{1}{x} + c$

Condone the omission of the constant of integration. Condone $-x^{-1} \ln x + (-x^{-1})$

(b)

B1: $\frac{3+2x+\ln x}{x^2} = 3x^{-2} + 2x^{-1} + x^{-2} \ln x$ or $\frac{3}{x^2} + \frac{2}{x} + \frac{\ln x}{x^2}$.

The $\frac{2}{x}$ may be implied by later work. That is if they state $\int \frac{2x}{x^2} dx = 2 \ln x$ or $\ln x^2$

M1: Area = $\left[-\frac{3}{x} + 2 \ln x - \frac{\ln x}{x} - \frac{1}{x} \right]_2^4$ following through on their part (a)

M1: For substituting 2 and 4 **into an expression of the form** $\left[\frac{p}{x} + q \ln x + r \frac{\ln x}{x} + \frac{s}{x} \right]_2^4$ and attempts to

simplify by using one log law correctly. (Note that the two terms in $\frac{1}{x}$ may be combined before substitution)

A1: Exact answer: Likely to be $1 + 2 \ln 2$ or $1 + \ln 4$

.....
For those who don't read the instruction in (b) to use (a) and do via parts score

$$B1: \int \frac{3+2x+\ln x}{x^2} dx = -x^{-1}(3+2x+\ln x) + \int \left(2 + \frac{1}{x}\right) \times \frac{1}{x} dx$$

$$M1: \text{Area} = \left[\pm x^{-1}(3+2x+\ln x) \pm 2 \ln x \pm x^{-1} \right]_2^4$$

Question Number	Scheme	Marks
6 (a)	$y = x^{\sin x} \Rightarrow \ln y = \sin x \ln x$ Differentiates $\frac{1}{y} \frac{dy}{dx} = \frac{\sin x}{x} + \ln x \cos x$ $\frac{dy}{dx} = \frac{y \sin x}{x} + y \ln x \cos x$ oe	B1 M1 M1 A1 A1 (5)
(b)	Puts $\frac{dy}{dx} = 0 \Rightarrow \frac{\sin x}{x} + \ln x \cos x = 0$ $\frac{\sin x}{\cos x} + x \ln x = 0 \Rightarrow \tan x + x \ln x = 0^*$	M1 A1* (2)
		(7 marks)

(a)

B1: $y = x^{\sin x} \Rightarrow \ln y = \sin x \ln x$ o.e. Condone this written $\log y = \sin x \log x$

This may have been found via $\log_x y = \sin x \rightarrow \frac{\ln y}{\ln x} = \sin x$

M1: For differentiation of $\ln y \rightarrow \frac{1}{y} \times \frac{dy}{dx}$ or $\log_e y \rightarrow \frac{1}{y} \times \frac{dy}{dx}$

M1: Attempts the product rule on $\sin x \ln x$ If the rule is quoted it must be correct.

If candidates set $u = \dots, u' = \dots, v = \dots, v' = \dots$ it may be implied by their $vu' + uv'$

If it is neither quoted nor implied score for $\frac{\sin x}{x} \pm \ln x \cos x$

A1: Correct differentiation to $\frac{1}{y} \frac{dy}{dx} = \frac{\sin x}{x} + \ln x \cos x$

A1: Achieves $\frac{dy}{dx} = \frac{y \sin x}{x} + y \ln x \cos x$ or exact equivalent such as $\frac{dy}{dx} = y \left(\frac{\sin x + x \ln x \cos x}{x} \right)$

Accept this in terms of just x . For instance $\frac{dy}{dx} = \frac{x^{\sin x} \sin x}{x} + x^{\sin x} \ln x \cos x$

ISW after sight of a correct answer.

.....
 Note that you will see candidates who write $\frac{1}{y} \frac{dy}{dx} = \frac{\sin x}{x} dx + \ln x \cos x dx$ which can be marked in the same

way

(b)

M1: Sets their $\frac{dy}{dx} = 0$, divides by y or $x^{\sin x}$ to form a simplified equation in x .

This may be implied by $\frac{dy}{dx} = \frac{y \sin x}{x} + y \ln x \cos x = 0 \Rightarrow \sin x + x \ln x \cos x = 0$ o.e.

For this to be scored $\frac{dy}{dx}$ **must be in the form**

$$\frac{y \sin x}{\dots} \pm y \ln x \cos x$$

$$\frac{x^{\sin x} \sin x}{\dots} \pm x^{\sin x} \ln x \cos x \text{ where } \dots \text{ could be } 1$$

A1*: CSO Proceeds to the given answer of $\tan x + x \ln x = 0$

showing at least the intermediate line $\frac{\sin x}{\cos x} + x \ln x = 0$

It can be implied by $\sin x + x \ln x \cos x = 0$ ($\div \cos x$) $\Rightarrow \tan x + x \ln x = 0$

6 (a)	$y = x^{\sin x} \Rightarrow y = e^{\sin x \ln x}$	B1
	Differentiates $y = e^{\sin x \ln x} \rightarrow \frac{dy}{dx} = e^{\sin x \ln x} \times \dots$	M1
	$\frac{d}{dx}(\sin x \ln x) = \frac{\sin x}{x} + \ln x \cos x$	M1
	$\frac{dy}{dx} = e^{\sin x \ln x} \times \left(\frac{\sin x}{x} + \ln x \cos x \right)$	A2
		(5)

B1: $y = x^{\sin x} \Rightarrow y = e^{\sin x \ln x}$

M1: Differentiates $e^{\sin x \ln x} \rightarrow e^{\sin x \ln x} \times \dots$ where ... could be 1

M1: Attempts the product rule on $\sin x \ln x$. **See main scheme on how to apply this mark.**

A2: Correct differentiation = $\frac{\sin x}{x} + \ln x \cos x$

Question	Scheme	Marks
7 (i)	<p>Chooses a suitable substitution for $\int_1^5 \frac{3x}{\sqrt{2x-1}} dx$</p> <p>E.g. $u = \sqrt{2x-1}$ $u = \sqrt{2x-1} \Rightarrow \frac{dx}{du} = u$</p> $\int \frac{3x}{\sqrt{2x-1}} dx$ $= \int \frac{3}{2}u^2 + \frac{3}{2} du$ $= \frac{1}{2}u^3 + \frac{3}{2}u$ <p>Limits = $\left[\frac{1}{2}u^3 + \frac{3}{2}u \right]_1^3 = \dots$</p> $\left(\frac{1}{2} \times 3^3 + \frac{3}{2} \times 3 \right) - \left(\frac{1}{2} \times 1^3 + \frac{3}{2} \times 1 \right) = 16$	<p>B1</p> <p>M1</p> <p>dM1 A1</p> <p>M1</p> <p>A1</p> <p>(6)</p>
7 (ii)	$\frac{6x^2 - 16}{(x+1)(2x-3)} = 3 + \dots$ $\frac{6x^2 - 16}{(x+1)(2x-3)} = 3 + \frac{2}{x+1} - \frac{1}{2x-3}$ $\int \frac{6x^2 - 16}{(x+1)(2x-3)} dx = \int 3 + \frac{2}{x+1} - \frac{1}{2x-3} dx$ $= 3x + 2 \ln x+1 - \frac{1}{2} \ln 2x-3 + c$	<p>B1</p> <p>M1 A1</p> <p>M1 A1ft A1</p> <p>(6)</p> <p>(12 marks)</p>

(i)

B1: States or uses a suitable substitution. That is one that will work.

M1: Attempts to change the integral in x to an integral in ' u 'Expect to see all aspects changed including the dx For $u = \sqrt{2x-1}$ expect to see $\int Au^2 + Bdu$

For $u = 2x - 1$ expect to see $\int Pu^{\frac{1}{2}} + Qu^{-\frac{1}{2}} du$

dM1: For correct method of integration. Allow for correct powers. Dependent upon previous M

A1: See scheme. Correct integration which may be left unsimplified

M1: Uses correct limits for their **valid** substitution.

A1: CSO 16. Look for 18 -2 or such before the 16.

16 with no working scores no marks. This is evidence of use of a calculator

(ii) **Note that open scoring is M1 M1 A1 M1 A1 A1**

B1: For a quotient of 3.

This can be scored by setting $6x^2 - 16 = A(x+1)(2x-3) + B(x+1) + C(2x-3)$ and reaching $A = 3$
or by division with 3 in the "correct place"

M1: A full attempt to find A , B and C .

Either sets $6x^2 - 16 = A(x+1)(2x-3) + B(x+1) + C(2x-3)$ condoning slips leading to values of A , B and C
or divides and uses partial fractions on the remainder to form an expression of the correct form.

A1: Correct partial fractions seen or implied. $\frac{6x^2 - 16}{(x+1)(2x-3)} = 3 + \frac{2}{x+1} - \frac{1}{2x-3}$

M1: Two of $A \rightarrow \dots x$, $\int \frac{B}{x+1} dx \rightarrow \dots \ln(x+1)$ $\int \frac{C}{2x-3} dx \rightarrow \dots \ln(2x-3)$

A1ft: Two correct of $A \rightarrow Ax$ $\int \frac{B}{x+1} dx \rightarrow B \ln(x+1)$ and $\int \frac{C}{2x-3} dx \rightarrow \frac{C}{2} \ln(2x-3)$

Please note that answers such as $\int \frac{1}{2(2x-3)} dx \rightarrow \frac{1}{2} \ln(4x-6)$ are correct

A1: $3x + 2 \ln|x+1| - \frac{1}{2} \ln|2x-3| + c$ but condone $3x + 2 \ln(x+1) - \frac{1}{2} \ln(2x-3) + c$ **The $+c$ is required.**

You should isw after a correct answer.

Note that it is possible to do **part (i)** by parts. This does not satisfy the demand of the question but can score some marks

B1: $\int \frac{3x}{\sqrt{2x-1}} dx = 3x\sqrt{2x-1} - \int 3\sqrt{2x-1} dx$

M1: $\int \frac{3x}{\sqrt{2x-1}} dx = ax\sqrt{2x-1} - b(2x-1)^{\frac{3}{2}}$ FYI: Correct when $a = 3, b = 1$

third M1 (5th mark on open): applies limits to a valid expression $\int_1^5 \frac{3x}{\sqrt{2x-1}} dx = \left[ax\sqrt{2x-1} - b(2x-1)^{\frac{3}{2}} \right]_1^5 = \dots$

second A1 (6th mark on open) $\int_1^5 \frac{3x}{\sqrt{2x-1}} dx = \left[3x\sqrt{2x-1} - (2x-1)^{\frac{3}{2}} \right]_1^5 = (45-27) - (3-1) = 16$ o.e.

to score on e-open 110011

Note that in **part (ii)** some candidates will write

$$\frac{6x^2 - 16}{(x+1)(2x-3)} = \frac{B}{x+1} + \frac{C}{2x-3} \text{ and via substitution will get "a correct" } \frac{2}{x+1} - \frac{1}{2x-3}$$

$$\int \frac{6x^2 - 16}{(x+1)(2x-3)} dx = \int \frac{"B"}{x+1} + \frac{"C"}{2x-3} dx = "B" \ln(x+1) + \frac{"C"}{2} \ln(2x-3)$$

Scores SC B0 M0 A0 M1 A1ft A0 for 2 out of 6 if ft integration is correct

Question Number	Scheme	Marks
8 (a)	$\begin{pmatrix} 4 \\ -3 \\ 2 \end{pmatrix} + \lambda \begin{pmatrix} 3 \\ -2 \\ -1 \end{pmatrix} = \begin{pmatrix} 2 \\ 0 \\ -9 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ -1 \\ -3 \end{pmatrix} \Rightarrow \begin{array}{l} 4 + 3\lambda = 2 + 2\mu \\ -3 - 2\lambda = 0 - 1\mu \\ 2 - 1\lambda = -9 - 3\mu \end{array} \text{ any two of these}$	M1
	<p>Full method to find either λ or μ eg (1) + 3(3) $\Rightarrow \mu = \dots$ Either $\lambda = -4$ or $\mu = -5$</p> <p>Position vector of intersection is $\begin{pmatrix} 4 \\ -3 \\ 2 \end{pmatrix} - 4 \begin{pmatrix} 3 \\ -2 \\ -1 \end{pmatrix} = \text{OR} \begin{pmatrix} 2 \\ 0 \\ -9 \end{pmatrix} - 5 \begin{pmatrix} 2 \\ -1 \\ -3 \end{pmatrix} =$</p> $= \begin{pmatrix} -8 \\ 5 \\ 6 \end{pmatrix}$	M1 A1 dM1
(b)	Co-ordinates or position vector of point $Q = \begin{pmatrix} 2 + 2\mu \\ 0 - 1\mu \\ -9 - 3\mu \end{pmatrix}$	
	$\overrightarrow{PQ} = \begin{pmatrix} 2 + 2\mu \\ 0 - 1\mu \\ -9 - 3\mu \end{pmatrix} - \begin{pmatrix} 10 \\ -7 \\ 0 \end{pmatrix} = \begin{pmatrix} 2\mu - 8 \\ 7 - 1\mu \\ -9 - 3\mu \end{pmatrix}$	M1
	Uses $\overrightarrow{PQ} \cdot \begin{pmatrix} 2 \\ -1 \\ -3 \end{pmatrix} = 0 \Rightarrow 4\mu - 16 - 7 + \mu + 27 + 9\mu = 0 \Rightarrow \mu = -\frac{2}{7}$	dM1 A1
Substitutes their μ into $\begin{pmatrix} 2 + 2\mu \\ 0 - 1\mu \\ -9 - 3\mu \end{pmatrix} \Rightarrow Q = \left(\frac{10}{7}, \frac{2}{7}, -\frac{57}{7} \right)$	ddM1 A1	
		(5) (10 marks)

(a)

M1: For writing down **any two equations** that give the coordinates of the point of intersection.Accept two of $4 + 3\lambda = 2 + 2\mu$, $-3 - 2\lambda = 0 - 1\mu$, $2 - 1\lambda = -9 - 3\mu$

There must be an attempt to set the coordinates equal but condone one slip.

Setting the vectors equal to each other is insufficient for this but correct calculations will imply this.

M1: A full method to find **either** λ **or** μ . There are a few answers with limited or no working.In such cases this M mark can be implied only if **either** λ **or** μ are correct for their equations.**A1:** Either value correct $\mu = -5$ or $\lambda = -4$. Correct value(s) following correct equations implies M1 A1**dm1:** Substitutes their value of λ into l_1 to find the coordinates or position vector of the point of intersection.It is dependent upon having scored second method mark. Alternatively substitutes their value of μ into l_2 to find the coordinates or position vector of the point of intersection. It can be implied in cases where there is limited working by two correct coordinates for their λ **or** μ .**A1:** Either $\begin{pmatrix} -8 \\ 5 \\ 6 \end{pmatrix}$ or $-8\mathbf{i} + 5\mathbf{j} + 6\mathbf{k}$ but not the coordinate $(-8, 5, 6)$ However ISW after sight of the correct **vector****

(b)

M1: States or uses a general point on $l_2 = \begin{pmatrix} 2 + 2\mu \\ 0 - 1\mu \\ -9 - 3\mu \end{pmatrix}$ and attempt $\overrightarrow{PQ} = \begin{pmatrix} 2 + 2\mu \\ 0 - 1\mu \\ -9 - 3\mu \end{pmatrix} - \begin{pmatrix} 10 \\ -7 \\ 0 \end{pmatrix} = \begin{pmatrix} 2\mu - 8 \\ 7 - 1\mu \\ -9 - 3\mu \end{pmatrix}$ either

way around with their general point. Condone slips

dm1: Uses the scalar product with fact that \overrightarrow{PQ} and $\begin{pmatrix} 2 \\ -1 \\ -3 \end{pmatrix}$ are perpendicular vectors to find μ

See scheme but again condone slips.

Alternatively uses the scalar product with fact that \overrightarrow{PQ} and \overrightarrow{XQ} are perpendicular vectors to find μ

$$\overrightarrow{PQ} \cdot \overrightarrow{XQ} = 0 \Rightarrow (2\mu - 8)(2\mu + 10) + (7 - \mu)(-\mu - 5) + (-9 - 3\mu)(-15 - 3\mu) = 0 \Rightarrow \mu = \left(-\frac{2}{7}\right)$$

A1: $\mu = -\frac{2}{7}$ **ddM1:** Uses their μ to find the coordinates of Q .If no method is seen imply by two correct coordinates for their μ **A1:** $Q = \left(\frac{10}{7}, \frac{2}{7}, -\frac{57}{7}\right)$ but not (**) $\frac{10}{7}\mathbf{i} + \frac{2}{7}\mathbf{j} - \frac{57}{7}\mathbf{k}$ o.e**** Only penalise incorrect notation once, the first time that it occurs. ****

Alt using distances

<p>(b)</p>	<p>Co-ordinates or position vector of point $Q = \begin{pmatrix} 2+2\mu \\ 0-1\mu \\ -9-3\mu \end{pmatrix}$</p> $\overrightarrow{PQ} = \begin{pmatrix} 2+2\mu \\ 0-1\mu \\ -9-3\mu \end{pmatrix} - \begin{pmatrix} 10 \\ -7 \\ 0 \end{pmatrix} = \begin{pmatrix} 2\mu-8 \\ 7-1\mu \\ -9-3\mu \end{pmatrix}$ <p>Uses Pythagoras with</p> $PQ^2 + QX^2 = PX^2 \Rightarrow 28\mu^2 + 148\mu + 544 = 504 \Rightarrow \mu = \cancel{8}, -\frac{2}{7}$ <p>Substitutes their μ into $\begin{pmatrix} 2+2\mu \\ 0-1\mu \\ -9-3\mu \end{pmatrix} \Rightarrow Q = \left(\frac{10}{7}, \frac{2}{7}, -\frac{57}{7}\right)$</p>	<p>M1</p> <p>dM1 A1</p> <p>ddM1 A1</p>
-------------------	---	--

Alt using minimum distance and differentiation

<p>(b)</p>	<p>Co-ordinates or position vector of point $Q = \begin{pmatrix} 2+2\mu \\ 0-1\mu \\ -9-3\mu \end{pmatrix}$</p> $\overrightarrow{PQ} = \begin{pmatrix} 2+2\mu \\ 0-1\mu \\ -9-3\mu \end{pmatrix} - \begin{pmatrix} 10 \\ -7 \\ 0 \end{pmatrix} = \begin{pmatrix} 2\mu-8 \\ 7-1\mu \\ -9-3\mu \end{pmatrix}$ <p>Uses Pythagoras with</p> $PQ^2 = (2\mu-8)^2 + (7-\mu)^2 + (-9-3\mu)^2$ $\frac{d}{d\mu} PQ^2 = 0 \Rightarrow 4(2\mu-8) - 2(7-\mu) - 6(-9-3\mu) = 0 \Rightarrow \mu = -\frac{2}{7}$ <p>Substitutes their μ into $\begin{pmatrix} 2+2\mu \\ 0-1\mu \\ -9-3\mu \end{pmatrix} \Rightarrow Q = \left(\frac{10}{7}, \frac{2}{7}, -\frac{57}{7}\right)$</p>	<p>M1</p> <p>dM1 A1</p> <p>ddM1 A1</p>
-------------------	---	--

Question Number	Scheme	Marks
9 (a)	$\frac{dA}{dt} = \frac{A^{\frac{3}{2}}}{5t^2} \Rightarrow \int \frac{dA}{A^{\frac{3}{2}}} = \int \frac{dt}{5t^2}$ oe	M1
	$-2A^{-\frac{1}{2}} = -\frac{1}{5}t^{-1}(+c)$	M1 M1 A1
	Substitutes $t = 3, A = 2.25 \Rightarrow c = \dots \left(-\frac{19}{15}\right)$	M1
	Uses their $\frac{2}{\sqrt{A}} = \frac{1}{5t} + \frac{19}{15} \Rightarrow A = \left(\frac{30t}{19t+3}\right)^2$	M1, A1
		(7)
(b)	As $t \rightarrow \infty, A \rightarrow \left(\frac{30}{19}\right)^2 = \frac{900}{361}$ or awrt 2.49 cm ²	M1 A1
		(2)
		(9 marks)

(a)

M1: Attempts to separate the variables either $\int \frac{dA}{A^{\frac{3}{2}}} = \int \dots \frac{dt}{t^2}$ oe such as $\int \dots \frac{dA}{A^{\frac{3}{2}}} = \int \frac{dt}{t^2}$

Condone without the integral signs but the $A^{\frac{3}{2}}$, dA , t^2 and dt must be in the correct positions.

Don't be too concerned on the position of the "5"

M1: Integrates one side correctly. Look for $pA^{-\frac{1}{2}}$ or qt^{-1}

dM1: Integrates both sides correctly. Look for $pA^{-\frac{1}{2}}$ and qt^{-1}

A1: Correct intermediate stage (without $+c$).

Either $-2A^{-\frac{1}{2}} = -\frac{1}{5}t^{-1}(+c)$ or $-10A^{-\frac{1}{2}} = -t^{-1}(+c)$ or equivalent such as $-\frac{A^{-0.5}}{0.5} = -\frac{1}{5}t^{-1}(+c)$

M1: Must have a "+c" now. It is for using $t = 3, A = 2.25 \Rightarrow c = \dots$

There must be some evidence for this award which may be implied by their value for c if no working is seen.

One index must have started out correct

This may be awarded after an incorrect change of subject has occurred.

M1: It is for proceeding to $A = \left(\frac{pt}{qt+r}\right)^2$ from an equation of the form $\frac{p}{\sqrt{A}} = \frac{q}{t} + r$ with correct operations.

Candidates do not need to have found a **value** for r (their constant) to score this mark

The attempt must involve

- using a common factor on the rhs of $\frac{p}{\sqrt{A}} = \frac{q}{t} + r$
- "inverting" both sides not each term
- an attempt to square both sides and not each term to reach $A = \dots$

A1: $A = \left(\frac{30t}{19t+3}\right)^2$ cso. This mark can be awarded independent of the first M1

Be aware that $A = \left(\frac{300t}{190t + 30} \right)^2$ is also correct

(b)

M1: $t \rightarrow \infty, A \rightarrow \left(\frac{a}{b} \right)^2$ The form of part (a) must be $A = \left(\frac{at}{bt + c} \right)^2$ where a, b and c are all positive.

The reason for this is that if we had " $bt - c$ ", the area would be infinite at $t = \frac{c}{b}$ and a limit would not exist

A1: $\frac{900}{361}$ or awrt 2.49 cm² following correct work.

Condone for both marks $A < \frac{900}{361}$ following correct work.

Pearson Education Limited. Registered company number 872828
with its registered office at 80 Strand, London, WC2R 0RL, United Kingdom