

Edexcel Maths S1

Mark Scheme Pack

2005-2013

GCE

Edexcel GCE

Statistics S1 (6683)

Summer 2005

advancing learning, changing lives

Mark Scheme (Results)

**June 2005**  
**6683 Statistics S1**  
**Mark Scheme**

Question Number	Scheme	Marks
1.	<p>Diagram A : <math>y</math> &amp; <math>x</math> : <math>r = -0.79</math>; As <math>x</math> increases, <math>y</math> decreases  <u>or</u> most points lie in the 2<sup>nd</sup> and 4<sup>th</sup> quadrant.</p> <p>Diagram B : <math>v</math> &amp; <math>u</math> : <math>r = 0.08</math>; No real pattern. Several values of <math>v</math> for one value of <math>u</math>  or points lie in all four quadrants, randomly scattered.</p> <p>Diagram C : <math>t</math> &amp; <math>s</math> : <math>r = 0.68</math>; As <math>s</math> increases, <math>t</math> increases or most points lie in the 1<sup>st</sup>  and 3<sup>rd</sup> quadrants</p>	<p>B1;B1dep</p> <p>B1;B1dep</p> <p>B1;B1dep (6)</p>
2. (a)	Distance is a continuous.	continuous B1 (1)
(b)	F.D = freq/class width $\Rightarrow$ 0.8, 3.8, 5.3, 3.7, 0.75, 0.1	or the same multiple of M1 A1 (2)
(c)	$Q_2 = 50.5 + \frac{(67 - 23)}{53} \times 10 = 58.8$ $Q_1 = 52.48; \quad Q_3 = 67.12$	<p>awrt 58.8/58.9 M1 A1</p> <p>awrt 52.5/52.6 67.1/67.3 A1 A1 (4)</p>
	Special case : no working B1 B1 B1 ( $\equiv$ A's on the open)	
(d)	$\bar{x} = \frac{8379.5}{134} = 62.5335\dots$ $s = \sqrt{\frac{557489.75}{134} - \left(\frac{8379.5}{134}\right)^2}$ $s = 15.8089\dots \quad (S_{n-1} = 15.86825\dots)$	<p>awrt 62.5 B1</p> <p>M1 A1√</p> <p>awrt 15.8 (15.9) A1 (4)</p>
	Special case : answer only B1 B1 ( $\equiv$ A's on the open)	
(e)	$\frac{Q_3 - 2Q_2 + Q_1}{Q_3 - Q_1} = \frac{67.12 - 2 \times 58.8 + 52.48}{67.12 - 52.48}$ $= 0.1366 \Rightarrow ; +ve \text{ skew}$	<p>subst their <math>Q_1, Q_2</math> &amp; <math>Q_3</math>  need to show working for A1√  and have reasonable values for  quartiles  awrt 0.14  M1 A1√  A1; B1 (4)</p>
(f)	<p>For +ve skew Mean &gt; Median &amp; <math>62.53 &gt; 58.80</math>  <u>or</u> <math>Q_3 - Q_2</math> (8.32) &gt; <math>Q_2 - Q_1</math> (6.32)  Therefore +ve skew</p>	<p>B1 (1)</p>

Question Number	Scheme	Marks
3. (a)	$S_{xy} = 8880 - \frac{130 \times 48}{8} = (8100)$ <p style="text-align: right;">may be implied</p> $S_{xx} = 20487.5$ $b = \frac{s_{xy}}{s_{xx}} = \frac{8100}{20487.5} = 0.395363\dots$ <p style="text-align: right;">allow use of their <math>S_{xy}</math> for M awrt 0.395</p> $a = \frac{48}{8} - (0.395363\dots) \frac{130}{8} = -0.424649\dots$ <p style="text-align: right;">allow use of their <math>b</math> for M awrt -0.425</p> $y = -0.425 + 0.395x$ <p style="text-align: right;">3s.f.</p> <p>Special case answer only B0 M0 B1 M0 B1 B1 (fully correct 3sf) (<math>\equiv</math> to B0 M0 A1 M0 A1 B1 on the open)</p>	<p>B1</p> <p>M1 A1</p> <p>M1 A1</p> <p>B1 <math>\checkmark</math></p> <p>(6)</p>
(b)	$f - 100 = -0.424649\dots + 0.395\dots(m - 250)$ $f = 0.735 + 0.395m$	<p style="text-align: right;">subst f - 100 &amp; m - 250</p> <p>M1 A1 <math>\checkmark</math></p> <p style="text-align: right;">3 s.f.</p> <p>A1</p> <p>(3)</p>
(c)	$m = 235 \Rightarrow f = 93.64489\dots$	<p style="text-align: right;">awrt 93.6/93.7</p> <p>B1</p> <p>(1)</p>


<p>4(a)</p>	<p> <math>1.5 (Q_3 - Q_1) = 1.5 (28 - 12) = 24</math>  <math>Q_3 + 24 = 52 \Rightarrow 63</math> is an outlier  <math>Q_1 - 24 &lt; 0 \Rightarrow</math> no outliers                 </p>	<p>                     may be implied                      att <math>Q_3 + \dots</math> or <math>Q_1 - \dots</math>,                      52 and -12 or <math>&lt; 0</math> or evidence of no lower outliers                      63 is an outlier                      B1                      M1,                      A1                      A1                      M1                      A1                      A1                      (7)                 </p>
<p>(b)</p>	<p>Distribution is +ve skew; <math>Q_2 - Q_1 (5) &lt; Q_3 - Q_2 (11)</math>;</p>	<p>B1; B1 (2)</p>
<p>(c)</p>	<p>Many delays are small so passengers should find these acceptable or sensible comment in the context of the question.</p>	<p>B1 (1)</p>

<p>5.(a)</p>	$k + 2k + 3k + 5k + 6k = 1$ $17k = 1$ $k = \frac{1}{17} = 0.0588$	<p>use of <math>\sum P(X = x) = 1</math></p>	<p>M1</p> <p>A1</p> <p>(2)</p>
<p>(b)</p>	$E(X) = 1 \times \frac{1}{17} + 2 \times \frac{2}{17} + \dots + 5 \times \frac{6}{17} = \frac{64}{17}$ $= 3 \frac{13}{17}$	<p>use of <math>\sum xP(X = x)</math> and at least 2 prob correct</p> <p>Do not ignore subsequent working</p>	<p>M1</p> <p>A1</p>
<p>(c)</p>	$E(X^2) = 1^2 \times \frac{1}{17} + 2^2 \times \frac{2}{17} + \dots + 5^2 \times \frac{6}{17} = \left( \frac{266}{17} = 15.6 \right)$ $\text{Var}(X) = \frac{266}{17} - \left( \frac{64}{17} \right)^2$ $(E(X))^2$ $= 1.4740\dots$	<p>use of <math>\sum x^2 P(X = x)</math></p> <p>and at least 2 prob correct</p> <p>use of <math>\sum x^2 P(X = x) -</math></p> <p>awrt 1.47</p>	<p>M1 A1</p> <p>A1</p> <p>(4)</p>
<p>(d)</p>	$\text{Var}(4 - 3X) = 9 \text{Var}(X) = 9 \times 1.47 = 13.23 \Rightarrow 13.2$ $\text{or } 9 \times 1.4740\dots = 13.266 \Rightarrow 13.3$	<p>cao <math>9 \text{Var} X</math></p>	<p>M1 A1</p> <p>(2)</p>

<p>6(a)</p> <p>(b)</p> <p>(c)</p>	<p><math>M \sim N(155, 3.5^2)</math></p> <p><math>P(M &gt; 160) = P\left(z &gt; \frac{160-155}{3.5}\right)</math>  <math>= P(z &gt; 1.43)</math>  <math>= 0.0764</math></p> <p><math>P(150 \leq M \leq 157) = P(-1.43 \leq z \leq 0.57)</math>  <math>= 0.7157 - (1 - 0.9236)</math>  <math>= 0.6393</math></p> <p>special case : answer only B0 B0 M1 A1</p> <p><math>P(M \leq m) = 0.3 \Rightarrow \frac{m-155}{3.5} = -0.5244</math>  <math>m = 153.2</math></p>	<p>standardising  <math>\pm(160-155), \sigma, \sigma^2, \sqrt{\sigma}</math></p> <p>M1 A1 A1 (3)</p> <p>awrt -1.43, 0.57  <math>p &gt; 0.5</math>          0.6393 - 0.6400 4dp</p> <p>B1 B1 M1 A1 (4)</p> <p>-0.5244          att stand = z value          for A1 may use awrt to -          0.52.</p> <p>B1 M1 A1 A1 (4)</p>																									
<p>7.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th></th> <th style="text-align: center;">Glasses</th> <th style="text-align: center;">No Glasses</th> <th style="text-align: center;">Totals</th> <th></th> </tr> </thead> <tbody> <tr> <td>Science</td> <td style="text-align: center;">18</td> <td style="text-align: center;"><b>12</b></td> <td style="text-align: center;">30</td> <td></td> </tr> <tr> <td>Arts</td> <td style="text-align: center;"><b>27</b></td> <td style="text-align: center;"><b>23</b></td> <td style="text-align: center;"><b>50</b></td> <td>50 may be seen in (a)</td> </tr> <tr> <td>Humanities</td> <td style="text-align: center;">44</td> <td style="text-align: center;"><b>24</b></td> <td style="text-align: center;">68</td> <td>23 may be seen in (b)</td> </tr> <tr> <td>Totals</td> <td style="text-align: center;">89</td> <td style="text-align: center;"><b>59</b></td> <td style="text-align: center;">148</td> <td></td> </tr> </tbody> </table> <p><math>P(\text{Arts}) = \frac{50}{148} = \frac{25}{74} = 0.338</math></p> <p><math>P(\text{No glasses} / \text{Arts}) = \frac{23/148}{50/148} = \frac{23}{50} = 0.46</math></p> <p><math>P(\text{Right Handed}) = \left(\frac{30}{148} \times 0.8\right) + \left(\frac{50}{148} \times 0.7\right) + \left(\frac{68}{148} \times 0.75\right)</math>  <math>= \frac{55}{74} = 0.743</math></p> <p><math>P(\text{Science} / \text{Right handed}) = \frac{\frac{30}{148} \times 0.8}{\frac{55}{74}} = \frac{12}{55} = 0.218</math></p>		Glasses	No Glasses	Totals		Science	18	<b>12</b>	30		Arts	<b>27</b>	<b>23</b>	<b>50</b>	50 may be seen in (a)	Humanities	44	<b>24</b>	68	23 may be seen in (b)	Totals	89	<b>59</b>	148		<p>B1 B1</p> <p>a number/148</p> <p>M1 A1 (4)</p> <p>prob their(a)prob or number their 50</p> <p>M1 A1 (2)</p> <p>attempt add three prob A1 ✓ on their (a)</p> <p>awrt 0.743</p> <p>M1 A1 ✓ A1 (3)</p> <p>✓ on their (c)</p> <p>M1 A1 ✓ A1 (3)</p>
	Glasses	No Glasses	Totals																								
Science	18	<b>12</b>	30																								
Arts	<b>27</b>	<b>23</b>	<b>50</b>	50 may be seen in (a)																							
Humanities	44	<b>24</b>	68	23 may be seen in (b)																							
Totals	89	<b>59</b>	148																								



Question Number	Scheme	Marks
1. (a)	Mode is 56	B1 (1)
(b)	$Q_1 = 35, Q_2 = 52, Q_3 = 60$	B1,B1,B1 (3)
(c)	$\bar{x} = \frac{1335}{27} = 49.\dot{4}$ or $49\frac{4}{9}$	exact or awrt 49.4 B1
	$\sigma^2 = \frac{71801}{27} - \left(\frac{1335}{27}\right)^2 = 214.5432\dots$ $\sigma = 14.6$ or $14.9$	M1A1ft awrt 14.6(5) or 14.9 A1 (4)
(d)	$\frac{49.4-56}{14.6} = -0.448$	awrt range -0.44 to -0.46 M1A1 (2)
(e)	For negative skew; Mean < median < mode (49.4 < 52 < 56 not required) $Q_3 - Q_2 < Q_2 - Q_1$ 8 and 17 Accept other valid reason eg. 3(mean - median)/sd as alt for M1A1	2 compared correctly 3 compared correctly M1 A1 M1 A1 ft (4)
<b>Total 14 marks</b>		
2. (a)	$p + q = 0.4$ $2p + 4q = 1.3$	B1 M1A1 Consider with (b). (3)
(b)	Attempt to solve $p = 0.15, q = 0.25$	M1 If both seen, award 3. A1A1 (3)
(c)	$E(X^2) = 1^2 \times 0.10 + 2^2 \times 0.15 + \dots + 5^2 \times 0.30 = 14$ $\text{Var}(X) = 14 - 3.5^2 = 1.75$	M1A1ft M1A1 (4)
(d)	$\text{Var}(3 - 2X) = 4\text{Var}(X) = 7.00$	M1A1ft (2) <b>Total 12 marks</b>

<p>3. (a)</p>	<p>Sensible graph scales, labels, shape</p> 	<p>B1,B1,B1</p>
<p>(b)</p>	<p>Points lie close to a straight line</p>	<p>B1 (3)</p>
<p>(c)</p>	$S_{xy} = 8354 - \frac{106 \times 704}{10} = 891.6$ $S_{xx} = 1352 - \frac{106^2}{10} = 228.4$ $b = \frac{891.6}{228.4} = 3.903677\dots$ $a = \frac{704}{10} - b \frac{106}{10} = 29.021015\dots$	<p>B1 (1)</p> <p>B1</p> <p>B1</p> <p>M1A1 awrt 3.9</p> <p>M1A1 awrt 29</p> <p>29.02, 3.90 A1ft (7)</p>
<p>(d)</p>	<p>For every extra week in storage, another 3.90 ml of chemical evaporates</p>	<p>B1 (1)</p>
<p>(e)</p>	<p>(i) 103.12      (ii) 165.52</p>	<p>B1B1 (2)</p>
<p>(f)</p>	<p>(i) Close to range of <math>x</math>, so reasonably reliable                  (ii) Well outside range of <math>x</math>, could be unreliable since no evidence that model will continue to hold</p>	<p>B1,B1 B1 (4)</p> <p>B1</p> <p><b>Total 18 marks</b></p>

4. (a)	<p style="text-align: right;">Tree</p> <p style="text-align: center;"><math>\frac{9}{12}, \frac{3}{12}</math></p> <p style="text-align: right;">Complete &amp; labels</p>	<p>M1</p> <p>A1</p> <p>A1 (3)</p>
(b)	$P(\text{Second ball is red}) = \frac{9}{12} \times \frac{3}{11} + \frac{3}{12} \times \frac{2}{11} = \frac{1}{4}$	<p>M1A1 (2)</p>
(c)	$P(\text{Both are red} \mid \text{Second ball is red}) = \frac{\frac{3}{12} \times \frac{2}{11}}{\frac{1}{4}} = \frac{2}{11}$ <p style="text-align: right;">exact or awrt 0.182</p>	<p>M1A 1 (2)</p> <p><b>Total 7 marks</b></p>
5. (a)	<p>To simplify a real world problem                  To improve understanding / describe / analyse a real world problem                  Quicker and cheaper than using real thing                  To predict possible future outcomes                  Refine model / change parameters possible</p> <p style="text-align: right;">Any 2</p>	<p>B1B1 (2)</p>
(b)	<p>(i) e.g.s height, weight                      (ii) score on a face after tossing a fair die</p>	<p>B1B1 (2)</p> <p><b>Total 4 marks</b></p>

6. (a)		$\mathcal{E}$	Venn Diagram 0.32, 0.11 & A, B 0.22, 0.35 & box	M1 A1 A1 (3)
(b)	$P(A) = 0.32 + 0.22 = 0.54; P(B) = 0.33$		M1A1ft; A1ft (3)	
(c)	$P(A B') = \frac{P(A \cap B')}{P(B')} = \frac{32}{67}$	awrt 0.478	M1A1 (2)	
(d)	For independence $P(A \cap B) = P(A)P(B)$ For these data $0.22 \neq 0.54 \times 0.33 = 0.1782$ (OR $P(A B') \neq P(A)$ for M1A1ft OR $\frac{2}{3} = P(A B) \neq P(A) = 0.54$ for M1A1ft) $\therefore$ NOT independent		M1A1ft  A1ft (3) <b>Total 11 marks</b>	
7. (a)	Let $H$ be rv height of athletes, so $H \sim N(180, 5.2^2)$ $P(H > 188) = P(Z > \frac{188 - 180}{5.2}) = P(Z > 1.54) = 0.0618$ $\pm$ stand. $\sqrt{\cdot}$ , sq, awrt 0.062		M1A1A1 (3)	
(b)	Let $W$ be rv weight of athletes, so $W \sim N(85, 7.1^2)$ $P(W < 97) = P(Z < 1.69) = 0.9545$ standardise, awrt 0.9545		M1A1 (2)	
(c)	$P(H > 188 \text{ \& } W < 97) = 0.0618(1 - 0.9545)$ $= 0.00281$ allow (a)x(b) for M awrt 0.0028		M1A1ft A1 (3)	
(d)	Evidence suggests height and weight are positively correlated / linked Assumption of independence is not sensible		B1 (1) <b>Total 9 marks</b>	

GCE

Edexcel GCE


Statistics S1 (6683)

June 2006

advancing learning, changing lives

Mark Scheme  
(Results)


**June 2006**  
**6683 Statistics S1**  
**Mark Scheme**

Question Number	Scheme	Marks
1(a)	Indicates max / median / min / upper quartile/ lower quartile (2 or more) Indicates outliers (or equivalent description) Illustrates skewness (or equivalent description e.g. shape)      Any 3 rows Allows comparisons Indicates range / IQR / spread	B1 B1 B1 (3)
(b)(i) (ii)	37 (minutes) Upper quartile or $Q_3$ or third quartile or 75 <sup>th</sup> percentile or $P_{75}$	B1 B1 (2)
(c)	Outlier s How to calculate correctly ‘Observations that are very different from the other observations and need to be treated with caution’ These two children probably walked / took a lot longer	Any 2 B1 B1 (2)
(d)	 <p style="text-align: right;">Box &amp; median &amp; whiskers Sensible scale 30, 37, 50 25, 55</p>	M1 B1 B1 B1 (4)
(e)	Children from school A generally took less time      Any correct 4 lines 50% of B $\leq$ 37 mins, 75% of A < 37 mins (similarly for 30) Median/Q1/Q3 of A < median/Q1/Q3 of B (1 or more) A has outliers, (B does not) Both positive skew IQR of A < IQR of B, range of A > range of B	B1 B1 B1 B1 (4) <b>Total 15</b>

Question Number	Scheme	Marks
2. (a)	$P(\text{both longer than } 24.5) = \frac{11}{55} \times \frac{10}{54} = \frac{1}{27} \text{ or } 0.\dot{0}3\dot{7} \text{ or } 0.037$ <p style="text-align: right;">2 fracs x w/o rep. awrt 0.037</p>	<p><b>M1A1</b></p> <p style="text-align: right;">(2)</p>
(b)	<p>Estimate of mean time spent on their conversations is</p> $\bar{x} = \frac{1060}{55} = 19\frac{3}{11} \text{ or } 19.\dot{2}\dot{7} \text{ or } 19.3$ <p style="text-align: right;">1060/total, awrt 19.3 or 19mins 16s</p>	<p><b>M1A1</b></p> <p style="text-align: right;">(2)</p>
(c)	$\frac{1060 + \sum fy}{80} = 21$ <p style="text-align: right;">21x80=1680</p> $\sum fy = 620$ <p style="text-align: right;">Subtracting 'their 1060'</p> $\therefore \bar{y} = \frac{620}{25} = 24.8$ <p style="text-align: right;">Dividing their 620 by 25</p>	<p><b>B1</b></p> <p><b>M1</b></p> <p><b>M1A1</b></p> <p style="text-align: right;">(4)</p>
(d)	<p>Increase in mean value. Length of conversations increased considerably during 25 weeks relative to 55 weeks</p> <p style="text-align: right;">context - ft only from <b>comment</b> above</p>	<p><b>B1</b></p> <p><b>B1</b> ∫</p> <p style="text-align: right;">(2)</p> <p style="text-align: right;"><b>Total 10</b></p>
3. (a)	$\sum x = \sum t = 337.1, \quad \sum y = 16.28$ <p style="text-align: right;">Can be implied</p> $S_{xy} = 757.467 - \frac{337.1 \times 16.28}{8} = 71.4685$ <p style="text-align: right;">either method, awrt 71.5</p> $S_{xx} = 15965.01 - \frac{337.1^2}{8} = 1760.45875$ <p style="text-align: right;">awrt 1760</p>	<p><b>B1, B1</b></p> <p><b>M1A1</b></p> <p><b>A1</b></p> <p style="text-align: right;">(5)</p>
(b)	$b = \frac{71.4685}{1760.45875} = 0.04059652$ <p style="text-align: right;">/ correct way up, awrt 0.0406</p> $a = \frac{16.28}{8} - b \times \frac{337.1}{8} = 0.324364$ <p style="text-align: right;">using correct formula, awrt 0.324</p> $y = 0.324 + 0.0406x$ <p style="text-align: right;">3 sf or better but award for copying from above</p>	<p><b>M1A1</b></p> <p><b>A1</b> ∫</p> <p style="text-align: right;">(5)</p>
(c)	<p>At <math>t = 40</math>, <math>x = 40</math>, <math>y = 1.948</math>, <math>l = 2461.948</math></p> <p style="text-align: right;">sub <math>x=40</math>, awrt 1.95, awrt 2461.95</p>	<p><b>M1A1A1</b> ∫</p> <p style="text-align: right;">(3)</p>
(d)	$l - 2460 = 0.324 + 0.0406t$ $l = 2460.324 + 0.0406t$ <p style="text-align: right;">LHS required awrt 2460.32, f.t. their 0.0406, / and t</p>	<p><b>M1</b></p> <p style="text-align: right;">(2)</p>
(e)	<p>At <math>t = 90</math>, <math>l = 2463.978</math></p> <p style="text-align: right;">awrt 2464</p>	<p><b>B1</b></p> <p style="text-align: right;">(1)</p>
(f)	<p>90°C outside range of data unlikely to be reliable</p>	<p><b>B1</b></p>

<p>4 (a)</p> <p>(b)</p> <p>M1A1 ∫</p> <p>(c)</p>	<p><math>E(X) = 3;</math></p> <p><math>Var(X) = \frac{25-1}{12} = 2</math> **AG**</p> <p><math>Var(X) = 1^2 \times \frac{1}{5} + 2^2 \times \frac{1}{5} + 3^2 \times \frac{1}{5} \dots - 3^2 = 11 - 9 = 2</math> **AG**</p> <p>Accept (55/5)-9 as minimum evidence.</p> <p><math>E(3X - 2) = 3E(X) - 2 = 7</math></p> <p><math>Var(4 - 3x) = 3^2 Var(X) = 18</math></p>	<p>B1</p> <p>M1A1</p> <p>(3)</p> <p>(2)</p> <p>M1A1</p> <p>(2)</p> <p>Total 7</p>
<p>5(a)</p> <p>(b)</p> <p>(c)</p>	<div data-bbox="411 808 1169 1218" data-label="Figure"> </div> <p>2 separate sketches OK.</p> <p>Bell Shape 1.78 &amp; 0.2 1.65 &amp; 0.3</p> <p>Accept clear alternatives to 0.3: 0.7/0.5/0.2</p> <p><math>\frac{1.78 - \mu}{\sigma} = 0.8416 \Rightarrow 1.78 - \mu = 0.8416\sigma</math> either for method 0.8416</p> <p><math>\frac{1.65 - \mu}{\sigma} = -0.5244 \Rightarrow 1.65 - \mu = -0.5244\sigma</math> (-)0.5244</p> <p>Solving gives <math>\mu = 1.70, \sigma = 0.095</math> N.B. awrt 0.84, 0.52 B1B0 awrt 1.7, 0.095 cao</p> <p><math>P(\text{height} \geq 1.74) = 1 - P(\text{height} &lt; 1.74)</math> 'one minus'</p> <p><math>= 1 - P\left(Z &lt; \frac{1.74 - 1.70}{0.095}\right)</math> standardise with their mu and sigma</p> <p><math>= 1 - P(Z &lt; 0.42) = 0.3372</math> awrt 0.337</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>(3)</p> <p>M1</p> <p>B1</p> <p>B1</p> <p>M1A1A1</p> <p>(6)</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>(3)</p> <p>Total 12</p>

6.(a)



3 closed curves that intersect  
 Subtract at either stage  
 9,7,5  
 10,6,12  
 41 & box

**M1**  
**M1**  
**A1**  
**A1**  
**A1**  
**(6)**

(b)  $P(G, \overline{LH}, \overline{D}) = \frac{10}{100} = \frac{1}{10}$

**B1** ∫  
**(1)**

(c)  $P(\overline{G}, \overline{LH}, \overline{D}) = \frac{41}{100}$

**B1** ∫  
**(1)**

(d)  $P(\text{Only two attributes}) = \frac{9+7+5}{100} = \frac{21}{100}$

**M1A1** ∫  
**(2)**

(e)  $P(G | LH \& DH) = \frac{P(G \& LH \& DH)}{P(LH \& DH)} = \frac{\frac{10}{100}}{\frac{15}{100}} = \frac{10}{15} = \frac{2}{3}$  awrt 0.667

**M1A1** ∫ **A1**

N.B. Assumption of independence M0

**(3)**  
**Total 13**

# Mark Scheme (Results)

## January 2007

advancing learning, changing lives


GCE

GCE Mathematics

Statistics (6683)

**January 2007  
6683 Statistics S1  
Mark Scheme**

Question number	Scheme	Marks
1. (a)	(£) 17	Just <b>17</b>
(b)	$\sum t = 212$ and $\sum m = 61$ (Accept as totals under each column in qu.)	B1, B1
	$S_m = 2485 - \frac{61 \times 212}{10}, = 1191.8$	awrt <b>1190</b> or 119 (3sf)
	$S_n = 983.6$ (awrt <b>984</b> ) and $S_{mm} = 1728.9$ (awrt <b>1730</b> )	(or 98.4 and 173)
(c)	$r = \frac{1191.8}{\sqrt{983.6 \times 1728.9}}$ $= 0.913922\dots$	M1, A1 A1, A1 (6) M1, A1f.t. awrt <b>0.914</b>
(d)	0.914 (Must be the same as (c) or awrt 0.914)	B1f.t. ( $ r  < 1$ )
	e.g. linear transformation, coding does not affect coefficient (or recalculate)	dB1 (2)
(e)	0.914 suggests longer spent shopping the more spent. (Idea more time, more spent)	B1
	0.178 different amounts spent for same time.	B1 (2)
(f)	e.g. might spend short time buying 1 expensive item <u>OR</u> might spend a long time checking for bargains, talking, buying lots of cheap items.	B1g (1)
<b>15 marks</b>		
(b)	M1 for one correct formula seen, f.t. their $\sum t, \sum m$ [Use 1 <sup>st</sup> A1 for 1 correct, 2 <sup>nd</sup> A1 for 2 etc]	
(c)	M1 for attempt at correct formula, $\frac{2485}{\sqrt{2101 \times 5478}}$ scores M1A0A0 A1ft f.t. their values for $S_n$ etc from (b) but don't give for $S_n = 5478$ etc (see above) Answer only (awrt 0.914) scores 3/3, 0.913 (i.e. truncation) can score M1A1ft by implication.	
(d)	2 <sup>nd</sup> B1 dependent on 1 <sup>st</sup> B1 Accept $\sum m = 261, \sum m^2 = 8541, \sum tm = 6725 \rightarrow 0.914$	
(e)	One mark for a sensible comment relating to each coefficient For 0.178 allow "little or no link between time and amount spent". Must be in context. Just saying 0.914 is strong +ve correlation between amount spent and time shopping and 0.178 is weak correlation ...scores B0B0.	
(f)	B1g for a sensible, practical suggestion showing that other factors might affect the amount spent. E.g. different day (weekend vs weekday) or time of day (time spent queuing if busy)	

Question number	Scheme	Marks
2. (a)	 <p style="text-align: right;">Correct tree shape</p> <p style="text-align: right;">A, B and C and 0.35 and 0.25</p> <p style="text-align: right;">D (x3) and 0.03, 0.06, 0.05 (May be implied by seeing P(A ∩ D) etc at the ends)</p>	<p>M1</p> <p>A1</p> <p>A1 (3)</p>
(b)(i)	$P(A \cap D) = 0.35 \times 0.03, = \underline{\mathbf{0.0105}}$ or $\frac{21}{2000}$	<p>M1, A1</p>
(ii)	$P(D) = (i) + 0.25 \times 0.06 + (0.4 \times 0.05)$ $= \underline{\mathbf{0.0455}}$ or $\frac{91}{2000}$	<p>P(C) = 0.4 (anywhere) B1</p> <p>M1</p> <p>A1 (5)</p>
(c)	$P(C D) = \frac{P(C \cap D)}{P(D)}, = \frac{0.4 \times 0.05}{(ii)}$ $= 0.43956... \text{ or } \frac{40}{91}$ <p style="text-align: right;"><b>0.44</b> or awrt <b>0.440</b></p> <p>[Correct answers only score full marks in each part]</p>	<p>M1, A1ft</p> <p>A1 (3)</p> <p style="text-align: right;"><b>11 marks</b></p>
(a)	<p>M1 for tree diagram, 3 branches and then two from each. At least one probability attempted.</p> <p>(b) 1<sup>st</sup> M1 for 0.35x0.03. Allow for equivalent from <u>their</u> tree diagram.</p> <p>B1 for P(C) = 0.4, can be in correct place on tree diagram or implied by 0.4x0.05 in P(D).</p> <p>2<sup>nd</sup> M1 for all 3 cases attempted and <u>some</u> correct probabilities seen, including +. Can ft their tree. Condone poor use of notation if correct calculations seen. E.g. P(C   D) for P(C ∩ D) .</p> <p>(c) M1 for attempting correct ratio of probabilities. There must be an attempt to substitute some values in a correct formula. If no correct formula and ration not correct ft score M0. Writing P(D C) and attempting to find this is M0. Writing P(D C) but calculating correct ratio – ignore notation and mark ratios.</p> <p>A1ft must have their 0.4 x0.05 divided by their (ii). If ratio is incorrect ft (0/3) unless correct formula seen and part of ratio is correct then M1.</p>	

Question number	Scheme	Marks														
3. (a)	<p>N.B. Part (a) doesn't have to be in a table, could be a list <math>P(X = 1) = \dots</math> etc</p> <table border="1" data-bbox="293 349 968 510"> <tr> <td><math>x</math></td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> </tr> <tr> <td><math>P(X = x)</math></td> <td><math>\frac{1}{36}</math></td> <td><math>\frac{3}{36}</math></td> <td><math>\frac{5}{36}</math></td> <td><math>\frac{7}{36}</math></td> <td><math>\frac{9}{36}</math></td> <td><math>\frac{11}{36}</math></td> </tr> </table> <p>0.0278, 0.0833, 0.139, 0.194, 0.25, 0.306 (Accept awrt 3 s.f)</p> <p>(b) <math>P(3) + P(4) + P(5) = \frac{21}{36}</math> or <math>\frac{7}{12}</math> or awrt 0.583</p> <p>(c) <math>E(X) = \frac{1}{36} + 2 \times \frac{3}{36} + \dots = \frac{161}{36}</math> or 4.472 or <math>4\frac{17}{36}</math></p> <p>(d) <math>E(X^2) = \frac{1}{36} + 2^2 \times \frac{3}{36} + \dots = \frac{791}{36}</math> or full expression or <math>21\frac{35}{36}</math> or awrt 21.97</p> <p><math>\text{Var}(X) = \frac{791}{36} - \left(\frac{161}{36}\right)^2</math>, = <b>1.9714... *</b></p> <p>(e) <math>\text{Var}(2 - 3X) = 9 \times 1.97</math> or <math>(-3)^2 \times 1.97</math>, = 17.73 awrt <b>17.7</b> or <math>\frac{2555}{144}</math></p>	$x$	1	2	3	4	5	6	$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$	<p>B1, B1, B1</p> <p>(3)</p> <p>M1, A1 (2)</p> <p>M1, A1 (2)</p> <p>M1, A1</p> <p>M1, A1c.s.o. (4)</p> <p>M1, A1 (2)</p> <p><b>13 marks</b></p>
$x$	1	2	3	4	5	6										
$P(X = x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{5}{36}$	$\frac{7}{36}$	$\frac{9}{36}$	$\frac{11}{36}$										
	<p>(a) 1<sup>st</sup> B1 for <math>x = 1, \dots, 6</math> and at least one correct probability N.B. <math>\frac{3}{36} = \frac{1}{12}</math> and <math>\frac{9}{36} = \frac{1}{4}</math>                  2<sup>nd</sup> B1 for at least 3 correct probabilities                  3<sup>rd</sup> B1 for a fully correct probability distribution.</p> <p>(b) M1 for attempt to add the correct three probabilities, ft their probability distribution</p> <p>(c) M1 for a correct attempt at <math>E(X)</math>. Minimum is as printed. Exact answer only scores M1A1.                  [Division by 6 at any point scores M0, no ISW. Non-exact answers with no working score M0.]</p> <p>(d) 1<sup>st</sup> M1 for a correct attempt at <math>E(X^2)</math>. Minimum as printed. <math>\frac{791}{36}</math> or awrt 21.97 scores M1A1.                  2<sup>nd</sup> M1 for their <math>E(X^2) - (\text{their } E(X))^2</math>.                  2<sup>nd</sup> A1 cso needs awrt 1.97 <u>and</u> <math>\frac{791}{36} - \left(\frac{161}{36}\right)^2</math> or <math>\frac{2555}{1296}</math> or any fully correct expression seen.                  Can accept <u>at least 4 sf</u> for both. i.e. 21.97 for <math>\frac{791}{36}</math>, 4.472 for <math>\frac{161}{36}</math>, 20.00 for <math>\left(\frac{161}{36}\right)^2</math>.</p> <p>(e) M1 for correct use of <math>\text{Var}(aX + b)</math> formula or a <u>full</u> method.                  NB <math>-3^2 \times 1.97</math> followed by awrt 17.7 scores M1A1 <u>BUT</u> <math>-3^2 \times 1.97</math> alone, or followed by <math>-17.7</math>, scores M0A0.</p>															

Question number	Scheme	Marks
4. (a)	Positive skew (both bits)	B1 (1)
(b)	$19.5 + \frac{(60-29)}{43} \times 10, = 26.7093\dots$ (N.B. Use of 60.5 gives 26.825... so allow awrt 26.8)	awrt <b>26.7</b> M1, A1 (2)
(c)	$\mu = \frac{3550}{120} = 29.5833\dots$ or $29\frac{7}{12}$ $\sigma^2 = \frac{138020}{120} - \mu^2$ or $\sigma = \sqrt{\frac{138020}{120} - \mu^2}$ $\sigma = 16.5829\dots$ or ( $s = 16.652\dots$ )	awrt <b>29.6</b> B1 M1 awrt <b>16.6</b> (or $s = 16.7$ ) A1 (3)
(d)	$\frac{3(29.6 - 26.7)}{16.6}$ $= 0.52\dots$ (N.B. 60.5 in (b) ...awrt 0.499[or with $s$ awrt 0.497])	M1A1ft awrt <b>0.520</b> (or with $s$ awrt 0.518) A1 (3)
(e)	$0.520 > 0$ So it is consistent with (a)	correct statement about their (d) being $>0$ or $<0$ ft their (d) B1ft dB1ft (2)
(f)	Use <u>Median</u> Since the data is skewed <u>or</u> less affected by outliers/extreme values	B1 dB1 (2)
(g)	If the data are <u>symmetrical</u> or <u>skewness is zero</u> or <u>normal/uniform distribution</u> (“mean =median” or “no outliers” or “evenly distributed” all score B0)	B1 (1) <b>14 marks</b>
(b)	M1 for $(19.5 \text{ or } 20) + \frac{(60-29)}{43} \times 10$ or better. Allow 60.5 giving awrt 26.8 for M1A1 Allow their $0.5n$ [or $0.5(n+1)$ ] instead of 60 [or 60.5] for M1.	
(c)	M1 for a correct expression for $\sigma, \sigma^2, s$ or $s^2$ . NB $\sigma^2 = 274.99$ and $s^2 = 277.30$ Condone poor notation if answer is awrt 16.6 (or 16.7 for $s$ )	
(d)	M1 for attempt to use this formula using their values to any accuracy. Condone missing 3. 1 <sup>st</sup> A1ft for using their values to at least 3sf. Must have the 3. 2 <sup>nd</sup> A1 for using accurate enough values to get awrt 0.520 (or 0.518 if using $s$ ) NB Using only 3 sf gives 0.524 and scores M1A1A0	
(e)	1 <sup>st</sup> B1 for saying or implying correct sign for their (d). B1g and B1ft. Ignore “correlation” if seen. 2 <sup>nd</sup> B1 for a comment about consistency with their (d) and (a) being positive skew, ft their (d) only This is dependent on 1 <sup>st</sup> B1: so if (d) $>0$ , they say yes, if (d) $<0$ they say no.	
(f)	2 <sup>nd</sup> B1 is dependent upon choosing median.	

Question number	Scheme	Marks
5. (a)	Time is a <u>continuous</u> variable <u>or</u> data is in a <u>grouped</u> frequency table	B1 (1)
(b)	Area is proportional to frequency <u>or</u> $A \propto f$ <u>or</u> $A = kf$	B1 (1)
(c)	$3.6 \times 2 = 0.8 \times 9$ <p>1 child represented by 0.8</p>	M1 dM1 A1 cso (3)
(d)	$(\text{Total}) = \frac{24}{0.8}, = \underline{\underline{30}}$	M1, A1 (2)
<b>7 marks</b>		
(b)	<p>1<sup>st</sup> B1 for one of these correct statements.                      “Area proportional to frequency density” or “Area = frequency” is B0</p>	
(c)	<p>1<sup>st</sup> M1 for a correct combination of any 2 of the 4 numbers: 3.6, 2, 0.8 and 9                      e.g. <math>3.6 \times 2</math> or <math>\frac{3.6}{0.8}</math> or <math>\frac{0.8}{2}</math> etc BUT e.g. <math>\frac{3.6}{2}</math> is M0                      2<sup>nd</sup> M1 dependent on 1<sup>st</sup> M1 and for a correct combination of 3 numbers leading to 4<sup>th</sup>.                      May be in separate stages but must see all 4 numbers                      A1cso for fully correct solution. Both Ms scored, no false working seen and <u>comment required</u>.</p>	
(d)	M1 for $\frac{24}{0.8}$ seen or implied.	

Question number	Scheme	Marks
6. (a)	Used to simplify <u>or</u> represent a real world problem Cheaper <u>or</u> quicker <u>or</u> easier (than the real situation) <u>or</u> more easily modified To improve understanding of the real world problem Used to predict outcomes from a real world problem (idea of predictions)	(any two lines) B1 B1 (2)
(b)	(3 or 4) Model used to make predictions. (Idea of predicted values based on the model)	B1
	(4 or 3) (Experimental) data collected	B1
	(7) Model is refined.	B1 (3)
<b>5 marks</b>		
(a)	1 <sup>st</sup> B1 For one line 2 <sup>nd</sup> B1 For a second line Be generous for 1 <sup>st</sup> B1 but stricter for B1B1	
(b)	1 <sup>st</sup> & 2 <sup>nd</sup> B1 These two points can be interchanged. Idea of values from (experimental) data and predicted values based on the model.	
	1 <sup>st</sup> B1 for predicted values from model e.g. “model used to gain suitable data”	
	2 <sup>nd</sup> B1 for data collected. Idea of experimental data but “experiment” needn’t be explicitly seen	
	3 <sup>rd</sup> B1 This should be stage 7. Idea of refinement or revision or adjustment	

Question number	Scheme	Marks
7. (a)	$P(X < 91) = P\left(Z < \frac{91-100}{15}\right)$ $= P(Z < -0.6)$ $= 1 - 0.7257$ $= 0.2743$	Attempt standardisation M1 A1 M1 awrt <b>0.274</b> A1 (4)
(b)	$1 - 0.2090 = 0.7910$ $P(X > 100+k) = 0.2090 \quad \text{or} \quad P(X < 100+k) = 0.7910 \quad (\text{May be implied})$ $\frac{100+k-100}{15} = 0.81 \quad (\text{ft their } z = 0.81, \text{ but must be } z \text{ not prob.})$ $\underline{k = 12}$	0.791 B1 M1 B1 M1, A1ft A1 cao (6) <b>10 marks</b>
(a)	$\pm \frac{(91-\mu)}{\sigma \text{ or } \sigma^2}$ 1 <sup>st</sup> M1 for attempting standardisation. Can use of 109 instead of 91. Use of 90.5 etc is M0 1 <sup>st</sup> A1 for -0.6 (or +0.6 if using 109) 2 <sup>nd</sup> M1 for 1 - probability from tables. Probability should be > 0.5	
(b)	1 <sup>st</sup> B1 for 0.791 seen or implied. 1 <sup>st</sup> M1 for a correct probability statement, but must use X or Z correctly. Shown on diagram is OK 2 <sup>nd</sup> B1 for awrt 0.81 seen (or implied by correct answer - see below) (Calculator gives 0.80989...) 2 <sup>nd</sup> M1 for attempting to standardise e.g. $\frac{100+k-100}{15}$ or $\frac{k}{15}$ $\frac{X-100}{15}$ scores 2 <sup>nd</sup> M0 until the 100+ k is substituted to give k, but may imply 1 <sup>st</sup> M1 if k= 112.15 seen 1 <sup>st</sup> A1ft for correct equation for k (as written or better). Can be implied by k = 12.15 (or better) 2 <sup>nd</sup> A1 for k = 12 only. <u>Answers only</u> k = 112 or 112.15 or better scores 3/6 (on EPEN give first 3 marks) k = 12.15 or better (calculator gives 12.148438...) scores 5/6 (i.e loses last A1 only) k = 12 (no incorrect working seen) scores 6/6 NB Using 0.7910 instead of 0.81 gives 11.865 which might be rounded to 12. This should score no more than B1M1B0M1A0A0.	

# Mark Scheme (Results)

## Summer 2007

GCE

GCE Mathematics


Statistics S1 (6681)

June 2007  
6683 Statistics S1  
Mark Scheme

Question Number	Scheme	Marks
1. (a)	$r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} = \frac{-808.917}{\sqrt{113573 \times 8.657}}$ $= -0.81579\dots$	<p style="text-align: center;"><b>M1</b></p> <p style="text-align: center;"><b>A1</b> (2)</p>
(b)	Houses are <u>cheaper</u> further away from the station or equivalent statement	<b>B1</b>
(c)	-0.816	<b>B1</b> (1)
<b>Total 4 marks</b>		
Notes:		
1(a)	<p>M1 for knowing formula and clear attempt to sub in correct values from question. Root required for method. Anything that rounds to -0.82 for A1. Correct answer with no working award 2/2</p>	
(b)	<p>Context based on negative correlation only required. Accept <u>Houses</u> are <u>more expensive</u> closer to the <u>station</u> or equivalent statement. Require 'house prices' or 'station' and a clear correct comparison.</p>	
(c)	<p>Accept anything that rounds to -0.82 or 'the same' or 'unchanged' or equivalent. Award B1 if value quoted same as answer to (a).</p>	

Question Number	Scheme	Marks
<p>2(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p><math>\frac{1}{2}</math></p> <p>54</p> <p>+ is an 'outlier' or 'extreme value' Any heavy musical instrument or a statement that the instrument is heavy</p> <p><math>Q_3 - Q_2 = Q_2 - Q_1</math> so symmetrical or no skew</p> <p><math>P(W &lt; 54) = 0.75</math> (or <math>P(W &gt; 54) = 0.25</math>) or correctly labelled and shaded diagram</p> <p><math>\frac{54 - 45}{\sigma} = 0.67</math> <math>\sigma = 13.43.....</math></p>	<p><b>B1</b> <b>(1)</b></p> <p><b>B1</b> <b>(1)</b></p> <p><b>B1</b> <b>(2)</b></p> <p><b>B1</b> <b>(2)</b></p> <p><b>M1</b> <b>(2)</b></p> <p><b>M1B1</b> <b>(4)</b></p> <p><b>A1</b> <b>(4)</b></p> <p><b>Total 10 marks</b></p>
<p>Notes</p> <p>2(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>Accept 50% or half or 0.5. Units not required.</p> <p>Correct answer only. Units not required.</p> <p>'Anomaly' only award B0 Accept '85kg was heaviest instrument on the trip' or equivalent for second B1. Examples of common acceptable instruments; double bass, cello, harp, piano, drums, tuba Examples of common unacceptable instruments: violin, viola, trombone, trumpet, french horn, guitar</p> <p>'Quartiles equidistant from median' or equivalent award B1 then symmetrical or no skew for B1 Alternative: 'Positive tail is longer than negative tail' or 'median closer to lowest value' or equivalent so slight positive skew. B0 for 'evenly' etc. instead of 'symmetrical' B0 for 'normal' only</p> <p><b>Please note that B mark appears first on ePEN</b> First line might be missing so first M1 can be implied by second. Second M1 for standardising with sigma and equating to z value NB Using 0.7734 should not be awarded second M1 Anything which rounds to 0.67 for B1. Accept 0.675 if to 3sf obtained by interpolation Anything that rounds to 13.3 – 13.4 for A1.</p>	

<p>3(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>Use overlay</p> $S_{xy} = 28750 - \frac{315 \times 620}{8} = 4337.5$ <p><b>**answer given**</b> so award for method</p> $S_{xx} = 15225 - \frac{315^2}{8} = 2821.875$ $b = \frac{4377.5}{S_{xx}} = 1.537... = 1.5$ $a = \bar{y} - b\bar{x} = \frac{620}{8} - b\frac{315}{8} = 16.97... = 17.0$ <p>Use overlay</p> <p>Brand D, since a long way above / from the line Using line: <math>y = 17 + 35 \times 1.5 = 69.5</math></p> <p style="text-align: right;">dependent upon 'Brand D' above</p>	<p><b>B2</b> <b>(2)</b></p> <p><b>M1</b></p> <p><b>M1A1</b> <b>(3)</b></p> <p><b>M1,A1</b></p> <p><b>M1,A1</b> <b>(4)</b></p> <p><b>B1</b> <b>B1</b> <b>(2)</b></p> <p><b>B1</b> <b>M1A1</b> <b>(4)</b></p> <p><b>Total 15 marks</b></p>
<p>Notes:</p> <p>3(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>Points B2, within 1 small square of correct point, subtract 1 mark each error minimum 0.</p> <p>Anything that rounds to 2820 for A1</p> <p>Anything that rounds to 1.5 and 17.0 (accept 17)</p> <p>Follow through for the intercept for first B1. Correct slope of straight line for second B1.</p> <p>Anything that rounds to 69p-71p for final A1. Reading from graph is acceptable for M1A1. If value read from graph at <math>x = 35</math> is answer given but out of range, then award M1A0.</p>	

<p>4(a)</p> <p>(b)</p> <p>(c)</p>	$P(Q \cup T) = 0.6$ $P(Q) + P(T) - P(Q \cap T) = 0.6$ $P(Q \cap T) = 0.1$  $P(Q \cap T   Q \cup T) = \frac{0.15}{0.60} = \frac{1}{4} \text{ or } 0.25 \text{ or } 25\%$	<p><b>B1</b> <b>M1</b> <b>A1</b> <b>(3)</b></p> <p><b>M1</b> <b>A1</b> <b>B1</b> <b>(3)</b></p> <p><b>M1A1</b> <b>A1</b> <b>(3)</b></p> <p><b>Total 9 marks</b></p>
<p>Notes:</p> <p>4(a)</p> <p>(b)</p> <p>(c)</p>	<p>B1 for 0.6 M1 for use of <math>P(Q) + P(T) - P(Q \cap T) = P(Q \cup T)</math> 0.1 Correct answer only for A1 Alternative: (25+45+40=)110% B1 110-100=10% M1A1 0.1 stated clearly as the final answer with no working gets 3/3</p> <p>Two intersecting closed curves for M1, no box required. At least one label (<math>Q</math> or <math>T</math>) required for first A1. Follow through (0.25- 'their 0.1') and (0.45- 'their 0.1') for A1. 0.4 and box required, correct answer only for B1 Using %, whole numbers in Venn diagram without % sign, whole numbers in correct ratio all OK</p> <p>Require fraction with denominator 0.6 or their equivalent from Venn diagram for M1 Follow through their values in fraction for A1 Final A1 is correct answer only. <u>No working no marks.</u></p>	

5(a)	18-25 group, area=7x5=35 25-40 group, area=15x1=15	<b>B1</b> <b>B1</b> (2)
(b)	$(25-20) \times 5 + (40-25) \times 1 = 40$	<b>M1A1</b> (2)
(c)	Mid points are 7.5, 12, 16, 21.5, 32.5 $\sum f = 100$ $\frac{\sum ft}{\sum f} = \frac{1891}{100} = 18.91$	<b>M1</b> <b>B1</b> <b>M1A1</b> (4)
(d)	$\sigma_t = \sqrt{\frac{41033}{100} - \bar{t}^2}$ $\sigma_t = \sqrt{52.74\dots} = 7.26$ $\sqrt{\frac{n}{n-1} \left( \frac{41033}{100} - \bar{t}^2 \right)}$ alternative OK	<b>M1</b> <b>M1</b> <b>A1</b> (3)
(e)	$Q_2 = 18$ or 18.1 if (n+1) used $Q_1 = 10 + \frac{15}{16} \times 4 = 13.75$ or 15.25 numerator gives 13.8125 $Q_3 = 18 + \frac{25}{35} \times 7 = 23$ or 25.75 numerator gives 23.15	<b>B1</b> <b>M1A1</b> <b>A1</b> (4)
(f)	0.376... Positive skew	<b>B1</b> <b>B1</b> (2) <b>Total 17 marks</b>
Notes: 5(b)	5x5 is enough evidence of method for M1. Condone 19.5, 20.5 instead of 20 etc. Award 2 if 40 seen. Look for working for this question in part (d) too. Use of some mid-points, at least 3 correct for M1. These may be tabulated in (d). Their $\frac{\sum ft}{\sum f}$ for M1 and anything that rounds to 18.9 for A1.	
(d)	Clear attempt at $\frac{41033}{100} - \bar{t}^2$ or $\frac{n}{n-1} \left( \frac{41033}{100} - \bar{t}^2 \right)$ alternative for first M1. They may use their $\bar{t}$ and gain the method mark. Square root of above for second M1 Anything that rounds to 7.3 for A1.	
(e)	Clear attempt at either quartile for M1 These will take the form 'their lower limit' + correct fraction x 'their class width'. Anything that rounds to 13.8 for lower quartile. 23 or anything that rounds to 23.2 dependent upon method used.	
(f)	Anything that rounds to 0.38 for B1 or 0.33 for B1 if (n+1) used. Correct answer or correct statement that follows from their value for B1.	

<p>6(a)</p>	$P(X > 25) = P\left(Z > \frac{25 - 20}{4}\right)$ $= P(Z > 1.25)$ $= 1 - 0.8944$ $= 0.1056$	<p><b>M1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>(3)</b></p>
<p>(b)</p>	$P(X < 20) = 0.5 \text{ so } P(X < d) = 0.5 + 0.4641 = 0.9641$ $P(Z < z) = 0.9641, z = 1.80$ $\frac{d - 20}{4} = 1.80$ $d = 27.2$	<p><b>B1</b></p> <p><b>B1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p><b>(4)</b></p> <p><b>Total 7 marks</b></p>
<p>Notes:</p>	<p>(a)</p> <p>Standardise with 20 and 4 for M1, allow numerator 20-25          1- probability for second M1          Anything that rounds to 0.106 for A1.          Correct answer with no working award 3/3</p> <p>(b)</p> <p>0.9641 seen or implied by 1.80 for B1          1.80 seen for B1          Standardise with 20 and 4 and equate to z value for M1          Z=0.8315 is M0          Anything that rounds to 27.2 for final A1.          Correct answer with no working 4/4</p>	

<p>7(a)</p>	$p + q = 0.45$ $\sum xP(X = x) = 4.5$ $3p + 7q = 1.95$	<p><b>B1</b> <b>M1</b> <b>A1</b> <b>(3)</b></p>
<p>(b)</p>	<p>Attempt to solve equations in (a)</p> $q = 0.15$ $p = 0.30$	<p><b>M1</b> <b>A1</b> <b>A1</b> <b>(3)</b></p>
<p>(c)</p>	$P(4 < X < 7) = P(5) + P(7)$ $= 0.2 + q = 0.35$	<p><b>M1</b> <b>A1</b> <b>(2)</b></p>
<p>(d)</p>	$\text{Var}(X) = E(X^2) - [E(X)]^2 = 27.4 - 4.5^2$ $= 7.15$	<p><b>M1</b> <b>A1</b> <b>(2)</b></p>
<p>(e)</p>	$E(19 - 4X) = 19 - 4 \times 4.5 = 1$	<p><b>B1</b> <b>(1)</b></p>
<p>(f)</p>	$\text{Var}(19 - 4X) = 16\text{Var}(X)$ $= 16 \times 7.15 = 114.4$	<p><b>M1</b> <b>A1</b> <b>(2)</b></p>
<b>Total 13 marks</b>		
<p>Notes:</p>	<p>7(a) <math>0.55 + p + q = 1</math> award B1. Not seen award B0.  <math>0.2 + 3p + 1 + 7q + 1.35 = 4.5</math> or equivalent award M1A1  <math>3p + 7q + k = 4.5</math> award M1.</p> <p>(b) Attempt to solve must involve 2 linear equations in 2 unknowns  Correct answers only for accuracy.  Correct answers with no working award 3/3</p> <p>(c) Follow through accuracy mark for their <math>q</math>, <math>0 &lt; q &lt; 0.8</math></p> <p>(d) Attempt to substitute <u>given</u> values <u>only</u> into correct formula for M1.  7.15 only for A1  7.15 seen award 2/2</p> <p>(f) Accept 'invisible brackets' i.e. <math>-4^2 \text{Var}(X)</math> provided answer positive.  Anything that rounds to 114 for A1.</p>	

# Mark Scheme (Results)

## January 2008

GCE

GCE Mathematics (6683/01)


January 2008  
6683 Statistics S1  
Mark Scheme

Question Number	Scheme	Marks
1. (a)	$\sum x = 773, \sum y = 724$ $r = \frac{10 \times 56076 - 773 \times 724}{\sqrt{(10 \times 60475 - 773^2)(10 \times 53122 - 724^2)}} \quad \text{o.e.}$ $r = 0.155357 \dots$	<b>B1, B1</b> <b>M1 A1ft</b> <b>A1</b> <b>(5)</b>
(b)	<p>Both weak correlation Neither score is a good indication of future performance Interview test is slightly better since correlation is positive</p>	<b>B1g B1h</b> <b>(2)</b> <b>Total 7 marks</b>
NB (a)	$S_{xx} = 60475 - \frac{(773)^2}{10} = 722.1, \quad S_{yy} = 53122 - \frac{(724)^2}{10} = 704.4, \quad S_{xy} = 56076 - \frac{773 \times 724}{10} = 110.8$ <p>1<sup>st</sup> B1 for <math>\sum x</math> and 2<sup>nd</sup> B1 for <math>\sum y</math>, should be seen or implied.</p> <p>M1 for at least one correct attempt at one of <math>S_{xx}</math>, <math>S_{yy}</math> or <math>S_{xy}</math> and then using in the correct formula</p> <p>1<sup>st</sup> A1ft for a fully correct expression. (ft their <math>\sum x</math> and their <math>\sum y</math>) or 3 correct expressions for <math>S_{xx}</math>, <math>S_{xy}</math>, and <math>S_{yy}</math> but possibly incorrect values for these placed correctly in <math>r</math>.</p>	
(b)	<p>2<sup>nd</sup> A1 for awrt 0.155</p> <p>If <math> r  &gt; 0.5</math> they can score B1g in (b) for saying that it (skills test) is not a good guide to performance but B0h since a second acceptable comment about both tests is not possible.</p> <p>Give B1 for one correct line, B1B1 for any 2. If the only comment is the test(s) <u>are</u> a good guide: scores B0B0 If the only comment is the tests are not good: scores B1B0 (second line)</p> <p>The third line is for a comment that suggests that the interview test is OK but the skills test is not since one is positive and the other is negative.</p> <p>Treat 1<sup>st</sup> B1 as B1g and 2<sup>nd</sup> as B1h</p> <p>An answer of “no” alone scores B0B0</p>	

Question Number	Scheme	Marks
<p>2.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>mean is <math>\frac{2757}{12}, = 229.75</math></p> <p>sd is <math>\sqrt{\frac{724961}{12} - (229.75)^2}, = 87.34045</math></p> <p>[Accept <math>s =</math> AWRT 91.2]</p> <p>Ordered list is: 125, 160, 169, 171, 175, 186, 210, 243, 250, 258, 390, 420</p> <p><math>Q_2 = \frac{1}{2}(186 + 210) = 198</math></p> <p><math>Q_1 = \frac{1}{2}(169 + 171) = 170</math></p> <p><math>Q_3 = \frac{1}{2}(250 + 258) = 254</math></p> <p><math>Q_3 + 1.5(Q_3 - Q_1) = 254 + 1.5(254 - 170), = 380</math>      Accept AWRT (370-392)</p> <p>Patients <math>F</math> (420) and <math>B</math> (390) are outliers.</p> <p><math>\frac{Q_1 - 2Q_2 + Q_3}{Q_3 - Q_1} = \frac{170 - 2 \times 198 + 254}{254 - 170}, = 0.\dot{3}</math></p> <p>Positive skew.</p>	<p>AWRT 230    <b>M1, A1</b></p> <p>AWRT 87.3    <b>M1, A1</b></p> <p><b>(4)</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>B1</b></p> <p><b>(3)</b></p> <p><b>M1, A1</b></p> <p><b>B1ft B1ft</b></p> <p><b>(4)</b></p> <p>AWRT 0.33    <b>M1, A1</b></p> <p><b>A1ft</b></p> <p><b>(3)</b></p> <p><b>Total 14 marks</b></p>
<p>(a)</p> <p>NB</p> <p>(b)</p> <p>S.C.</p> <p>(c)</p> <p>(d)</p>	<p>1<sup>st</sup> M1      for using <math>\frac{\sum x}{n}</math> with a credible numerator and <math>n = 12</math>.</p> <p>2<sup>nd</sup> M1      for using a correct formula, root required but can ft their mean</p> <p>Use of <math>s = \sqrt{8321.84...} = 91.22...</math> is OK for M1A1 here.</p> <p><b>Answers only from a calculator in (a) can score full marks</b></p> <p>1<sup>st</sup> B1 for median= 198 only, 2<sup>nd</sup> B1 for lower quartile 3<sup>rd</sup> B1 for upper quartile</p> <p>If all <math>Q_1</math> and <math>Q_3</math> are incorrect but an ordered list (with <math>\geq 6</math> correctly placed) is seen and used then award B0B1 as a special case for these last two marks.</p> <p>M1            for a clear attempt using their quartiles in given formula,</p> <p>A1            for any value in the range 370 - 392</p> <p>1<sup>st</sup> B1ft      for any one correct decision about <math>B</math> or <math>F</math> - ft their limit in range (258, 420)</p> <p>2<sup>nd</sup> B1ft      for correct decision about both <math>F</math> and <math>B</math> - ft their limit in range (258, 420)</p> <p>If more points are given score B0 here for the second B mark. ( Can score M0A0B1B1 here)</p> <p>M1            for an attempt to use their figures in the correct formula – must be seen (<math>\geq 2</math> correct substitutions)</p> <p>1<sup>st</sup> A1        for AWRT 0.33</p> <p>2<sup>nd</sup> A1ft      for positive skew. Follow through their value/sign of skewness . Ignore any further calculations. “positive correlation” scores A0</p>	

<p>3.</p>	<table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>Width</td> <td>1</td> <td>1</td> <td>4</td> <td>2</td> <td>3</td> <td>5</td> <td>3</td> <td>12</td> </tr> <tr> <td>Freq. Density</td> <td>6</td> <td>7</td> <td>2</td> <td>6</td> <td>5.5</td> <td>2</td> <td>1.5</td> <td>0.5</td> </tr> </table> <p style="text-align: right; margin-right: 20px;"><math>0.5 \times 12</math> or 6</p> <p>Total area is <math>(1 \times 6) + (1 \times 7) + (4 \times 2) + \dots = 70</math></p> <p><math>(90.5 - 78.5) \times \frac{1}{2} \times \frac{140}{\text{their } 70}</math></p> <p style="text-align: right; margin-right: 20px;">“70 seen anywhere”</p> <p>Number of runners is 12</p>	Width	1	1	4	2	3	5	3	12	Freq. Density	6	7	2	6	5.5	2	1.5	0.5	<p><b>M1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>B1</b></p> <p><b>A1</b></p> <p style="text-align: right;"><b>(5)</b></p> <p><b>Total 5 marks</b></p>
Width	1	1	4	2	3	5	3	12												
Freq. Density	6	7	2	6	5.5	2	1.5	0.5												
	<p>1<sup>st</sup> M1 for attempt at width of the correct bar (90.5 - 78.5) [Maybe on histogram or in table]</p> <p>1<sup>st</sup> A1 for <math>0.5 \times 12</math> or 6 (may be seen on the histogram. Must be related to the area of the bar above 78.5 - 90.5.</p> <p>2<sup>nd</sup> M1 for attempting area of correct bar <math>\times \frac{140}{\text{their } 70}</math></p> <p>B1 for 70 seen anywhere in their working</p> <p>2<sup>nd</sup> A1 for correct answer of 12.</p> <p>Minimum working required is <math>2 \times 0.5 \times 12</math> where the 2 should come from <math>\frac{140}{70}</math></p> <p>Beware <math>90.5 - 78.5 = 12</math> (this scores M1A0M0B0A0)</p> <p>Common answer is <math>0.5 \times 12 = 6</math> (this scores M1A1M0B0A0)</p> <p>If unsure send to review e.g. <math>2 \times 0.5 \times 12 = 12</math> without 70 being seen</p>																			

<p>4.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	$S_{xy} = 1818.5 - \frac{41 \times 406}{10}, = 153.9$ <p>(could be seen in (b))</p> $S_{xx} = 188 - \frac{41^2}{10} = 19.9$ <p>(could be seen in (b))</p> $b = \frac{153.9}{19.9}, = 7.733668....$ $a = 40.6 - b \times 4.1 (= 8.89796....)$ $y = 8.89 + 7.73x$ <p>A typical car will travel 7700 miles every year</p> $x = 5, y = 8.89 + 7.73 \times 5 (= 47.5 - 47.6)$ <p>So mileage predicted is</p> <p style="text-align: right;">AWRT 48000</p>	<p>AWRT 154</p> <p>AWRT 7.73</p> <p>AWRT 48000</p> <p>M1, A1</p> <p>A1</p> <p>M1, A1</p> <p>M1</p> <p>A1</p> <p>B1ft</p> <p>M1</p> <p>A1</p> <p>(3)</p> <p>(4)</p> <p>(1)</p> <p>(2)</p> <p><b>Total 10 marks</b></p>
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p style="text-align: center;"><b>Accept calculations for <math>S_{xx}</math> and <math>S_{xy}</math> in (a) or (b)</b></p> <p>M1 for correct attempt or expression for either</p> <p>1<sup>st</sup> A1 for one correct</p> <p>2<sup>nd</sup> A1 for both correct</p> <p><b>Ignore the open marks for part (b) they should be awarded as per this scheme</b></p> <p>1<sup>st</sup> M1 for <math>\frac{\text{their } S_{xy}}{\text{their } S_{xx}}</math></p> <p>1<sup>st</sup> A1 for AWRT 7.73</p> <p>2<sup>nd</sup> M1 for attempt at correct formula for <math>a</math> (minus required). Ft their <math>b</math>. Quoting a correct formula but making one slip in sub.eg. <math>\bar{y} = 406</math> is OK</p> <p>2<sup>nd</sup> A1 for correct equation with 2dp accuracy. Accept <math>a = 8.89</math>, and <math>b = 7.73</math> even if not written as final equation.</p> <p><b>Correct answers only (from calc) score 4/4 if correct to 2dp or 3/4 if AWRT 2dp</b></p> <p>B1ft for their <math>b \times 1000</math> to at least 2 sf. Accept “7.7 thousand” but value is needed</p> <p>M1 for substituting <math>x = 5</math> into their final answer to (b).</p> <p>A1 for AWRT 48000 (Accept “48 thousands”)</p>	

<p>5. (a)</p>	<p><b>Diagram may be drawn with <math>B \subset (A \cup C)</math> or with the 0 for <math>B \cap (A \cup C)</math>' simply left blank</b></p> <div style="text-align: center;">  </div> <p>Accept decimals or probs. in Venn diagram</p>	<p>3cc 90,3,2,1 1,(0),2 1 outside Box</p> <p><b>M1</b> <b>A1</b> <b>M1A1</b> <b>A1</b> <b>B1</b></p> <p style="text-align: right;"><b>(6)</b></p>
<p>(b)</p>	<p>P(none)=0.01</p>	<p><b>B1ft</b></p>
<p>(c)</p>	<p>P(A but not B)=0.04</p>	<p><b>M1 A1ft</b></p>
<p>(d)</p>	<p>P(any wine but C)=0.03</p>	<p><b>M1A1ft</b></p>
<p>(e)</p>	<p>P(exactly two)=0.06</p>	<p><b>M1A1ft</b></p>
<p>(f)</p>	<p><math>P(C A) = \frac{P(C \cap A)}{P(A)} = \frac{93}{96}</math> or <math>\frac{31}{32}</math> or AWRT 0.969</p>	<p><b>M1A1ft,A1</b></p>
<p><b>Total 16 marks</b></p>		
<p>(a)</p>	<p>1<sup>st</sup> M1 for 3 closed, labelled curves that overlap. A1 for the 90, 3, 2 and 1                  2<sup>nd</sup> M1 for one of 1, 0 or 2 correct <u>or</u> a correct sum of 4 values for A, B or C                  2<sup>nd</sup> A1 for all 7 values correct. Accept a blank instead of 0.  <b>NB final mark is a B1 for the box not an A mark as on EPEN</b>  <b>In parts (b) to (f) full marks can be scored for correct answers or correct ft</b></p>	<div style="border: 1px solid black; padding: 5px; width: fit-content; margin-left: auto;"> <p>For M marks in (c) to (e) they must have a fraction</p> </div>
<p>(b)</p>	<p>B1ft Follow through their '1' from outside divided by 100</p>	
<p>(c)</p>	<p>M1 for correct expression eg <math>P(A \cup B) - P(B)</math> or calculation e.g. 3 + 1 or 4 on top</p>	
<p>(d)</p>	<p>A1 for a correct probability, follow through with their '3+1' from diagram</p>	
<p>(d)</p>	<p>M1 for correct expression or calculation e.g. 1+2+0 or 99-96 or 3 on top</p>	
<p>(e)</p>	<p>A1 for a correct probability, follow through their '2+1+0' from diagram</p>	
<p>(e)</p>	<p>M1 for a correct expression or calculation e.g. 3+2+1 or 6 on top</p>	
<p>(f)</p>	<p>M1 for a correct expression upto " , " and <u>some</u> correct substitution, ft their values.</p>	
<p>(f)</p>	<p>One of these probabilities must be correct or correct ft. If P(C) on bottom M0</p>	
<p>(f)</p>	<p>1<sup>st</sup> A1ft follow through their <math>A \cap C</math> and their A but the ratio must be in (0, 1)</p>	
<p>(f)</p>	<p>2<sup>nd</sup> A1 for correct answer only. Answer only scores 3/3, but check working <math>P(A \cap C) / P(C)</math> is M0</p>	

<p>6. (a) (b) (c)</p>	<p>200 or 200g</p> <p><math>P(190 &lt; X &lt; 210) = 0.6</math> or <math>P(X &lt; 210) = 0.8</math> or <math>P(X &gt; 210) = 0.2</math> or diagram (o.e.) Correct use of 0.8 or 0.2</p> $Z = (\pm) \frac{210 - 200}{\sigma}$ $\frac{10}{\sigma} = 0.8416$ $\sigma = 11.882129\dots$ <p>0.8416 AWRT 11.9</p> <p><math>P(X &lt; 180) = P\left(Z &lt; \frac{180 - 200}{\sigma}\right)</math>  <math>= P(Z &lt; -1.6832)</math>  <math>= 1 - 0.9535</math>  <math>= 0.0465</math> or AWRT 0.046</p>	<p><b>B1</b> <b>M1</b> <b>A1</b> <b>M1</b> <b>B1</b> <b>A1</b> <b>M1</b> <b>M1</b> <b>A1</b> <b>A1</b> <b>(1)</b> <b>(5)</b> <b>(3)</b> <b>Total 9 marks</b></p>
<p>(a) (b) (c)</p>	<p>“mean = 200g” is B0 but “median = 200” or just “200” alone is B1</p> <p><b>Standardization in (b) and (c).</b> They must use <math>\sigma</math> not <math>\sigma^2</math> or <math>\sqrt{\sigma}</math>.</p> <p>1<sup>st</sup> M1 for a correct probability statement (as given or eg <math>P(200 &lt; X &lt; 210) = 0.3</math> o.e.) or shaded diagram - must have values on z-axis and probability areas shown  1<sup>st</sup> A1 for correct use of 0.8 or <math>p = 0.2</math>. Need a correct probability statement. May be implied by a suitable value for z seen (e.g. <math>z = 0.84</math>)  2<sup>nd</sup> M1 for attempting to standardise. Values for <math>x</math> and <math>\mu</math> used in formula. Don't need <math>z =</math> for this M1 nor a z-value, just mark standardization.  B1 for <math>z = 0.8416</math> (or better) [<math>z = 0.84</math> usually just loses this mark in (a)]  2<sup>nd</sup> A1 for AWRT 11.9</p> <p>1<sup>st</sup> M1 for attempting to Standardise with 200 and their sd(&gt;0) e.g. <math>(\pm) \frac{180 - 200}{\text{their } \sigma}</math>  2<sup>nd</sup> M1 <b>NB on open this is an A mark ignore and treat it as 2<sup>nd</sup> M1</b> for 1 – a probability from tables provided compatible with their probability statement.  A1 for 0.0465 or AWRT 0.046 (Dependent on both Ms in part (c))</p>	

7.(a)	$P(R = 3 \cap B = 0) = \frac{1}{4} \times \frac{1}{4} = \frac{1}{16}$	<b>M1, A1</b>  (2)																														
(b)	<table border="1" style="margin-left: auto; margin-right: auto;"> <tbody> <tr><td>3</td><td>0</td><td>3</td><td>6</td><td>9</td></tr> <tr><td>2</td><td>0</td><td>2</td><td>4</td><td>6</td></tr> <tr><td>1</td><td>0</td><td>1</td><td>2</td><td>3</td></tr> <tr><td>0</td><td>0</td><td>0</td><td>0</td><td>0</td></tr> <tr><td><i>B</i></td><td></td><td></td><td></td><td></td></tr> <tr><td><i>R</i></td><td>0</td><td>1</td><td>2</td><td>3</td></tr> </tbody> </table>	3	0	3	6	9	2	0	2	4	6	1	0	1	2	3	0	0	0	0	0	<i>B</i>					<i>R</i>	0	1	2	3	All 0s All 1,2,3s All 4,6,9s  <b>B1</b> <b>B1</b> <b>B1</b>  (3)
3	0	3	6	9																												
2	0	2	4	6																												
1	0	1	2	3																												
0	0	0	0	0																												
<i>B</i>																																
<i>R</i>	0	1	2	3																												
(c)	$a = \frac{7}{16}, b = c = d = \frac{1}{16}$	<b>B1, B1 B1</b>  (3)																														
(d)	$E(T) = \left(1 \times \frac{1}{16}\right) + \left(2 \times \frac{1}{8}\right) + \left(3 \times \frac{1}{8}\right) + \left(4 \times \frac{1}{16}\right) + \dots$ $= 2\frac{1}{4} \text{ or exact equivalent e.g. } 2.25, \frac{9}{4}$	<b>M1</b>  <b>A1</b>  (2)																														
(e)	$\text{Var}(T) = \left(1^2 \times \frac{1}{16}\right) + \left(2^2 \times \frac{1}{8}\right) + \left(3^2 \times \frac{1}{8}\right) + \left(4^2 \times \frac{1}{16}\right) + \dots - \left(\frac{9}{4}\right)^2$ $= \frac{49}{4} - \frac{81}{16} = 7\frac{3}{16} \text{ or } \frac{115}{16} \quad (\text{o.e.})$	<b>M1A1, M1</b>  AWRT 7.19 <b>A1</b> (4)																														
<b>Total 14 marks</b>																																
(a)	<b>M1</b> for $\frac{1}{4} \times \frac{1}{4}$																															
(c)	<b>1<sup>st</sup> B1</b> for $\frac{7}{16}$ ,																															
	<b>2<sup>nd</sup> B1</b> for only one error in $b, c, d$ ( $b = c = d \neq \frac{1}{16}$ or $b = c = \frac{1}{16} \neq d$ etc), <b>3<sup>rd</sup> B1</b> all of $b, c, d = \frac{1}{16}$																															
(d)	<b>M1</b> for attempting $\sum tP(T = t)$ , 3 or more terms correct or correct ft. Must Attempt to sum. NB calculating $E(T)$ and then dividing by a number other than 1 scores M0.																															
(e)	<b>1<sup>st</sup> M1</b> for attempt at $E(T^2)$ , 3 or more terms correct or correct ft.																															
	<b>1<sup>st</sup> A1</b> for $\frac{49}{4}$ (o.e.) or a fully correct expression (all non-zero terms must be seen)																															
	<b>2<sup>nd</sup> M1</b> for subtracting their $[E(T)]^2$ , Must be some attempt to square $-\frac{9}{4}$ is M0 but $-\frac{9}{16}$ could be M1																															
	<b>2<sup>nd</sup> A1</b> for correct fraction or AWRT 7.19 Full marks can still be scored in (d) and (e) if $a$ is incorrect																															


# Mark Scheme (Results)

## June 2008

GCE

### GCE Mathematics (6683/01)


**June 2008  
6683 Statistics S1  
Mark Scheme**

Question Number	Scheme	Marks
Q1 (a)	 <p style="text-align: right; margin-right: 10%;">                     Tree without probabilities or labels                      0.02(Disease), 0.95(Positive) on correct branches                      0.03(Positive) on correct branch.                 </p>	<p><b>M1</b> <b>A1</b> <b>A1</b></p> <p>[3]</p>
(b)	<p>P(Positive Test) = <math>0.02 \times 0.95 + 0.98 \times 0.03</math>                      = 0.0484</p>	<p><b>M1A1ft</b> <b>A1</b></p> <p>[3]</p>
(c)	<p>P(Do not have disease   Postive test) = <math>\frac{0.98 \times 0.03}{0.0484}</math>                      = 0.607438..</p> <p style="text-align: right; margin-right: 10%;">awrt 0.607</p>	<p><b>M1</b> <b>A1</b></p> <p>[2]</p>
(d)	<p>Test not very useful OR                      High probability of not having the disease for a person with a positive test</p>	<p><b>B1</b></p> <p>[1]</p>
	<p><u>Notes:</u>                      (a) M1: All 6 branches.                      Bracketed probabilities not required.                      (b) M1 for sum of two products, at least one correct from their diagram                      A1ft follows from the probabilities on their tree                      A1 for correct answer only or <math>\frac{121}{2500}</math>                      (c) M1 for conditional probability with numerator following from their tree and denominator their answer to part (b).                      A1 also for <math>\frac{147}{242}</math>.</p>	<p><b>Total 9</b></p>

Question Number	Scheme	Marks
<p>Q2</p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>50</p> <p><math>Q_1 = 45</math> <math>Q_2 = 50.5</math> <math>Q_3 = 63</math></p> <p>Mean = <math>\frac{1469}{28} = 52.464286..</math></p> <p>Sd = <math>\sqrt{\frac{81213}{28} - \left(\frac{1469}{28}\right)^2}</math> = 12.164.... or 12.387216...for divisor <math>n-1</math></p> <p><math>\frac{52.46.. - 50}{sd} = \text{awrt } 0.20 \text{ or } 0.21</math></p> <p>1. mode/median/mean Balmoral &gt; mode/median/mean Abbey 2. Balmoral sd &lt; Abbey sd or similar sd or correct comment from their values, Balmoral range &lt; Abbey range, Balmoral IQR &gt; Abbey IQR or similar IQR 3. Balmoral positive skew or almost symmetrical AND Abbey negative skew, Balmoral is less skew than Abbey or correct comment from their value in (d) 4. Balmoral residents generally older than Abbey residents or equivalent. Only one comment of each type max 3 marks</p>	<p><b>B1</b> [1]</p> <p><b>B1</b> <b>B1</b> <b>B1</b> [3]</p> <p>ONLY</p> <p>awrt 52.5 <b>M1A1</b></p> <p><b>M1</b> <b>A1</b> [4]</p> <p>awrt 12.2 or 12.4</p> <p><b>M1A1</b> [2]</p> <p><b>B1B1B1</b> [3]</p> <p><b>Total 13</b></p>
	<p><u>Notes:</u></p> <p>(c) M1 for their 1469 between 1300 and 1600, divided by 28, A1 for awrt 52.5 .. Please note this is B1B1 on Epen M1 use of correct formula including sq root A1 awrt 12.2 or 12.4 Correct answers with no working award full marks.</p> <p>(d) M1 for their values correctly substituted A1 Accept 0.2 as a special case of awrt 0.20 with 0 missing</p> <p>(e) Technical terms required in correct context in lines 1 to 3 e.g. 'average' and 'spread' B0 1 correct comment B1B0B0 2 correct comments B1B1B0 3 correct comments B1B1B1</p>	

Question Number	Scheme	Marks
<p>Q3 (a)</p> <p>(b)</p> <p>(c)</p>	$-1 \times p + 1 \times 0.2 + 2 \times 0.15 + 3 \times 0.15 = 0.55$ $p = 0.4$ $p + q + 0.2 + 0.15 + 0.15 = 1$ $q = 0.1$ $\text{Var}(X) = (-1)^2 \times p + 1^2 \times 0.2 + 2^2 \times 0.15 + 3^2 \times 0.15, -0.55^2$ $= 2.55 - 0.3025 = 2.2475$ $E(2X-4) = 2E(X) - 4$ $= -2.9$	<p><b>M1dM1</b></p> <p><b>A1</b></p> <p><b>M1</b></p> <p><b>A1</b></p> <p>[5]</p> <p><b>M1A1,M1</b></p> <p><b>A1</b></p> <p>awrt 2.25</p> <p>[4]</p> <p><b>M1</b></p> <p><b>A1</b></p> <p>[2]</p> <p><b>Total 11</b></p>
	<p><u>Notes:</u></p> <p>(a) M1 for at least 2 correct terms on LHS                  Division by constant e.g. 5 then M0                  dM1 dependent on first M1 for equate to 0.55 and attempt to solve.                  Award M1M1A1 for <math>p=0.4</math> with no working                  M1 for adding probabilities and equating to 1. All terms or equivalent required e.g. <math>p+q=0.5</math>                  Award M1A1 for <math>q=0.1</math> with no working</p> <p>(b) M1 attempting <math>E(X^2)</math> with at least 2 correct terms                  A1 for fully correct expression or 2.55                  Division by constant at any point e.g. 5 then M0                  M1 for subtracting their mean squared                  A1 for awrt 2.25                  Award awrt 2.25 only with no working then 4 marks</p> <p>(c) M1 for <math>2x(\text{their mean}) - 4</math>                  Award 2 marks for -2.9 with no working</p>	

Question Number	Scheme	Marks
Q4 (a)	$S_{tt} = 10922.81 - \frac{401.3^2}{15} = 186.6973$	awrt 187 <b>M1A1</b>
	$S_{vv} = 42.3356 - \frac{25.08^2}{15} = 0.40184$	awrt 0.402 <b>A1</b>
	$S_{tv} = 677.971 - \frac{401.3 \times 25.08}{15} = 6.9974$	awrt 7.00 <b>A1</b>
(b)	$r = \frac{6.9974}{\sqrt{186.6973 \times 0.40184}} = 0.807869$	awrt 0.808 <b>M1A1ft</b> <b>A1</b>
(c)	<p><math>t</math> is the explanatory variable as we can control temperature but not frequency of noise or equivalent comment</p>	<b>B1</b> <b>B1</b>
(d)	<p>High value of <math>r</math> or <math>r</math> close to 1 or Strong correlation</p>	<b>B1</b>
(e)	$b = \frac{6.9974}{186.6973} = 0.03748$	awrt 0.0375 <b>M1A1</b>
	$a = \frac{25.08}{15} - b \times \frac{401.3}{15} = 0.6692874$	awrt 0.669 <b>M1A1</b>
(f)	$t = 19, v = 0.6692874 + 0.03748 \times 19 = 1.381406$	awrt 1.4 <b>B1</b>
	<p><u>Notes:</u></p> <p>(a) M1 any one attempt at a correct use of a formula. Award full marks for correct answers with no working. Epen order of awarding marks as above.</p> <p>(b) M1 for correct formula and attempt to use A1ft for their values from part (a)</p> <p>NB Special Case for <math>\frac{677.971}{\sqrt{10922.81 \times 42.3356}}</math> M1A0</p> <p>A1 awrt 0.808 Award 3 marks for awrt 0.808 with no working</p> <p>(c) Marks are independent. Second mark requires some interpretation in context and can be statements such as ‘temperature effects / influences pitch or noise’ B1 ‘temperature is being changed’ BUT B0 for ‘temperature is changing’</p> <p>(e) M1 their values the right way up A1 for awrt 0.0375 M1 attempt to use correct formula with their value of <math>b</math> A1 awrt 0.669</p> <p>(f) awrt 1.4</p>	<b>[4]</b> <b>[3]</b> <b>[2]</b> <b>[1]</b> <b>[4]</b> <b>[1]</b> <b>Total 15</b>

Question Number	Scheme	Marks
<p>Q5 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<div style="display: flex; align-items: center;"> <div style="flex: 1;">  </div> <div style="flex: 1; padding-left: 20px;"> <p>3 closed intersecting curves with labels 100 100,30 12,10,3,25 Box</p> </div> </div> <p>(b) <math>P(\text{Substance } C) = \frac{100+100+10+25}{300} = \frac{235}{300} = \frac{47}{60}</math> or exact equivalent</p> <p>(c) <math>P(\text{All 3}   A) = \frac{10}{30+3+10+100} = \frac{10}{143}</math> or exact equivalent</p> <p>(d) <math>P(\text{Universal donor}) = \frac{20}{300} = \frac{1}{15}</math> or exact equivalent</p>	<p><b>M1</b> <b>A1</b> <b>A1</b> <b>B1</b> [4]</p> <p><b>M1A1ft</b> [2]</p> <p><b>M1A1ft</b> [2]</p> <p><b>M1A1 cao</b> [2] <b>Total 10</b></p>
	<p><u>Notes:</u></p> <p>(a) 20 not required. Fractions and exact equivalent decimals or percentages.</p> <p>(b) M1 For adding their positive values in C and finding a probability A1ft for correct answer or answer from their working</p> <p>(c) M1 their 10 divided by their sum of values in A A1ft for correct answer or answer from their working</p> <p>(d) M1 for 'their 20' divided by 300 A1 correct answer only</p>	

Question Number	Scheme	Marks								
<p>Q6 (a)</p> <p>(b)</p>	<p><math>F(4)=1</math>  <math>(4+k)^2 = 25</math>  <math>k = 1</math> as <math>k &gt; 0</math></p> <table border="1" data-bbox="220 472 1326 595"> <tr> <td><math>x</math></td> <td>2</td> <td>3</td> <td>4</td> </tr> <tr> <td><math>P(X=x)</math></td> <td><math>\frac{9}{25}</math></td> <td><math>\frac{7}{25}</math></td> <td><math>\frac{9}{25}</math></td> </tr> </table>	$x$	2	3	4	$P(X=x)$	$\frac{9}{25}$	$\frac{7}{25}$	$\frac{9}{25}$	<p><b>M1</b>  <b>A1</b></p> <p>[2]</p> <p><b>B1ftB1B1</b></p> <p>[3]</p> <p><b>Total 5</b></p>
$x$	2	3	4							
$P(X=x)$	$\frac{9}{25}$	$\frac{7}{25}$	$\frac{9}{25}$							
	<p><u>Notes:</u>                  (a) M1 for use of <math>F(4) = 1</math> only If <math>F(2)=1</math> and / or <math>F(3)=1</math> seen then M0.  <math>F(2)+F(3)+F(4)=1</math> M0                  A1 for <math>k=1</math> and ignore <math>k= -9</math></p> <p>(b) B1ft follow through their <math>k</math> for <math>P(X=2)</math> either exact or 3sf between 0 and 1 inclusive.                  B1 correct answer only or exact equivalent                  B1 correct answer only or exact equivalent</p>									

Question Number	Scheme	Marks
<p>Q7 (a)</p> <p>(b)</p> <p>(c)</p>	$z = \frac{53 - 50}{2}$ <p>Attempt to standardise</p> <p>1-probability required can be implied</p> $P(X > 53) = 1 - P(Z < 1.5)$ $= 1 - 0.9332$ $= 0.0668$ $P(X \leq x_0) = 0.01$ $\frac{x_0 - 50}{2} = -2.3263$ $x_0 = 45.3474$ <p>awrt 45.3 or 45.4</p> $P(2 \text{ weigh more than } 53\text{kg and } 1 \text{ less}) = 3 \times 0.0668^2 (1 - 0.0668)$ $= 0.012492487..$ <p>awrt 0.012</p>	<p><b>M1</b></p> <p><b>B1</b></p> <p><b>A1</b></p> <p>[3]</p> <p><b>M1</b></p> <p><b>M1B1</b></p> <p><b>M1A1</b></p> <p>awrt 45.3 or 45.4</p> <p><b>M1A1</b></p> <p>[5]</p> <p><b>B1M1A1ft</b></p> <p><b>A1</b></p> <p>[4]</p> <p><b>Total 12</b></p>
	<p><u>Notes:</u></p> <p>(a) M1 for using 53,50 and 2, either way around on numerator            B1 1- any probability for mark            A1 0.0668 cao</p> <p>(b) M1 can be implied or seen in a diagram            or equivalent with correct use of 0.01 or 0.99            M1 for attempt to standardise with 50 and 2 numerator either way around            B1 for <math>\pm 2.3263</math>            M1 Equate expression with 50 and 2 to a <math>z</math> value to form an equation with consistent signs and attempt to solve            A1 awrt 45.3 or 45.4</p> <p>(c) B1 for 3,            M1 <math>p^2(1-p)</math> for any value of <math>p</math>            A1ft for <math>p</math> is their answer to part (a) without 3            A1 awrt 0.012 or 0.0125</p>	

# Mark Scheme (Results)

## January 2009



GCE

GCE Mathematics (6683/01)


**January 2009  
6683 Statistics S1  
Mark Scheme**

Question Number	Scheme	Marks
<b>1</b>	$S_{xx} = 57.22 - \frac{(21.4)^2}{10} = 11.424$	M1 A1
(a)	$S_{xy} = 313.7 - \frac{21.4 \times 96}{10} = 108.26$	A1 (3)
(b)	$b = \frac{S_{xy}}{S_{xx}} = 9.4765\dots$ $a = \bar{y} - b\bar{x} = 9.6 - 2.14b = (-10.679\dots)$	M1 A1 M1 A1 (4)
(c)	$y = -10.7 + 9.48x$ <p>Every (extra) <u>hour</u> spent using the programme produces about <u>9.5 marks improvement</u></p>	B1ft (1)
(d)	$y = -10.7 + 9.48 \times 3.3, = 20.6$ <p style="text-align: right;">awrt 21</p>	M1,A1 (2)
(e)	Model may not be valid since [8h is] outside the range [0.5 - 4].	B1 (1)
<b>[11]</b>		
(a)	<p>M1 for a correct expression 1<sup>st</sup> A1 for AWRT 11.4 for <math>S_{xx}</math> 2<sup>nd</sup> A1 for AWRT 108 for <math>S_{xy}</math></p> <p>Correct answers only: One value correct scores M1 and appropriate A1, both correct M1A1A1</p>	
(b)	<p>1<sup>st</sup> M1 for using their values in correct formula 1<sup>st</sup> A1 for AWRT 9.5 2<sup>nd</sup> M1 for correct method for <math>a</math> (minus sign required) 2<sup>nd</sup> A1 for equation with <math>a</math> and <math>b</math> AWRT 3 sf (e.g. <math>y = -10.68 + 9.48x</math> is fine) Must have a full equation with <math>a</math> and <math>b</math> correct to awrt 3 sf</p>	
(c)	B1ft for comment conveying the idea of <u><math>b</math> marks per hour</u> . Must mention value of $b$ but can fit their value of $b$ . No need to mention “extra” but must mention “marks” and “hour(s)” e.g. “...9.5 times per hour...” scores B0	
(d)	<p>M1 for sub <math>x = 3.3</math> into their regression equation from the end of part (b) A1 for awrt 21</p>	
(e)	B1 for a statement that says or implies that it may <u>not</u> be valid because <u>outside the range</u> . They do not have to mention the values concerned here namely 8 h or 0.5 - 4	

Question Number	Scheme	Marks
<p>2</p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	<p><math>E = \text{take regular exercise}</math>      <math>B = \text{always eat breakfast}</math></p> <p><math>P(E \cap B) = P(E   B) \times P(B)</math></p> <p><math>= \frac{9}{25} \times \frac{2}{3} = 0.24</math>    or    <math>\frac{6}{25}</math>    or    <math>\frac{18}{75}</math></p> <p><math>P(E \cup B) = \frac{2}{3} + \frac{2}{5} - \frac{6}{25}</math>      or <math>P(E'   B')</math>      or <math>P(B' \cap E)</math>      or <math>P(B \cap E')</math></p> <p><math>= \frac{62}{75}</math>      <math>= \frac{13}{25}</math>      <math>= \frac{12}{75}</math>      <math>= \frac{32}{75}</math></p> <p><math>P(E' \cap B') = 1 - P(E \cup B) = \frac{13}{75}</math>    or    0.173</p> <p><math>P(E   B) = 0.36 \neq 0.40 = P(E)</math>    or    <math>P(E \cap B) = \frac{6}{25} \neq \frac{2}{5} \times \frac{2}{3} = P(E) \times P(B)</math></p> <p>So <math>E</math> and <math>B</math> are <u>not</u> statistically independent</p>	<p>M1</p> <p>A1    (2)</p> <p>M1</p> <p>A1</p> <p>M1 A1    (4)</p> <p>M1</p> <p>A1    (2)</p> <p>[8]</p>
<p>(a)</p> <p>(b)</p> <p>(c)</p>	<p>M1    for <math>\frac{9}{25} \times \frac{2}{3}</math> or <math>P(E B) \times P(B)</math> <u>and</u> at least one correct value seen. A1 for 0.24 or exact equiv. NB <math>\frac{2}{5} \times \frac{2}{3}</math> alone or <math>\frac{2}{5} \times \frac{9}{25}</math> alone scores M0A0. Correct answer scores full marks.</p> <p>1<sup>st</sup> M1    for use of the addition rule. Must have 3 terms and some values, can ft their (a) <u>Or</u> a full method for <math>P(E'   B')</math> requires <math>1 - P(E B')</math> and equation for <math>P(E B')</math>: <math>(a) + \frac{x}{3} = \frac{2}{5}</math> <u>Or</u> a full method for <math>P(B' \cap E)</math> <u>or</u> <math>P(B \cap E')</math> [ or other valid method]</p> <p>2<sup>nd</sup> M1    for a method leading to answer e.g. <math>1 - P(E \cup B)</math> <u>or</u> <math>P(B') \times P(E'   B')</math> <u>or</u> <math>P(B') - P(B' \cap E)</math> <u>or</u> <math>P(E') - P(B \cap E')</math></p> <p><u>Venn Diagram</u> 1<sup>st</sup> M1 for diagram with attempt at <math>\frac{2}{5} - P(B \cap E)</math> or <math>\frac{2}{3} - P(B \cap E)</math>. Can ft their (a)</p> <p>1<sup>st</sup> A1    for a correct first probability as listed or 32, 18 and 12 on Venn Diagram</p> <p>2<sup>nd</sup> M1    for attempting 75 - their (18 + 32 + 12)</p> <p>M1    for identifying suitable values to test for independence e.g. <math>P(E) = 0.40</math> and <math>P(E B) = 0.36</math> <u>Or</u> <math>P(E) \times P(B) = \dots</math> and <math>P(E \cap B) = \text{their (a)}</math> [but their (a) <math>\neq \frac{2}{5} \times \frac{2}{3}</math>]. Values seen somewhere</p> <p>A1    for correct values and a correct comment</p> <p><b>Diagrams</b> You may see these or find these useful for identifying probabilities.</p>	<p><b>Common Errors</b></p> <p>(a) <math>\frac{9}{25}</math> is M0A0</p> <p>(b) <math>P(E \cup B) = \frac{53}{75}</math> scores M1A0</p> <p><math>1 - P(E \cup B) = \frac{22}{75}</math> scores M1A0</p> <p>(b) <math>P(B') \times P(E') = \frac{1}{3} \times \frac{3}{5}</math> scores 0/4</p>



Question Number	Scheme	Marks																		
3	<p>(a) <math>E(X) = 0 \times 0.4 + 1 \times 0.3 + \dots + 3 \times 0.1, = 1</math></p> <p>(b) <math>F(1.5) = [P(X \leq 1.5) =] P(X \leq 1), = 0.4 + 0.3 = 0.7</math></p> <p>(c) <math>E(X^2) = 0^2 \times 0.4 + 1^2 \times 0.3 + \dots + 3^2 \times 0.1, = 2</math>  <math>\text{Var}(X) = 2 - 1^2, = 1 \quad (*)</math></p> <p>(d) <math>\text{Var}(5 - 3X) = (-3)^2 \text{Var}(X), = 9</math></p> <p>(e)</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Total</th> <th>Cases</th> <th>Probability</th> </tr> </thead> <tbody> <tr> <td rowspan="3" style="text-align: center;">4</td> <td><math>(X = 3) \cap (X = 1)</math></td> <td><math>0.1 \times 0.3 = 0.03</math></td> </tr> <tr> <td><math>(X = 1) \cap (X = 3)</math></td> <td><math>0.3 \times 0.1 = 0.03</math></td> </tr> <tr> <td><math>(X = 2) \cap (X = 2)</math></td> <td><math>0.2 \times 0.2 = 0.04</math></td> </tr> <tr> <td rowspan="2" style="text-align: center;">5</td> <td><math>(X = 3) \cap (X = 2)</math></td> <td><math>0.1 \times 0.2 = 0.02</math></td> </tr> <tr> <td><math>(X = 2) \cap (X = 3)</math></td> <td><math>0.2 \times 0.1 = 0.02</math></td> </tr> <tr> <td style="text-align: center;">6</td> <td><math>(X = 3) \cap (X = 3)</math></td> <td><math>0.1 \times 0.1 = 0.01</math></td> </tr> </tbody> </table> <p>Total probability = <math>0.03 + 0.03 + 0.04 + 0.02 + 0.02 + 0.01 = 0.15</math></p>	Total	Cases	Probability	4	$(X = 3) \cap (X = 1)$	$0.1 \times 0.3 = 0.03$	$(X = 1) \cap (X = 3)$	$0.3 \times 0.1 = 0.03$	$(X = 2) \cap (X = 2)$	$0.2 \times 0.2 = 0.04$	5	$(X = 3) \cap (X = 2)$	$0.1 \times 0.2 = 0.02$	$(X = 2) \cap (X = 3)$	$0.2 \times 0.1 = 0.02$	6	$(X = 3) \cap (X = 3)$	$0.1 \times 0.1 = 0.01$	<p>M1, A1 (2)</p> <p>M1, A1 (2)</p> <p>M1, A1 M1, A1cso (4)</p> <p>M1, A1 (2)</p> <p>B1B1B1</p> <p>M1 A1</p> <p>A1 (6) [16]</p>
Total	Cases	Probability																		
4	$(X = 3) \cap (X = 1)$	$0.1 \times 0.3 = 0.03$																		
	$(X = 1) \cap (X = 3)$	$0.3 \times 0.1 = 0.03$																		
	$(X = 2) \cap (X = 2)$	$0.2 \times 0.2 = 0.04$																		
5	$(X = 3) \cap (X = 2)$	$0.1 \times 0.2 = 0.02$																		
	$(X = 2) \cap (X = 3)$	$0.2 \times 0.1 = 0.02$																		
6	$(X = 3) \cap (X = 3)$	$0.1 \times 0.1 = 0.01$																		
ALT	<p>(a) M1 for at least 3 terms seen. Correct answer only scores M1A1. Dividing by <math>k (\neq 1)</math> is M0.</p> <p>(b) M1 for <math>F(1.5) = P(X \leq 1)</math>. [Beware: <math>2 \times 0.2 + 3 \times 0.1 = 0.7</math> but scores M0A0]</p> <p>(c) 1<sup>st</sup> M1 for at least 2 non-zero terms seen. <math>E(X^2) = 2</math> alone is M0. Condone calling <math>E(X^2) = \text{Var}(X)</math>.                      1<sup>st</sup> A1 is for an answer of 2 or a fully correct expression.                      2<sup>nd</sup> M1 for <math>-\mu^2</math>, condone <math>2 - 1</math>, unless clearly <math>2 -</math>. Allow <math>2 - \mu^2</math>, with <math>= 1</math> even if <math>E(X) \neq 1</math>                      2<sup>nd</sup> A1 for a fully correct solution with no incorrect working seen, <b>both</b> Ms required.  <math display="block">\underline{\sum (x - \mu)^2 \times P(X = x)}</math>                     1<sup>st</sup> M1 for an attempt at a full list of <math>(x - \mu)^2</math> values and probabilities. 1<sup>st</sup> A1 if all correct                      2<sup>nd</sup> M1 for at least 2 non-zero terms of <math>(x - \mu)^2 \times P(X = x)</math> seen. 2<sup>nd</sup> A1 for <math>0.4 + 0.2 + 0.4 = 1</math></p> <p>(d) M1 for use of the correct formula. <math>-3^2 \text{Var}(X)</math> is M0 unless the final answer is <math>&gt; 0</math>.</p> <p>(e) Can follow through their <math>\text{Var}(X)</math> for M1</p>																			
ALT	<p>1<sup>st</sup> B1 for all cases listed for a total of 4 or 5 or 6 . e.g. (2,2) counted twice for a total of 4 is B0</p> <p>2<sup>nd</sup> B1 for all cases listed for 2 totals }                      3<sup>rd</sup> B1 for a complete list of all 6 cases } These may be highlighted in a table</p> <p><u>Using Cumulative probabilities</u></p> <p>1<sup>st</sup> B1 for one or more cumulative probabilities used e.g. 2 then 2 or more or 3 then 1 or more</p> <p>2<sup>nd</sup> B1 for both cumulative probabilities used. 3<sup>rd</sup> B1 for a complete list 1, 3; 2, <math>\geq 2</math>; 3, <math>\geq 1</math></p> <p>M1 for one correct pair of correct probabilities multiplied</p> <p>1<sup>st</sup> A1 for all 6 correct probabilities listed (0.03, 0.03, 0.04, 0.02, 0.02, 0.01) needn't be added.</p> <p>2<sup>nd</sup> A1 for 0.15 or exact equivalent only as the final answer.</p>																			

Question Number	Scheme	Marks
4	<p>(a) <math>Q_2 = 53, Q_1 = 35, Q_3 = 60</math></p> <p>(b) <math>Q_3 - Q_1 = 25 \Rightarrow Q_1 - 1.5 \times 25 = -2.5</math> (no outlier)  <math>Q_3 + 1.5 \times 25 = 97.5</math> (so 110 is an outlier)</p> <p>(c) </p> <p>(d) <math>\sum y = 461, \sum y^2 = 24\ 219 \therefore S_{yy} = 24219 - \frac{461^2}{10}, = 2966.9</math> (*)</p> <p>(e) <math>r = \frac{-18.3}{\sqrt{3463.6 \times 2966.9}}</math> or <math>\frac{-18.3}{3205.64\dots} = -0.0057</math> AWR T - 0.006 or <math>-6 \times 10^{-3}</math></p> <p>(f) <math>r</math> suggests correlation is close to zero so parent's claim is not justified</p>	<p>B1, B1, B1 (3)</p> <p>M1</p> <p>A1 (2)</p> <p>M1</p> <p>A1ft</p> <p>A1ft (3)</p> <p>B1, B1, B1cso (3)</p> <p>M1 A1 (2)</p> <p>B1 (1)</p> <p>[14]</p>
	<p>(a) 1<sup>st</sup> B1 for median                  2<sup>nd</sup> B1 for lower quartile                  3<sup>rd</sup> B1 for upper quartile</p> <p>(b) M1 for attempt to find one limit                  A1 for both limits found and correct. No explicit comment about outliers needed.</p> <p>(c) M1 for a box and two whiskers                  1<sup>st</sup> A1ft for correct position of box, median and quartiles. Follow through their values.                  2<sup>nd</sup> A1ft for 17 and 77 or "their" 97.5 and *. If 110 is not an outlier then score A0 here.                  Penalise no gap between end of whisker and outlier. Must label outlier, needn't be with *.  <b>Accuracy</b> should be within the correct square so 97 or 98 will do for 97.5</p> <p>(d) 1<sup>st</sup> B1 for <math>\sum y</math> N.B. <math>(\sum y)^2 = 212521</math> and can imply this mark                  2<sup>nd</sup> B1 for <math>\sum y^2</math> or at least three correct terms of <math>\sum (y - \bar{y})^2</math> seen.                  3<sup>rd</sup> B1 for complete correct expression seen leading to 2966.9. So all 10 terms of <math>\sum (y - \bar{y})^2</math></p> <p>(e) M1 for attempt at correct expression for <math>r</math>. Can ft their <math>S_{yy}</math> for M1.</p> <p>(f) B1 for comment <u>rejecting</u> parent's claim on basis of <u>weak or zero</u> correlation                  Typical error is "negative correlation so comment is true" which scores B0                  Weak negative or weak positive correlation is OK as the basis for their rejection.</p>	

Question Number	Scheme	Marks
5	<p>(a) 8-10 hours: width = 10.5 - 7.5 = 3 represented by 1.5cm                      16-25 hours: width = 25.5 - 15.5 = 10 so represented by <u>5 cm</u>                      8- 10 hours: height = fd = 18/3 = 6 represented by 3 cm                      16-25 hours: height = fd = 15/10 = 1.5 represented by <u>0.75 cm</u></p> <p>(b) <math>Q_2 = 7.5 + \frac{(52-36)}{18} \times 3 = 10.2</math>  <math>Q_1 = 5.5 + \frac{(26-20)}{16} \times 2 [= 6.25 \text{ or } 6.3]</math> or <math>5.5 + \frac{(26.25-20)}{16} \times 2 [=6.3]</math>  <math>Q_3 = 10.5 + \frac{(78-54)}{25} \times 5 [= 15.3]</math> or <math>10.5 + \frac{(78.75-54)}{25} \times 5 [=15.45 \text{ \15.5}]</math>                      IQR = (15.3 - 6.3) = 9</p> <p>(c) <math>\sum fx = 1333.5 \Rightarrow \bar{x} = \frac{1333.5}{104} =</math> AWRT <u>12.8</u></p> <p>(d) <math>\sum fx^2 = 27254 \Rightarrow \sigma_x = \sqrt{\frac{27254}{104} - \bar{x}^2} = \sqrt{262.05 - \bar{x}^2}</math> AWRT <u>9.88</u>  <math>Q_3 - Q_2 [= 5.1] &gt; Q_2 - Q_1 [= 3.9]</math> or <math>Q_2 &lt; \bar{x}</math></p> <p>(e) So data is positively skew</p> <p>Use median and IQR,                      since data is skewed <u>or</u> not affected by extreme values or outliers</p>	<p>B1 M1 A1 (3)</p> <p>M1 A1</p> <p>A1</p> <p>A1 A1ft (5)</p> <p>M1 A1</p> <p>M1 A1 (4)</p> <p>B1ft dB1 (2)</p> <p>B1 B1 (2)</p> <p>[16]</p>
	<p>(a) M1 For attempting both frequency densities <math>\frac{18}{3} (= 6)</math> and <math>\frac{15}{10}</math>, <u>and</u> <math>\frac{15}{10} \times \text{SF}</math>, where <math>\text{SF} \neq 1</math></p> <p>(b) NB Wrong class widths( 2 and 9) gives <math>\frac{h}{1.66...} = \frac{3}{9} \rightarrow h = \frac{5}{9}</math> or 0.55... and scores M1A0</p> <p>M1 for identifying correct interval and a correct fraction e.g. <math>\frac{\frac{1}{2}(104)-36}{18}</math>. Condone 52.5 or 53</p> <p>1<sup>st</sup> A1 for 10.2 for median. Using (n + 1) allow awrt 10.3</p> <p>2<sup>nd</sup> A1 for a correct expression for either <math>Q_1</math> or <math>Q_3</math> (allow 26.25 and 78.75) NB: <u>Must see</u></p> <p>3<sup>rd</sup> A1 for correct expressions for both <math>Q_1</math> and <math>Q_3</math> <u>some</u></p> <p>(c) 4<sup>th</sup> A1ft for IQR, ft their quartiles. Using (n + 1) gives 6.28 and 15.45 <u>method</u></p> <p>1<sup>st</sup> M1 for attempting <math>\sum fx</math> and <math>\bar{x}</math></p> <p>(d) 2<sup>nd</sup> M1 for attempting <math>\sum fx^2</math> and <math>\sigma_x, \sqrt{\quad}</math> is needed for M1. Allow s = awrt 9.93</p> <p>1<sup>st</sup> B1ft for suitable test, values need not be seen but statement must be compatible with values used. Follow through their values</p> <p>2<sup>nd</sup> dB1 Dependent upon their test showing positive and for stating positive skew                      If their test shows negative skew they can score 1<sup>st</sup> B1 but lose the second</p> <p>(e) 1<sup>st</sup> B1 for choosing median and IQR. Must mention <u>both</u>. } <u>Award independently</u>                      2<sup>nd</sup> B1 for suitable reason }                      e.g. “use median because data is skewed” scores B0B1 since IQR is not mentioned</p>	

Question Number	Scheme	Marks
<p>6 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	$P(X < 39) = P\left(Z < \frac{39-30}{5}\right)$ $= P(Z < 1.8) = \underline{0.9641} \quad (\text{allow awrt } 0.964)$ $P(X < d) = P\left(Z < \frac{d-30}{5}\right) = 0.1151$ $1 - 0.1151 = 0.8849 \quad (\text{allow } \pm 1.2)$ $\Rightarrow z = -1.2$ $\therefore \frac{d-30}{5} = -1.2 \quad \underline{d = 24}$ $P(X > e) = 0.1151 \quad \text{so } e = \mu + (\mu - \text{their } d) \text{ or } \frac{e-30}{5} = 1.2 \text{ or } - \text{their } z$ $\underline{e = 36}$ $P(d < X < e) = 1 - 2 \times 0.1151$ $= 0.7698 \quad \text{AWRT } \underline{0.770}$	<p>M1</p> <p>A1 (2)</p> <p>M1</p> <p>B1</p> <p>M1A1 (4)</p> <p>M1</p> <p>A1 (2)</p> <p>M1</p> <p>A1 (2)</p> <p>[10]</p>
	<p><b>Answer only scores all marks in each section BUT check (b) and (c) are in correct order</b></p> <p>(a) M1 for standardising with <math>\sigma</math>, <math>z = \pm \frac{39-30}{5}</math> is OK  A1 for 0.9641 or awrt 0.964 but if they go on to calculate <math>1 - 0.9641</math> they get M1A0</p> <p>(b) 1<sup>st</sup> M1 for attempting <math>1 - 0.1151</math>. Must be seen in (b) in connection with finding <math>d</math>  B1 for <math>z = \pm 1.2</math>. They must state <math>z = \pm 1.2</math> or imply it is a <math>z</math> value by its use.  This mark is only available in part (b).  2<sup>nd</sup> M1 for <math>\left(\frac{d-30}{5}\right) =</math> their negative <math>z</math> value (or equivalent)</p> <p>(c) M1 for a full method to find <math>e</math>. If they used <math>z = 1.2</math> in (b) they can get M1 for <math>z = \pm 1.2</math> here  If they use symmetry about the mean <math>\mu + (\mu - \text{their } d)</math> then ft their <math>d</math> for M1  Must explicitly <u>see</u> the method used unless the answer is correct.</p> <p>(d) M1 for a complete method or use of a correct expression e.g. “their 0.8849” - 0.1151  <u>or If their <math>d &lt;</math> their <math>e</math> using their values with <math>P(X &lt; e) - P(X &lt; d)</math></u>  If their <math>d \geq</math> their <math>e</math> then they can only score from an argument like <math>1 - 2 \times 0.1151</math>  A negative probability or probability <math>&gt; 1</math> for part (d) scores M0A0</p>	

# Mark Scheme (Results)

## Summer 2009

GCE

GCE Mathematics (6683/01)

June 2009  
6683 Statistics S1  
Mark Scheme

Question Number	Scheme	Marks
Q1 (a)	$(S_{pp}) = 38125 - \frac{445^2}{10}$ $= 18322.5$ <p style="text-align: right;">awrt 18300</p> $(S_{pt}) = 26830 - \frac{445 \times 240}{10}$ $= 16150$ <p style="text-align: right;">awrt 16200</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p style="text-align: right;">(3)</p>
(b)	$r = \frac{"16150"}{\sqrt{"18322.5" \times 21760}}$ $= 0.8088\dots$ <p style="text-align: right;">awrt 0.809</p>	<p style="text-align: center;">Using their values for method</p> <p>M1</p> <p>A1</p> <p style="text-align: right;">(2)</p>
(c)	As the temperature increases the pressure increases.	<p>B1</p> <p style="text-align: right;">(1)</p> <p style="text-align: right;">[6]</p>
Notes	<p>1(a) M1 for seeing a correct expression <math>38125 - \frac{445^2}{10}</math> or <math>26830 - \frac{445 \times 240}{10}</math></p> <p>If no working seen, at least one answer must be exact to score M1 by implication.</p> <p>1(b) Square root and their values with 21760 all in the right places required for method. Anything which rounds to (awrt) 0.809 for A1.</p> <p>1(c) Require a correct statement in <b>context</b> using <u>temperature/heat</u> and <u>pressure</u> for B1.</p> <p>Don't allow "as <math>t</math> increases <math>p</math> increases".</p> <p>Don't allow proportionality.</p> <p>Positive correlation only is B0 since there is no interpretation.</p>	

Question Number	Scheme	Marks
<p>Q2 (a)</p> <p>(b)(i)</p> <p>(ii)</p> <p>(c)</p>	<div style="text-align: center;"> </div> <p>Correct tree All labels Probabilities on correct branches</p> <p><math>\frac{1}{3} \times \frac{1}{10} = \frac{1}{30}</math> or equivalent</p> <p><math>CNL + BNL + FNL = \frac{1}{2} \times \frac{4}{5} + \frac{1}{6} \times \frac{3}{5} + \frac{1}{3} \times \frac{9}{10}</math> <math>= \frac{4}{5}</math> or equivalent</p> <p><math>P(F' / L) = \frac{P(F' \cap L)}{P(L)}</math>      Attempt correct conditional probability <b>but see notes</b></p> <p><math>= \frac{\frac{1}{6} \times \frac{2}{5} + \frac{1}{2} \times \frac{1}{5}}{1 - (ii)}</math>      <math>\frac{\text{numerator}}{\text{denominator}}</math></p> <p><math>= \frac{5}{30} = \frac{5}{6}</math> or equivalent      cao</p>	<p>B1 B1</p> <p>B1</p> <p>(3)</p> <p>M1 A1</p> <p>(2)</p> <p>M1</p> <p>A1</p> <p>(2)</p> <p>M1</p> <p>A1 A1ft</p> <p>A1</p> <p>(4) [11]</p>
<p>Notes</p>	<p>Exact decimal equivalents required throughout if fractions not used e.g. 2(b)(i) 0.03</p> <p>Correct path through their tree given in their probabilities award Ms</p> <p>2(a) All branches required for first B1. Labels can be words rather than symbols for second B1. Probabilities from question enough for third B1 i.e. bracketed probabilities not required. Probabilities and labels swapped i.e. labels on branches and probabilities at end can be awarded the marks if correct.</p> <p>2(b)(i) Correct answer only award both marks.</p> <p>2(b)(ii) At least one correct path identified and attempt at adding all three multiplied pairs award M1</p> <p>2(c) Require probability on numerator and division by probability for M1. Require numerator correct for their tree for M1.</p> <p>Correct formula seen and used, accept denominator as attempt and award M1</p> <p>No formula, denominator must be correct for their tree or 1-(ii) for M1</p> <p>1/30 on numerator only is M0, P(L/F') is M0.</p>	

Question Number	Scheme	Marks
Q3 (a)  (b)	<p>1(cm) cao</p> <p>10 cm<sup>2</sup> represents 15 10/15 cm<sup>2</sup> represents 1</p> <p>Therefore frequency of 9 is <math>\frac{10}{15} \times 9</math> or <math>\frac{9}{1.5}</math></p> <p>height = 6(cm)</p> <p style="text-align: right;">or 1cm<sup>2</sup> represents 1.5</p> <p style="text-align: right;">Require <math>\times \frac{2}{3}</math> or <math>\div 1.5</math></p>	B1     M1 A1  [3]
Notes	<p>If 3(a) and 3(b) incorrect, but their (a) x their (b)=6 then award BOM1A0</p> <p>3(b) Alternative method: f/cw=15/6=2.5 represented by 5 so factor x2 award M1 So f/cw=9/3=3 represented by 3x2=6. Award A1.</p>	

Question Number	Scheme	Marks
<p>Q4 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p><math>Q_2 = 17 + \left(\frac{60-58}{29}\right) \times 2</math>  <math>= 17.1</math> (17.2 if use 60.5)      awrt 17.1 (or17.2)</p> <p><math>\sum fx = 2055.5</math>      <math>\sum fx^2 = 36500.25</math>      Exact answers can be seen below or implied by correct answers.                      Evidence of attempt to use midpoints with at least one correct</p> <p>Mean = 17.129...      awrt 17.1</p> <p><math>\sigma = \sqrt{\frac{36500.25}{120} - \left(\frac{2055.5}{120}\right)^2}</math>  <math>= 3.28</math> (s= 3.294)      awrt 3.3</p> <p><math>\frac{3(17.129 - 17.1379...)}{3.28} = -0.00802</math>      Accept 0 or awrt 0.0</p> <p>No skew/ slight skew</p> <p>The skewness is very small. Possible.</p>	<p>M1</p> <p>A1 (2)</p> <p>B1 B1</p> <p>M1</p> <p>B1</p> <p>M1</p> <p>A1 (6)</p> <p>M1 A1</p> <p>B1 (3)</p> <p>B1 B1dep (2) [13]</p>
<p>Notes</p>	<p>4(a) Statement of <math>17 + \frac{\text{freq into class}}{\text{class freq}} \times cw</math> and attempt to sub or</p> <p><math>\frac{m-17}{19-17} = \frac{60(.5)-58}{87-58}</math> or equivalent award M1                      cw=2 or 3 required for M1.                      17.2 from cw=3 award A0.</p> <p>4(b) Correct <math>\sum fx</math> and <math>\sum fx^2</math> can be seen in working for both B1s                      Midpoints seen in table and used in calculation award M1                      Require complete correct formula including use of square root and attempt to sub for M1. No formula stated then numbers as above or follow from (b) for M1  <math>(\sum fx)^2, \sum (fx)^2</math> or <math>\sum f^2x</math> used instead of <math>\sum fx^2</math> in sd award M0                      Correct answers only with no working award 2/2 and 6/6</p> <p>4(c) Sub in their values into given formula for M1</p> <p>4(d) No skew / slight skew / 'Distribution is almost symmetrical' / 'Mean approximately equal to median' or equivalent award first B1. Don't award second B1 if this is not the case. Second statement should imply 'Greg's suggestion that a normal distribution is suitable is possible' for second B1 dep.                      If B0 awarded for comment in (c).and (d) incorrect, allow follow through from the <b>comment</b> in (c).</p>	

Question Number	Scheme	Marks
Q5 (a) $b = \frac{59.99}{33.381}$ $= 1.79713\dots$ $a = 32.7 - 1.79713\dots \times 51.83$ $= -60.44525\dots$ $w = -60.445251\dots + 1.79713\dots l$ (b) $w = -60.445251\dots + 1.79713\dots \times 60$ $= 47.3825\dots$ (c) It is extrapolating so (may be) unreliable.	1.8 or awrt 1.80 awrt -60 <i>l</i> and <i>w</i> required and awrt 2sf In range 47.3 – 47.6 inclusive	M1 A1 M1 A1 A1ft (5) M1 A1 (2) B1, B1dep (2) [9]
Notes	5(a) Special case $b = \frac{59.99}{120.1} = 0.4995 \text{ M0A0}$ $a = 32.7 - 0.4995 \times 51.83 \text{ M1A1}$ $w = 6.8 + 0.50l \text{ at least 2 sf required for A1}$ 5(b) Substitute into their answer for (a) for M1 5(c) ‘Outside the range on the table’ or equivalent award first B1	

Question Number	Scheme	Marks								
Q6 (a)	<table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td style="padding: 2px;">0</td> <td style="padding: 2px;">1</td> <td style="padding: 2px;">2</td> <td style="padding: 2px;">3</td> </tr> <tr> <td style="padding: 2px;"><math>3a</math></td> <td style="padding: 2px;"><math>2a</math></td> <td style="padding: 2px;"><math>a</math></td> <td style="padding: 2px;"><math>b</math></td> </tr> </table>	0	1	2	3	$3a$	$2a$	$a$	$b$	B1 (1)
0	1	2	3							
$3a$	$2a$	$a$	$b$							
(b)	$3a + 2a + a + b = 1$ $2a + 2a + 3b = 1.6$ $14a = 1.4$ $a = 0.1$ $b = 0.4$	or equivalent, using Sum of probabilities =1 or equivalent, using $E(X)=1.6$ Attempt to solve cao cao M1 M1 M1dep B1 B1 (5)								
(c)	$P(0.5 < x < 3) = P(1) + P(2)$ $= 0.2 + 0.1$ $= 0.3$	$3a$ or their $2a$ +their $a$ M1 Require $0 < 3a < 1$ to award follow through A1 ft (2)								
(d)	$E(3X - 2) = 3E(X) - 2$ $= 3 \times 1.6 - 2$ $= 2.8$	M1 cao A1 (2)								
(e)	$E(X^2) = 1 \times 0.2 + 4 \times 0.1 + 9 \times 0.4 (= 4.2)$ $\text{Var}(X) = "4.2" - 1.6^2$ $= 1.64$	**given answer** M1 M1 A1 (3)								
(f)	$\text{Var}(3X - 2) = 9 \text{Var}(X)$ $= 14.76$	awrt 14.8 M1 A1 (2)								
<b>[15]</b>										
Notes	<p>6(a) Condone <math>a</math> clearly stated in text but not put in table.</p> <p>6(b) Must be attempting to solve 2 different equations so third M dependent upon first two Ms being awarded.                      Correct answers seen with no working B1B1 only, 2/5                      Correctly verified values can be awarded M1 for correctly verifying sum of probabilities =1, M1 for using <math>E(X)=1.6</math> M0 as no attempt to solve and B1B1 if answers correct.</p> <p>6(d) 2.8 only award M1A1</p> <p>6(e) Award first M for at least two non-zero terms correct. Allow first M for correct expression with <math>a</math> and <math>b</math> e.g. <math>E(X^2) = 6a+9b</math>                      Given answer so award final A1 for correct solution.</p> <p>6(f) 14.76 only award M1A1</p>									

Question Number	Scheme	Marks
Q7(a) (i)  (ii)  (b)  (c)  (d)	$P(A \cup B) = a + b$ $P(A \cup B) = a + b - ab$ $P(R \cup Q) = 0.15 + 0.35 = 0.5$ $P(R \cap Q) = P(R Q) \times P(Q) = 0.1 \times 0.35 = 0.035$ $P(R \cup Q) = P(R) + P(Q) - P(R \cap Q) \quad \text{OR} \quad P(R) = P(R \cap Q') + P(R \cap Q)$ $0.5 = P(R) + 0.35 - 0.035$ $P(R) = 0.185$ $P(R) = 0.15 + \text{their (c)} = 0.15 + 0.035 = 0.185$	cao B1  or equivalent B1 (2)  0.5 B1 (1)  0.035 M1 A1 (2)  M1  0.185 A1 (2) <b>[7]</b>
Notes	7(a) (i) Accept $a + b - 0$ for B1 <b>Special Case</b> If answers to (i) and (ii) are (i) $P(A)+P(B)$ and (ii) $P(A)+P(B)-P(A)P(B)$ award B0B1 7(a)(i) and (ii) answers must be clearly labelled or in correct order for marks to be awarded.	

Question Number	Scheme	Marks
Q8 (a)	<p>Let the random variable <math>X</math> be the lifetime in hours of bulb</p> $P(X < 830) = P\left(Z < \frac{\pm(830 - 850)}{50}\right)$ <p style="text-align: right;">Standardising with 850 and 50</p> $= P(Z < -0.4)$ $= 1 - P(Z < 0.4)$ <p style="text-align: right;">Using 1-(probability&gt;0.5)</p> $= 1 - 0.6554$ $= 0.3446 \text{ or } 0.344578 \text{ by calculator}$ <p style="text-align: right;">awrt 0.345</p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>(3)</p>
(b)	$0.3446 \times 500$ $= 172.3$ <p style="text-align: right;">Their (a) x 500</p> <p style="text-align: right;">Accept 172.3 or 172 or 173</p>	<p>M1</p> <p>A1</p> <p>(2)</p>
(c)	<p>Standardise with 860 and <math>\sigma</math> and equate to <math>z</math> value <math>\frac{\pm(818 - 860)}{\sigma} = z</math> value</p> $\frac{818 - 860}{\sigma} = -0.84(16) \text{ or } \frac{860 - 818}{\sigma} = 0.84(16) \text{ or } \frac{902 - 860}{\sigma} = 0.84(16) \text{ or equiv.}$ <p style="text-align: right;"><math>\pm 0.8416(2)</math></p> $\sigma = 49.9$ <p style="text-align: right;">50 or awrt 49.9</p>	<p>M1</p> <p>A1</p> <p>B1</p> <p>A1</p> <p>(4)</p>
(d)	<p>Company <math>Y</math> as the <u>mean</u> is greater for <math>Y</math>.</p> <p>They have (approximately) the same <u>standard deviation</u> or <u>sd</u></p> <p style="text-align: right;">both</p>	<p>B1</p> <p>B1</p> <p>(2)</p> <p>[11]</p>
Notes	<p>8(a) If 1-<math>z</math> used e.g. 1-0.4=0.6 then award second M0</p> <p>8(c) M1 can be implied by correct line 2</p> <p>A1 for completely correct statement or equivalent.</p> <p>Award B1 if 0.8416(2) seen</p> <p>Do not award final A1 if any errors in solution e.g. negative sign lost.</p> <p>8(d) Must use statistical terms as underlined.</p>	

# Mark Scheme (Results) January 2010

GCE

## Statistics S1 (6683)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>


January 2010


Publications Code UA023026

All the material in this publication is copyright

© Edexcel Ltd 2010

January 2010  
6683 Statistics S1  
Mark Scheme

Question Number	Scheme	Marks
Q1 (a)		<p>M1 A1 A1 (3)</p>
(b)	<p>P(Blue bead and a green bead) = <math>\left(\frac{1}{4} \times \frac{1}{3}\right) + \left(\frac{1}{4} \times \frac{1}{3}\right) = \frac{1}{6}</math> (or any exact equivalent)</p>	<p>M1 A1 (2) Total [5]</p>
Q1 (a)	<p>M1 for shape and labels: 3 branches followed by 3,2,2 with some <i>R</i>, <i>B</i> and <i>G</i> seen Allow 3 branches followed by 3, 3, 3 if 0 probabilities are seen implying that 3, 2, 2 intended Allow blank branches if the other probabilities imply probability on blanks is zero Ignore further sets of branches</p> <p>1<sup>st</sup> A1 for correct probabilities and correct labels on 1<sup>st</sup> set of branches. 2<sup>nd</sup> A1 for correct probabilities and correct labels on 2<sup>nd</sup> set of branches. (accept 0.33, 0.67 etc or better here)</p>	
Special Case	<p>(b) M1 for identifying the 2 cases <i>BG</i> and <i>GB</i> and adding 2 products of probabilities. These cases may be identified by their probabilities e.g. <math>\left(\frac{1}{4} \times \frac{1}{3}\right) + \left(\frac{1}{4} \times \frac{1}{3}\right)</math> NB <math>\frac{1}{6}</math> (or exact equivalent) with no working scores 2/2</p> <p><u>With Replacement</u> (This oversimplifies so do not apply Mis-Read: max mark 2/5)</p> <p>(a) B1 for 3 branches followed by 3, 3, 3 with correct labels and probabilities of <math>\frac{1}{2}, \frac{1}{4}, \frac{1}{4}</math> on each.</p> <p>(b) M1 for identifying 2, possibly correct cases and adding 2 products of probabilities but A0 for wrong answer <math>\left[\left(\frac{1}{4} \times \frac{1}{4}\right) + \left(\frac{1}{4} \times \frac{1}{4}\right)\right]</math> will be sufficient for M1A0 here but <math>\frac{1}{4} \times \frac{1}{2} + \dots</math> would score M0</p>	

Question Number	Scheme	Marks
Q2 (a)	Median is 33	B1 (1)
(b)	$Q_1 = 24, Q_3 = 40, \text{IQR} = 16$	B1 B1 B1ft (3)
(c)	$Q_1 - \text{IQR} = 24 - 16 = 8$ So 7 is only outlier	M1 A1ft (2)
(d)	 <p style="text-align: right;">(accept either whisker)</p>	B1ft B1 B1ft (3)  Total [9]
Q2 (b)	<p>1<sup>st</sup> B1 for <math>Q_1 = 24</math> and 2<sup>nd</sup> B1 for <math>Q_3 = 40</math>                      3<sup>rd</sup> B1ft for their IQR based on their lower and upper quartile.                      Calculation of range (<math>40 - 7 = 33</math>) is B0B0B0  <u>Answer only</u> of IQR = 16 scores 3/3. For any other answer we must see working in (b) or on stem and leaf diagram</p> <p>(c) M1 for evidence that <math>Q_1 - \text{IQR}</math> has been attempted, their “8” (&gt;7) seen or clearly attempted is sufficient                      A1 ft must have seen their “8” and a suitable comment that only one person scored below this.</p> <p>(d) 1<sup>st</sup> B1ft for a clear box shape and ft their <math>Q_1, Q_2</math> and <math>Q_3</math> readable off the scale.                      Allow this mark for a box shape even if <math>Q_3 = 40, Q_1 = 7</math> and <math>Q_2 = 33</math> are used                      2<sup>nd</sup> B1 for only one outlier appropriately marked at 7                      3<sup>rd</sup> B1ft for either lower whisker. If they choose the whisker to their lower limit for outliers then follow through their “8”.                      (There should be no upper whisker unless their <math>Q_3 &lt; 40</math>, in which case there should be a whisker to 40)</p> <p>A typical error in (d) is to draw the lower whisker to 7, this can only score B1B0B0</p>	

Question Number	Scheme	Marks
Q3 (a)	2.75 or $2\frac{3}{4}$ , 5.5 or 5.50 or $5\frac{1}{2}$	B1 B1 (2)
(b)	Mean birth weight = $\frac{4841}{1500} = 3.227\bar{3}$ <b>awrt 3.23</b>	M1 A1 (2)
(c)	Standard deviation = $\sqrt{\frac{15889.5}{1500} - \left(\frac{4841}{1500}\right)^2} = 0.421093\dots$ or $s = 0.4212337\dots$	M1 A1ft A1 (3)
(d)	$Q_2 = 3.00 + \frac{403}{820} \times 0.5 = 3.2457\dots$ (allow 403.5..... $\rightarrow$ 3.25)	M1 A1 (2)
(e)	Mean(3.23) < Median(3.25) (or very close) Negative Skew (or symmetrical)	B1ft dB1ft (2) <b>Total [11]</b>
Q3 (b)	<p>M1 for a correct expression for mean. Answer only scores both.</p> <p>(c) M1 for a correct expression (ft their mean) for sd or variance. Condone mis-labelling eg sd=... with no square root or no labelling                      1<sup>st</sup> A1ft for a correct expression (ft their mean) including square root and no mis-labelling                      Allow 1<sup>st</sup> A1 for <math>\sigma^2 = 0.177\dots \rightarrow \sigma = 0.42\dots</math>                      2<sup>nd</sup> A1 for awrt 0.421. Answer only scores 3/3</p> <p>(d) M1 for a correct expression (allow 403.5 i.e. use of <math>n + 1</math>) but must have 3.00, 820 and 0.5                      A1 for awrt 3.25 provided M1 is scored.                      NB 3.25 with no working scores 0/2 as some candidates think mode is 3.25.</p> <p>(e) 1<sup>st</sup> B1ft for a comparison of their mean and median (may be in a formula but if <math>\pm(\text{mean} - \text{median})</math> is calculated that's OK. We are not checking the <u>value</u> but the <u>sign</u> must be consistent.)                      Also allow for use of quartiles <u>provided correct values seen</u>: <math>Q_1 = 3.02, Q_3 = 3.47</math>                      [They should get <math>(0.22 =) Q_3 - Q_2 &lt; Q_2 - Q_1 (= 0.23)</math> and say (slight) negative skew or symmetric]                      2<sup>nd</sup> dB1ft for a compatible comment based on their comparison. Dependent upon a suitable, correct comparison. Mention of "correlation" rather than "skewness" loses this mark.</p>	

Question Number	Scheme	Marks
<p>4 (a)</p> <div data-bbox="225 264 903 875" style="border: 1px solid black; padding: 10px; margin: 10px auto; width: fit-content;"> </div>	<p>3 closed curves and 4 in centre Evidence of subtraction</p> <p>31,36,24 41,17,11 Labels on loops, 16 and box</p>	<p>M1 M1 A1 A1 B1</p> <p>(5)</p> <p>B1ft (1)</p> <p>B1ft (1)</p> <p>M1 A1 (2)</p> <p><b>Total [9]</b></p>
<p>4 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>2<sup>nd</sup> M1 There may be evidence of subtraction in “outer” portions, so with 4 in the centre then 35, 40 28 (instead of 31,36,24) along with 33, 9, 3 can score this mark but A0A0 N.B. This is a common error and their “16” becomes 28 but still scores B0 in part (a)</p> <p>B1ft for <math>\frac{16}{180}</math> or any exact equivalent. Can fit their “16” from their box. If there is no value for their “16” in the box only allow this mark if they have <u>shown</u> some working.</p> <p>B1ft ft their “17”. Accept any exact equivalent</p> <p>If a probability greater than 1 is found in part (d) score M0A0</p> <p>M1 for clear sight of <math>\frac{P(S \cap D \cap N)}{P(S \cap N)}</math> and an attempt at one of the probabilities, ft their values.</p> <p>Allow <math>P(\text{all 3}   S \cap N) = \frac{4}{36}</math> or <math>\frac{1}{9}</math> to score M1 A0.</p> <p>Allow a correct ft from their diagram to score M1A0 e.g. in 33,3,9 case in (a): <math>\frac{4}{44}</math> or <math>\frac{1}{11}</math> is M1A0 A ratio of probabilities with a <u>product</u> of probabilities on top is M0, even with a correct formula.</p> <p>A1 for <math>\frac{4}{40}</math> or <math>\frac{1}{10}</math> or an exact equivalent</p> <p>Allow <math>\frac{4}{40}</math> or <math>\frac{1}{10}</math> to score both marks if this follows from their diagram, otherwise some explanation (method) is required.</p>	

Question Number	Scheme	Marks
Q5 (a)	$k + 4k + 9k = 1$ $14k = 1$ $k = \frac{1}{14} \quad \text{**given**}$	M1 A1 (2)
(b)	$P(X \geq 2) = 1 - P(X = 1) \quad \text{or} \quad P(X = 2) + P(X = 3)$ $= 1 - k = \frac{13}{14} \quad \text{or} \quad 0.92857\dots$	M1 A1 awrt 0.929 (2)
(c)	$E(X) = 1 \times k + 2 \times k \times 4 + 3 \times k \times 9 \quad \text{or} \quad 36k$ $= \frac{36}{14} = \frac{18}{7} \quad \text{or} \quad 2\frac{4}{7} \quad \text{(or exact equivalent)}$	M1 A1 (2)
(d)	$\text{Var}(X) = 1 \times k + 4 \times k \times 4 + 9 \times k \times 9 - \left(\frac{18}{7}\right)^2$ $\text{Var}(1 - X) = \text{Var}(X)$ $= \frac{19}{49} \quad \text{or} \quad 0.387755\dots$	M1 M1 M1 awrt 0.388 A1 (4) Total [10]
Q5 (a)	M1 for clear attempt to use $\sum p(x) = 1$ , full expression needed and the “1” must be clearly seen. This may be seen in a table. A1cso for no incorrect working seen. The sum and “= 1” must be explicitly seen somewhere.  A verification approach to (a) must show addition for M1 and have a suitable comment e.g. “therefore $k = \frac{1}{14}$ ” for A1 cso	
(b)	M1 for $1 - P(X \leq 1)$ or $P(X = 2) + P(X = 3)$ A1 for awrt 0.929. Answer only scores 2/2	
(c)	M1 for a full expression for $E(X)$ with at least two terms correct. NB If there is evidence of division (usually by 3) then score M0 A1 for any exact equivalent - answer only scores 2/2	
(d)	1 <sup>st</sup> M1 for clear attempt at $E(X^2)$ , need at least 2 terms correct in $1 \times k + 4 \times 4k + 9 \times 9k$ or $E(X^2) = 7$ 2 <sup>nd</sup> M1 for their $E(X^2) - (\text{their } \mu)^2$ 3 <sup>rd</sup> M1 for clearly stating that $\text{Var}(1 - X) = \text{Var}(X)$ , wherever seen A1 accept awrt 0.388. All 3 M marks are required. Allow 4/4 for correct answer only but must be for $\text{Var}(1 - X)$ .	


Question Number	Scheme	Marks
<p>Q6 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>+</p> <p>(f)</p> <p>(f)</p> <p>(e)</p> <p>(g)</p>	$S_{pp} = 106397 - \frac{833^2}{7} = 7270$ $S_{tp} = 42948 - \frac{341 \times 833}{7} = 2369, \quad S_{tt} = 18181 - \frac{341^2}{7} = 1569.42857... \quad \text{or} \quad \frac{10986}{7}$ $r = \frac{2369}{\sqrt{7270 \times 1569.42857..}} = 0.7013375$ <p>(Pmcc shows positive correlation.) Older patients have higher blood pressure</p> <p>(d) Points plotted correctly on graph: -1 each error or omission (within one square of correct position) * see diagram below for correct points</p> <p>(f) Line drawn with correct intercept, and gradient</p> $b = \frac{2369}{1569.42857..} = 1.509466...$ $a = \frac{833}{7} - b \times \frac{341}{7} = 45.467413...$ $p = 45.5 + 1.51t$ <p><math>t = 40, p = 105.84...</math> from equation or graph.</p>	<p>M1 A1</p> <p>A1 A1 (4)</p> <p>M1 A1ft</p> <p>A1 (3)</p> <p>B1 (1)</p> <p>B2</p> <p>B1ft B1 (2+2)</p> <p>M1 A1</p> <p>M1</p> <p>A1 (4)</p> <p>M1 A1 (2)</p> <p><b>Total [18]</b></p>
<p>Q6 (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p> <p>(g)</p>	<p>M1 for at least one correct expression 1<sup>st</sup> A1 for <math>S_{pp} = 7270</math>, 2<sup>nd</sup> A1 for <math>S_{tp} = 2369</math> or 2370, 3<sup>rd</sup> A1 for <math>S_{tt} =</math> awrt 1570</p> <p>M1 for attempt at correct formula and at least one correct value (or correct ft) M0 for <math>\frac{42948}{\sqrt{106397 \times 18181}}</math></p> <p>A1ft All values correct or correct ft. Allow for an answer of 0.7 or 0.70 <u>Answer only:</u> awrt 0.701 is 3/3, answer of 0.7 or 0.70 is 2/3</p> <p>B1 for comment in context that <u>interprets</u> the fact that correlation is positive, as in scheme. Must mention age and blood pressure in words, not just “t” and “p”.</p> <p>Record 1 point incorrect as B1B0 on open. [NB overlay for (60, 135) is slightly wrong]</p> <p>1<sup>st</sup> M1 for use of the correct formula for <math>b</math>, ft their values from (a) 1<sup>st</sup> A1 allow 1.5 or better 2<sup>nd</sup> M1 for use of <math>\bar{y} - b\bar{x}</math> with their values 2<sup>nd</sup> A1 for full equation with <math>a =</math> awrt 45.5 and <math>b =</math> awrt 1.51. Must be <math>p</math> in terms of <math>t</math>, not <math>x</math> and <math>y</math>.</p> <p>1<sup>st</sup> B1ft ft their intercept (within one square). You may have to extend their line. 2<sup>nd</sup> B1 for correct gradient i.e. parallel to given line (Allow 1 square out when <math>t = 80</math>)</p> <p>M1 for clear use of their equation with <math>t = 40</math> or correct value from their graph. A1 for awrt 106. Correct answer only (2/2) otherwise look for evidence on graph to award M1</p>	


Question Number

Scheme

Diagram for Q6 (d) + (f)

Q6 (d) + (f)



Question Number	Scheme	Marks
<p>Q7 (a)</p>  <p>(b) <math>P(X &lt; 154) = 0.05</math>  <math>\frac{154 - \mu}{\sigma} = -1.6449</math> or <math>\frac{\mu - 154}{\sigma} = 1.6449</math>  <math>\mu = 154 + 1.6449\sigma</math> <b>**given**</b></p> <p>(c) <math>172 - \mu = 0.5244\sigma</math> or <math>\frac{172 - \mu}{\sigma} = 0.5244</math> (allow <math>z = 0.52</math> or better here but must be in an equation)  Solving gives <math>\sigma = 8.2976075</math> (<b>awrt 8.30</b>) and <math>\mu = 167.64873</math> (<b>awrt 168</b>)</p> <p>(d) <math>P(\text{Taller than 160cm}) = P\left(Z &gt; \frac{160 - \mu}{\sigma}\right)</math>  <math>= P(Z &lt; 0.9217994)</math>  <math>= 0.8212</math> <b>awrt 0.82</b></p>	<p>bell shaped, must have inflexions</p> <p>154,172 on axis</p> <p>5% and 30%</p> <p>M1 B1 A1 cso</p> <p>B1</p> <p>M1 A1 A1</p> <p>M1 B1 A1</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>(3)</p> <p>(3)</p> <p>(4)</p> <p>(3)</p> <p>(3)</p> <p><b>Total [13]</b></p>
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>2<sup>nd</sup> B1 for 154 and 172 marked but 154 must be <math>&lt; \mu</math> and <math>172 &gt; \mu</math>. But <math>\mu</math> need not be marked.  Allow for <math>\frac{154 - \mu}{\sigma}</math> and <math>\frac{172 - \mu}{\sigma}</math> marked on appropriate sides of the peak.</p> <p>3<sup>rd</sup> B1 the 5% and 30% should be clearly indicated in the correct regions i.e. LH tail and RH tail.</p> <p>M1 for <math>\pm \frac{(154 - \mu)}{\sigma} = z</math> value (<math>z</math> must be recognizable e.g. 1.64, 1.65, 1.96 but NOT 0.5199 etc)</p> <p>B1 for <math>\pm 1.6449</math> seen in a line before the final answer.</p> <p>A1cso for no incorrect statements (in <math>\mu</math>, <math>\sigma</math>) equating a <math>z</math> value and a probability or incorrect signs e.g. <math>\frac{154 - \mu}{\sigma} = 0.05</math> or <math>\frac{154 - \mu}{\sigma} = 1.6449</math> or <math>P(Z &lt; \frac{\mu - 154}{\sigma}) = 1.6449</math></p> <p>B1 for a correct 2<sup>nd</sup> equation (NB <math>172 - \mu = 0.525\sigma</math> is B0, since <math>z</math> is incorrect)</p> <p>M1 for solving their two linear equations leading to <math>\mu = \dots</math> or <math>\sigma = \dots</math></p> <p>1<sup>st</sup> A1 for <math>\sigma =</math> awrt 8.30, 2<sup>nd</sup> A1 for <math>\mu =</math> awrt 168 [NB the 168 can come from false working. These A marks require use of correct equation from (b), and a <math>z</math> value for “0.5244” in (c)]  NB use of <math>z = 0.52</math> will typically get <math>\sigma = 8.31</math> and <math>\mu = 167.67\dots</math> and score B1M1A0A1  <u>No working</u> and both correct scores 4/4, only one correct scores 0/4  Provided the M1 is scored the A1s can be scored even with B0 (e.g. for <math>z = 0.525</math>)</p> <p>M1 for attempt to standardise with 160, their <math>\mu</math> and their <math>\sigma (&gt; 0)</math>. Even allow with symbols <math>\mu</math> and <math>\sigma</math>.</p> <p>B1 for <math>z =</math> awrt <math>\pm 0.92</math>  <u>No working</u> and a correct answer can score 3/3 provided <math>\sigma</math> and <math>\mu</math> are correct to 2sf.</p>	



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@linneydirect.com](mailto:publications@linneydirect.com)

Order Code UA023026 January 2010

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH

# Mark Scheme (Results) Summer 2010

GCE

## Statistics S1 (6683)

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

Summer 2010

Publications Code UA024765

All the material in this publication is copyright

© Edexcel Ltd 2010

## General Marking Guidance

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - M marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - B marks are unconditional accuracy marks (independent of M marks)
3. Abbreviations


These are some of the marking abbreviations that will appear in the mark scheme

- ft - follow through
- awrt - answers which round to
- oe - or equivalent (and appropriate)
- isw - ignore subsequent working
- cao - correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- SC: special case



**June 2010**  
**Statistics S1 6683**  
**Mark Scheme**

Question Number	Scheme	Marks
Q1 (a)	$r = \frac{8825}{\sqrt{1022500 \times 130.9}}, \quad = \text{awrt } \underline{\mathbf{0.763}}$	M1 A1 (2)
(b)	Teams with high attendance scored more goals (oe, statement in context)	B1 (1)
(c)	0.76(3)	B1ft (1)
<b>Total 4</b>		
(a)	M1 for a correct expression, square root required Correct answer award 2/2	
(b)	Context required (attendance and goals). Condone causality. B0 for 'strong positive correlation between attendance and goals' on its own oe	
(c)	Value required. Must be a correlation coefficient between -1 and +1 inclusive. B1ft for 0.76 or better or same answer as their value from part (a) to at least 2 d.p.	

Question Number	Scheme	Marks
Q2 (a)	 <p style="text-align: center;">P(R) and P(B) 2<sup>nd</sup> set of probabilities</p>	<p>B1 B1 (2) M1 A1 (2) M1 A1ft A1 (3) M1 A1ft A1 (3) <b>Total 10</b></p>
	<p>(b) <math>P(H) = \frac{5}{12} \times \frac{2}{3} + \frac{7}{12} \times \frac{1}{2}, = \frac{41}{72}</math> or awrt 0.569</p> <p>(c) <math>P(R H) = \frac{\frac{5}{12} \times \frac{2}{3}}{\frac{41}{72}}, = \frac{20}{41}</math> or awrt 0.488</p> <p>(d) <math>\left(\frac{5}{12}\right)^2 + \left(\frac{7}{12}\right)^2</math>  <math>= \frac{25}{144} + \frac{49}{144} = \frac{74}{144}</math> or <math>\frac{37}{72}</math> or awrt 0.514</p>	<p>(a) 1<sup>st</sup> B1 for the probabilities on the first 2 branches. Accept 0.416 and 0.583                  2<sup>nd</sup> B1 for probabilities on the second set of branches. Accept 0.6, 0.3, 0.5 and <math>\frac{1.5}{3}</math>                  Allow exact decimal equivalents using clear recurring notation if required.</p> <p>(b) M1 for an expression for P(H) that follows through their sum of two products of <b>probabilities</b> from their tree diagram</p> <p>(c) M1 for <math>\frac{P(R \cap H)}{P(H)}</math> with denominator their (b) substituted e.g. <math>\frac{P(R \cap H)}{P(H)} = \frac{5}{12}</math> award M1.                  Formula seen                  Formula not seen M1 for <math>\frac{\text{probability} \times \text{probability}}{\text{their } b}</math> but M0 if fraction repeated e.g. <math>\frac{5}{12} \times \frac{2}{3}</math>.  <math>\frac{2}{3}</math></p> <p>1<sup>st</sup> A1ft for a fully correct expression or correct follow through                  2<sup>nd</sup> A1 for <math>\frac{20}{41}</math> o.e.</p> <p>(d) M1 for <math>\left(\frac{5}{12}\right)^2</math> or <math>\left(\frac{7}{12}\right)^2</math> can follow through their equivalent values from tree diagram                  1<sup>st</sup> A1 for both values correct or follow through from their original tree and +                  2<sup>nd</sup> A1 for a correct answer                  Special Case <math>\frac{5}{12} \times \frac{4}{11}</math> or <math>\frac{7}{12} \times \frac{6}{11}</math> seen award M1A0A0</p>

Question Number	Scheme	Marks
Q3 (a)	$2a + \frac{2}{5} + \frac{1}{10} = 1$ <p style="text-align: right;">(or equivalent)</p> $\underline{a = \frac{1}{4} \text{ or } 0.25}$	M1 A1 (2)
(b)	$E(X) = \underline{1}$	B1 (1)
(c)	$E(X^2) = 1 \times \frac{1}{5} + 1 \times \frac{1}{10} + 4 \times \frac{1}{4} + 9 \times \frac{1}{5}$ <p style="text-align: right;">(= 3.1)</p>	M1
(d)	$\text{Var}(X) = 3.1 - 1^2,$ <p style="text-align: right;"><math>= \underline{2.1 \text{ or } \frac{21}{10} \text{ oe}}</math></p>	M1 A1 (3)
(e)	$\text{Var}(Y) = (-2)^2 \text{Var}(X),$ <p style="text-align: right;"><math>= \underline{8.4 \text{ or } \frac{42}{5} \text{ oe}}</math></p>	M1 A1 (2)
(e)	$X \geq Y$ when $X = 3$ or $2,$ so probability = " $\frac{1}{4}$ " + " $\frac{1}{5}$ " <p style="text-align: right;"><math>= \underline{\frac{9}{20} \text{ oe}}</math></p>	M1 A1ft A1 (3)
<b>Total 11</b>		
(a)	M1 for a clear attempt to use $\sum P(X = x) = 1$ Correct answer only 2/2. <b>NB Division by 5 in parts (b), (c) and (d) seen scores 0. Do not apply ISW.</b>	
(b)	B1 for 1	
(c)	1 <sup>st</sup> M1 for attempting $\sum x^2 P(X = x)$ at least two terms correct. Can follow through. 2 <sup>nd</sup> M1 for attempting $E(X^2) - [E(X)]^2$ or allow subtracting 1 from their attempt at $E(X^2)$ provided no incorrect formula seen. Correct answer only 3/3.	
(d)	M1 for $(-2)^2 \text{Var}(X)$ or $4\text{Var}(X)$ Condone missing brackets provided final answer correct for their $\text{Var}(X)$ . Correct answer only 2/2.	
(e)	Allow M1 for distribution of $Y = 6 - 2X$ and correct attempt at $E(Y^2) - [E(Y)]^2$ M1 for identifying $X = 2, 3$ 1 <sup>st</sup> A1ft for attempting to find their $P(X=2) + P(X=3)$ 2 <sup>nd</sup> A1 for $\frac{9}{20}$ or 0.45	

Question Number	Scheme	Marks
Q4	<p>(a) <math>\frac{2+3}{\text{their total}} = \frac{5}{\text{their total}} = \frac{1}{6}</math> (** given answer**)</p> <p>(b) <math>\frac{4+2+5+3}{\text{total}}, = \frac{14}{30}</math> or <math>\frac{7}{15}</math> or 0.46</p> <p>(c) <math>P(A \cap C) = 0</math></p> <p>(d) <math>P(C   \text{reads at least one magazine}) = \frac{6+3}{20} = \frac{9}{20}</math></p> <p>(e) <math>P(B) = \frac{10}{30} = \frac{1}{3}</math>, <math>P(C) = \frac{9}{30} = \frac{3}{10}</math>, <math>P(B \cap C) = \frac{3}{30} = \frac{1}{10}</math> or <math>P(B C) = \frac{3}{9}</math></p> <p><math>P(B) \times P(C) = \frac{1}{3} \times \frac{3}{10} = \frac{1}{10} = P(B \cap C)</math> or <math>P(B C) = \frac{3}{9} = \frac{1}{3} = P(B)</math></p> <p>So yes they are statistically independent</p>	<p>M1 A1cso (2)</p> <p>M1 A1 (2)</p> <p>B1 (1)</p> <p>M1 A1 (2)</p> <p>M1</p> <p>M1</p> <p>A1cso (3)</p> <p><b>Total 10</b></p>
	<p>(a) M1 for <math>\frac{2+3}{\text{their total}}</math> or <math>\frac{5}{30}</math></p> <p>(b) M1 for adding at least 3 of “4, 2, 5, 3” and dividing by their total to give a probability Can be written as separate fractions substituted into the completely correct Addition Rule</p> <p>(c) B1 for 0 or 0/30</p> <p>(d) M1 for a <b>denominator of 20</b> or <math>\frac{20}{30}</math> leading to an answer with denominator of 20 <math>\frac{9}{20}</math> only, 2/2</p> <p>(e) 1<sup>st</sup> M1 for attempting all the required probabilities for a suitable test 2<sup>nd</sup> M1 for use of a correct test - must have attempted all the correct probabilities. Equality can be implied in line 2. A1 for fully correct test carried out with a comment</p>	

Question Number	Scheme	Marks
Q5	<p>(a) 23, 35.5 (may be in the table)</p> <p>(b) Width of 10 units is 4 cm so width of 5 units is <b>2 cm</b> Height = <math>2.6 \times 4 = \mathbf{10.4 \text{ cm}}</math></p> <p>(c) <math>\sum fx = 1316.5 \Rightarrow \bar{x} = \frac{1316.5}{56} =</math> awrt <b>23.5</b> <math>\sum fx^2 = 37378.25</math> can be implied So <math>\sigma = \sqrt{\frac{37378.25}{56} - \bar{x}^2} =</math> awrt <b>10.7</b> allow <math>s = 10.8</math></p> <p>(d) <math>Q_2 = (20.5) + \frac{(28 - 21)}{11} \times 5 = 23.68\dots</math> awrt <b>23.7 or 23.9</b></p> <p>(e) <math>Q_3 - Q_2 = 5.6, Q_2 - Q_1 = 7.9</math> (or <math>\bar{x} &lt; Q_2</math>) [7.9 &gt; 5.6 so ] <b>negative skew</b></p>	<p>B1 B1 (2)</p> <p>B1</p> <p>M1 A1 (3)</p> <p>M1 A1</p> <p>B1</p> <p>M1 A1 (5)</p> <p>M1 A1 (2)</p> <p>M1</p> <p>A1 (2)</p> <p><b>Total 14</b></p>
	<p>(b) M1 for their width x their height=20.8. Without labels assume width first, height second and award marks accordingly.</p> <p>(c) 1<sup>st</sup> M1 for reasonable attempt at <math>\sum x</math> and /56 2<sup>nd</sup> M1 for a method for <math>\sigma</math> or <math>s</math>, <math>\sqrt{\quad}</math> is required Typical errors <math>\sum (fx)^2 = 354806.3</math> M0, <math>\sum f^2x = 13922.5</math> M0 and <math>(\sum fx)^2 = 1733172</math> M0 Correct answers only, award full marks.</p> <p>(d) Use of <math>\sum f(x - \bar{x})^2 =</math> awrt 6428.75 for B1 lcb can be 20, 20.5 or 21, width can be 4 or 5 and the fraction part of the formula correct for M1 - Allow 28.5 in fraction that gives awrt 23.9 for M1A1</p> <p>(e) M1 for attempting a test for skewness using quartiles or mean and median. Provided median greater than 22.55 and less than 29.3 award for M1 for <math>Q_3 - Q_2 &lt; Q_2 - Q_1</math> without values as a valid reason. SC Accept mean close to median and no skew oe for M1A1</p>	

Question Number	Scheme	Marks
Q6	<p>(a) See overlay</p> <p>(b) The <b>points</b> lie reasonably close to a straight <b>line</b> (o.e.)</p> <p>(c) <math>\sum d = 27.7, \quad \sum f = 146</math> (both, may be implied)</p> $S_{dd} = 152.09 - \frac{(27.7)^2}{6} = 24.208.....$ <p style="text-align: right;"><b>awrt <u>24.2</u></b></p> $S_{fd} = 723.1 - \frac{27.7 \times 146}{6} = 49.06....$ <p style="text-align: right;"><b>awrt <u>49.1</u></b></p> <p>(d) <math>b = \frac{S_{fd}}{S_{dd}} = 2.026....</math> <b>awrt <u>2.03</u></b></p> $a = \frac{146}{6} - b \times \frac{27.7}{6} = 14.97..... \quad \text{so } \underline{f = 15.0 + 2.03d}$ <p>(e) A flight costs <b>£2.03 (or about £2)</b> for every extra <b>100km</b> or about <b>2p</b> per <b>km</b>.</p> <p>(f) <math>15.0 + 2.03d &lt; 5d</math> so <math>d &gt; \frac{15.0}{(5 - 2.03)} = 5.00 \sim 5.05</math></p> <p>So <math>t &gt; 500 \sim 505</math></p>	<p>B1 B1 (2)</p> <p>B1 (1)</p> <p>B1</p> <p>M1 A1</p> <p>A1 (4)</p> <p>M1 A1</p> <p>M1 A1 (4)</p> <p>B1ft (1)</p> <p>M1</p> <p>A1 (2)</p> <p style="text-align: right;"><b>Total 14</b></p>
	<p>(a) 1<sup>st</sup> B1 for at least 4 points correct (allow <math>\pm</math> one 2mm square) 2<sup>nd</sup> B1 for all points correct (allow <math>\pm</math> one 2 mm square)</p> <p>(b) Ignore extra points and lines Require reference to points and line for B1.</p> <p>(c) M1 for a correct method seen for either - a correct expression 1<sup>st</sup> A1 for <math>S_{dd}</math> awrt 24.2 2<sup>nd</sup> A1 for <math>S_{fd}</math> awrt 49.1</p> <p>(d) 1<sup>st</sup> M1 for a correct expression for <math>b</math> - can follow through their answers from (c) 2<sup>nd</sup> M1 for a correct method to find <math>a</math> - follow through their <math>b</math> and their means 2<sup>nd</sup> A1 for <math>f = \dots</math> in terms of <math>d</math> and all values awrt given expressions. Accept 15 as rounding from correct answer only.</p> <p>(e) Context of cost and distance required. Follow through their value of <math>b</math></p> <p>(f) M1 for an attempt to find the intersection of the 2 lines. Value of <math>t</math> in range 500 to 505 seen award M1. Value of <math>d</math> in range 5 to 5.05 award M1. Accept <math>t</math> greater than 500 to 505 inclusive to include graphical solution for M 1A1</p>	

Question Number	Scheme	Marks
Q7 (a)	$P(D > 20) = P\left(Z > \frac{20-30}{8}\right)$ $= P(Z > -1.25)$ $= \underline{\underline{0.8944}} \qquad \qquad \qquad \underline{\underline{\text{awrt } 0.894}}$ (b) $P(D < Q_3) = 0.75$ so $\frac{Q_3 - 30}{8} = 0.67$ $Q_3 = \text{awrt } \underline{\underline{35.4}}$ (c) $35.4 - 30 = 5.4$ so $Q_1 = 30 - 5.4 = \text{awrt } \underline{\underline{24.6}}$ (d) $Q_3 - Q_1 = 10.8$ so $1.5(Q_3 - Q_1) = 16.2$ so $Q_1 - 16.2 = h$ or $Q_3 + 16.2 = k$ $h = \underline{\underline{8.4 \text{ to } 8.6}}$ and $k = \underline{\underline{51.4 \text{ to } 51.6}}$ both	M1 A1 A1 (3)  M1 B1  A1 (3)  B1ft (1)  M1  A1 (2)  M1  M1 A1 (3)  Total 12
	(a) M1 for an attempt to standardise 20 or 40 using 30 and 8. 1 <sup>st</sup> A1 for $z = \pm 1.25$ 2 <sup>nd</sup> A1 for awrt 0.894	
	(b) M1 for $\frac{Q_3 - 30}{8} =$ to a $z$ value M0 for 0.7734 on RHS. B1 for ( $z$ value) between 0.67~0.675 seen. M1B0A1 for use of $z = 0.68$ in correct expression with awrt 35.4	
	(c) Follow through using their of quartile values.	
	(d) M1 for an attempt to calculate 1.5(IQR) and attempt to add or subtract using one of the formulae given in the question - follow through their quartiles	
	(e) 1 <sup>st</sup> M1 for attempting $2P(D > \text{their } k)$ or $(P(D > \text{their } k) + P(D < \text{their } h))$ 2 <sup>nd</sup> M1 for standardising their $h$ or $k$ (may have missed the 2) so allow for standardising $P(D > 51.6)$ or $P(D < 8.4)$ Require boths Ms to award A mark.	





Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@linneydirect.com](mailto:publications@linneydirect.com)

Order Code UA024765 Summer 2010

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH

# Mark Scheme (Results) January 2011

GCE

## GCE Statistics S1 (6683) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our [Ask The Expert](#) email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

January 2011

Publications Code UA026664

All the material in this publication is copyright

© Edexcel Ltd 2011

## General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

### 3. Abbreviations


These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod - benefit of doubt
- ft - follow through
- the symbol  $\surd$  will be used for correct ft
- cao - correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw - ignore subsequent working
- awrt - answers which round to
- SC: special case
- oe - or equivalent (and appropriate)
- dep - dependent
- indep - independent
- dp decimal places
- sf significant figures
- \* The answer is printed on the paper
- The second mark is dependent on gaining the first mark

**January 2011  
Statistics S1 6683  
Mark Scheme**

Question Number	Scheme	Marks
1.		
(a)	$S_{ll} = 327754.5 - \frac{4027^2}{50} = 3419.92$ $S_{lw} = 29330.5 - \frac{357.1 \times 4027}{50} = 569.666$	M1 A1 A1 (3)
(b)	$r = \frac{569.666}{\sqrt{3419.92 \times 289.6}} = 0.572$ awrt 0.572 or 0.573	M1 A1 (2)
(c)	As the length of the salmon increases the weight increases	B1ft (1) <b>[6]</b>
<b>Notes</b>		
(a)	M1 for at least one correct expression 1 <sup>st</sup> A1 for $S_{ll} =$ awrt 3420 (Condone $S_{xx} = \dots$ or even $S_{yy} = \dots$ ) 2 <sup>nd</sup> A1 for $S_{lw} =$ awrt 570 (Condone $S_{xy} = \dots$ )	
(b)	M1 for attempt at correct formula. Must have their $S_{ll}$ , $S_{lw}$ and given $S_{ww}$ in the correct places If $S_{ll}$ , $S_{lw}$ are correct and an answer of awrt 0.57 is seen then award M1A0 M0 for $\frac{29330.5}{\sqrt{327754.5 \times 289.6}}$	
(c)	B1ft for a comment mentioning “length” and “weight”, not just $l$ and $w$ , and the idea of longer salmon weighing more. e.g. “positive correlation between weight and length” is B0 since the idea of positive correlation is not explained. Allow “larger” instead of “heavier” or “longer” Ignore any spurious values mentioned such as 0.572 If their $r$ is negative (but must be $r > -1$ ) ft an appropriate comment. Condone $r > 1$ if comment is correct. If $ r  < 0.4$ allow a comment of no or little relationship between weight and length but for $0 < r < 0.4$ the printed answer is still acceptable too.  Treat mention of “skewness” as ISW if a correct interpretation is given	

Question Number	Scheme	Marks
2.  (a)	$2.8 + 5.6 + 2.3 + 9.4 + 0.5 + 1.8 + 84.6 = 107$ mean = $107 / 28 (= 3.821\dots)$	M1 A1 (2)
(b)	It will have no effect since one is 4.5 under what it should be and the other is 4.5 above what it should be.	B1 dB1 (2) [4]
<b>Notes</b>		
(a)	M1 for a clear attempt to add the two sums. Accept a full expression or $2.8 + 5.6 + \dots + 84.6 = x$ where $100 < x < 110$ i.e. seeing at least two correct terms of Keith's and the 84.6 with a slip. A1 for awrt 3.8 (Condone 1 dp/2sf here since data is given to 1 dp or 2 sf) Accept $\frac{107}{28}$ or $3\frac{23}{28}$ or any exact equivalent <b>Correct answer implies M1A1</b>	
(b)	1 <sup>st</sup> B1 for clearly stating that it will have no effect. ("roughly the same" is B0 B0) 2 <sup>nd</sup> dB1 for a supporting reason that mentions the fact that the increase and decrease are the same and gives some numerical value(s) to support this. e.g. Sum of Keith's observations is still 22.4 ( or mean is still 3.2) <u>or</u> Sum is still 107 <u>or</u> $9.4 - 4.9 = 5 - 0.5$ (o.e.) This second B1 is dependent on their saying there is no effect so BOB1 is not possible.	

Question Number	Scheme	Marks
<p>3.</p> <p>(a) Outliers  <math>14 + 1.5 \times (14 - 7) = 24.5</math>  <math>7 - 1.5 \times (14 - 7) = -3.5</math></p> <p>Outlier 25                      either upper limit acceptable on diagram</p>  <p style="text-align: right;">Sales in £'000</p>	<p>M1 A1</p> <p>M1 A1ft B1</p> <p style="text-align: right;">(5)</p>	
<p>(b) Since <math>Q_3 - Q_2 &lt; Q_2 - Q_1</math>. Allow written explanation negatively skew</p>	<p>B1 dB1</p> <p style="text-align: right;">(2)</p>	
<p>(c) not true                      since the lower quartile is 7000 and therefore 75% above 7000 not 10000 or 10 is inside the box or any other sensible comment</p>	<p>B1 dB1</p> <p style="text-align: right;">(2) [9]</p>	
<b>Notes</b>		
<p>(a)</p>	<p><b>A fully correct box-plot (either version) with no supporting work scores 5/5. Otherwise read on</b></p> <p>1<sup>st</sup> M1 for at least one correct calculation seen                      1<sup>st</sup> A1 for 24.5 and -3.5 (or just negative noted) seen. May be read off the graph.                      If both values are seen but no calculation is given then M1A1, one value M1A0.                      2<sup>nd</sup> M1 for a box with an upper and a lower whisker(s) with at least 2 correct values (condone no median marked)                      2<sup>nd</sup> A1ft for 3, 7, 12, 14 and 20 or 24.5 in appropriate places and readable off their scale                      If <u>both</u> upper whiskers are seen A0                      Apply ft for their <u>whiskers</u> being compatible with their <u>outlier limits</u>                      e.g. if their lower limit is + 3.5 then a lower whisker ending at 4 or 3.5 is OK                      B1 for only one outlier appropriately marked at 25</p> <p style="text-align: center;"><b>Apply ± 0.5 square accuracy for diagram</b></p>	
<p>(b)</p>	<p>1<sup>st</sup> B1 for <math>Q_3 - Q_2 &lt; Q_2 - Q_1</math> statement or an equivalent statement in words                      Use of <math>Q_3 - Q_2 &lt; Q_2 - Q_1</math> does not require differences to be seen.                      2<sup>nd</sup> dB1 for “negative skew” dependent on suitable reason given above. “correlation” is B0                      “positive skew” with a supporting argument based on whiskers can score B1B1                      e.g. “right hand whisker is longer than LH one so positive skew”  <math>Q_3 - Q_2 &lt; Q_2 - Q_1</math> followed by “positive skew” is B1B0</p>	
<p>(c)</p>	<p>1<sup>st</sup> B1 for rejecting the company’s claim                      2<sup>nd</sup> dB1 for an appropriate supporting reason. Dependent on rejecting company’s claim.</p>	

Question Number	Scheme	Marks
4. (a)	$b = \frac{1.688}{5.753} = 0.293$ $a = 3.22 - 4.42 \times 0.293 = 1.9231\dots$ $p = 1.92 + 0.293v$	M1A1  M1 A1  (4)
(b)	$v = \frac{85 - 5}{10} = 8$ $p = 1.92 + 0.293 \times 8 = 4.3$	M1  A1  (awrt 4.3)  (2) [6]
<b>Notes</b>		
<b>Can ignore (a) and (b) labels here</b>		
(a)	1 <sup>st</sup> M1 for a correct expression for $b$ . $\frac{1.688}{1.168}$ is M0 1 <sup>st</sup> A1 for awrt 0.29  2 <sup>nd</sup> M1 for use of $a = \bar{p} - b\bar{v}$ follow through their value of $b$ (or even just the letter $b$ ) 2 <sup>nd</sup> A1 for a complete equation with $a =$ awrt 1.92 and $b =$ awrt 0.293 $y$ or $p = 1.92 + 0.293x$ is A0 Correct answer with no working is 4/4	
(b)	M1 for an attempt to find the value of $v$ when $x = 85$ ( at least 2 correct terms in $\pm \frac{85-5}{10}$ ) or for an attempt to find an equation for $p$ in terms of $x$ and using $x = 85$ Attempt at equation of $p$ in $x$ requires $p = 1.92 + 0.293 \frac{(x-5)}{10}$  A1 for awrt 4.3 (award when first seen and apply ISW) N.B. $p = 1.92 + 0.293 \times 85$ (o.e.) is M0A0	

Question Number	Scheme	Marks
5.		
(a)	Median = $32/2 = 16^{\text{th}}$ term (16.5) $\frac{x - 39.5}{49.5 - 39.5} = \frac{16 - 14}{25 - 14}$ or $x = 39.5 + \left(\frac{2}{11} \times 10\right)$ Median = 41.3 (use of $n + 1$ gives 41.8) <b>(awrt 41.3)</b>	M1 A1 (2)
(b)	Mean = $\frac{1414}{32} = 44.1875$ <b>(awrt 44.2)</b> Standard deviation = $\sqrt{\frac{69378}{32} - \left(\frac{1414}{32}\right)^2}$ $= 14.7$ (or $s = 14.9$ )	B1 M1 A1 (3)
(c)	mean > median therefore <u>positive skew</u>	B1ft B1ft (2) [7]
<b>Notes</b>		
(a)	M1 for an attempt to use interpolation to find the median. Condone use of 39 or 40 for 39.5 e.g. allow $39 + \frac{2}{11} \times 10$ (o.e.) or $40 + \frac{2}{11} \times 10$ (o.e.) to score M1A0 but must have the 10 A1 for awrt 41.3 (or awrt 41.8 if using $(n + 1)$ )	
(b)	B1 for awrt 44.2 M1 for a correct expression including square root. (Allow ft of their mean) A1 for awrt 14.7 (If using $s$ for awrt 14.9) You may see $\sum t = 1339 \rightarrow \bar{t} = 41.8$ and $\sum t^2 = 62928 \rightarrow \sigma 14.7$ or $s = 14.9$ this scores B0 for the mean but can score M1 for a correct st.dev expression and A1 for ans. <b>Correct answer only in (a) and (b) can score full marks but check <math>(n + 1)</math> case in (a)</b>	
(c)	1 <sup>st</sup> B1ft for a correct comparison of their mean and their median (may be in a formula) Calculating median – mean as negative is OK for this B1 but must say +ve skew for 2 <sup>nd</sup> B1 Only allow comparison to be $\approx 0$ if $ \text{mean} - \text{median}  \leq 0.5$ 2 <sup>nd</sup> B1ft for a correct description of skewness <u>based on their values of mean and median.</u> ft their values for mean and median not their previous calculation/comparison Must be compatible with their previous comparison (if they have one) “Positive skew” with no reason is B0B1 provided you can see their values that imply that. Description should be “positive” or “negative” or “no” skew or “symmetric” “Positive correlation” is B0	
Quartiles	1 <sup>st</sup> B1ft if $Q_1 = \text{awrt } 32$ <u>and</u> $Q_3 = \text{awrt } 49$ seen and a correct comparison made. ft $Q_2$ 2 <sup>nd</sup> B1ft if $Q_1 = \text{awrt } 32$ <u>or</u> $Q_3 = \text{awrt } 49$ seen and a correct description based on their quartiles and their comparison is made. (Should get “negative skew”)	

Question Number	Scheme	Marks																
6.																		
(a)	$k + 2k + 3k + 4k = 1$ or $10k = 1$ $k = 0.1$ (*) [allow verification with a comment e.g. "so $k = 0.1$ "]	B1cso (1)																
(b)	$E(X) = 1 \times 0.1 + 2 \times 0.2 + 3 \times 0.3 + 4 \times 0.4 = 3$	M1 A1 (2)																
(c)	$E(X^2) = 1 \times 0.1 + 4 \times 0.2 + 9 \times 0.3 + 16 \times 0.4 = 10$	M1 A1 (2)																
(d)	$\text{Var}(X) = 10 - 9 (= 1)$ $\text{Var}(2 - 5X) = 5^2 \text{Var}(X) = 25$	M1 M1 A1 (3)																
(e)	$P(1,3) + P(2,2) = 2 \times 0.1 \times 0.3 + 0.2 \times 0.2 = 0.1$ (*)	M1 A1cso (2)																
(f)	<table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td><math>X_1 + X_2</math></td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> <td>7</td> <td>8</td> </tr> <tr> <td><math>p</math></td> <td>0.01</td> <td>0.04</td> <td>0.1</td> <td><b>0.2</b></td> <td>0.25</td> <td>0.24</td> <td><b>0.16</b></td> </tr> </table>	$X_1 + X_2$	2	3	4	5	6	7	8	$p$	0.01	0.04	0.1	<b>0.2</b>	0.25	0.24	<b>0.16</b>	B1 B1 (2)
$X_1 + X_2$	2	3	4	5	6	7	8											
$p$	0.01	0.04	0.1	<b>0.2</b>	0.25	0.24	<b>0.16</b>											
(g)	$P(2) + P(3) = 0.05$	M1A1 (2) <b>[14]</b>																

Question Number	Scheme	Marks
	<b>Notes</b>	
(a)	B1 for a clear attempt to use sum of probabilities = 1. Must see previous line as well as $k = 0.1$ <b>A correct expression for <math>E(X)</math> or <math>E(X^2)</math> that is later divided by 4 scores M0</b>	
(b)	M1 for a completely correct expression. May be implied by correct answer of 3 or $30k$ A1 for 3 only.	
(c)	M1 for a completely correct expression. May be implied by correct answer of 10 or $100k$ A1 for 10 only. [ For $E(X^2) = 0.1 + 0.8 + 2.7 + 6.4 - 9 = 1$ scores M0A0 but accept this as $\text{Var}(X)$ in (d)]	
(d)	1 <sup>st</sup> M1 for using $\text{Var}(X) = E(X^2) - E(X)^2$ , f.t their values from (b) and (c) Allow this mark for $\text{Var}(X) = 10 - 9$ or better. May be implied if this is seen in (c). 2 <sup>nd</sup> M1 for $5^2 \text{Var}(X)$ or $25\text{Var}(X)$ can f.t. their $\text{Var}(X)$ . Allow $-5^2$ if it later becomes $+25$ A1 for 25 only. Dependent upon both Ms Forming distribution for $Y = 2 - 5X$ gets M1 for $E(Y^2) = 194$ then M1A1 for $194 - 169 = 25$	
(e)	M1 for correctly identifying (1, 3) <u>or</u> (3, 1) <u>and</u> (2, 2) as required cases ( $3k^2 + 4k^2$ or better) A1 cso for 0.1 only but must see evidence for M1	
(f)	1 <sup>st</sup> B1 for 0.2 correctly assigned. May be in table. 2 <sup>nd</sup> B1 for 0.16 correctly assigned. May be in table	
(g)	M1 for $P(2) + P(3)$ . May be implied by correct answer of 0.05 A1 for 0.05 only. <b>Correct answer only can score full marks in parts (b), (c), (f) and (g)</b>	

Question Number	Scheme	Marks
7. (a)		<p>both <math>\frac{2}{3}, \frac{1}{3}</math> B1</p> <p><math>\frac{4}{9}</math> B1</p> <p>both <math>\frac{3}{5}, \frac{2}{5}</math> B1</p> <p>all three of <math>\frac{4}{9}, \frac{4}{9}, \frac{5}{9}</math> B1</p> <p>(4)</p>
(b)	$P(A) = P(RR) + P(YY) = \frac{1}{2} \times \frac{2}{5} + \frac{1}{2} \times \frac{2}{5} = \frac{2}{5}$	<p>B1 for <math>\frac{1}{2} \times \frac{2}{5}</math> (oe) seen at least once</p> <p>B1 M1 A1 (3)</p>
(c)	$P(B) = P(RRR) + P(RYR) + P(YRR) + P(YYY)$ $\left(\frac{1}{2} \times \frac{2}{5} \times \frac{2}{3}\right) + \left(\frac{1}{2} \times \frac{3}{5} \times \frac{5}{9}\right) + \left(\frac{1}{2} \times \frac{3}{5} \times \frac{5}{9}\right) + \left(\frac{1}{2} \times \frac{2}{5} \times \frac{4}{9}\right) = \frac{5}{9} \quad (*)$	<p>M1 for at least 1 case of 3 balls identified. (Implied by 2<sup>nd</sup> M1)</p> <p>M1</p> <p>M1, A1cso (3)</p>
(d)	$P(A \cap B) = P(RRR) + P(YYY)$ $= \left(\frac{1}{2} \times \frac{2}{5} \times \frac{2}{3}\right) + \left(\frac{1}{2} \times \frac{2}{5} \times \frac{4}{9}\right) = \frac{2}{9} \quad (*)$	<p>M1 for identifying both cases and + probs. may be implied by correct expressions</p> <p>M1</p> <p>A1cso (2)</p>
(e)	$P(A \cup B) = P(A) + P(B) - P(A \cap B)$ $= \frac{2}{5} + \frac{5}{9} - \frac{2}{9} = \frac{11}{9}$	<p>Must have some attempt to <u>use</u></p> <p>M1</p> <p>A1cao (2)</p>

Question Number	Scheme	Marks
(f)	$\frac{P(RRR)}{P(RRR) + P(YYY)} = \frac{\frac{1}{2} \times \frac{2}{5} \times \frac{2}{3}}{\left(\frac{1}{2} \times \frac{2}{5} \times \frac{2}{3}\right) + \left(\frac{1}{2} \times \frac{2}{5} \times \frac{5}{9}\right)} = \frac{6}{11}$	<p>Probabilities must come from the product of 3 probs. from their tree diagram.</p> <p>M1 A1ft A1 cao (3) [17]</p>
<b>Notes</b>		
(b)	M1 for both cases, and +, attempted, ft their values from tree diagram. May be 4 cases of 3 balls.	
(c)	2 <sup>nd</sup> M1 for all 4 correct expressions, ft their values from tree diagram. A1 is cso	
(e)	M1 for clear attempt to <u>use</u> the correct formula, must have some correct substitution. ft their (b)	
(f)	M1 for identifying the correct probabilities and forming appropriate fraction of probs. 1 <sup>st</sup> A1ft for a correct expression using probabilities from their tree <b>Accept exact decimal equivalents. Correct answer only is full marks except in (c) and (d)</b>	

Question Number	Scheme	Marks
8.  (a)	$P(X > 168) = P\left(Z > \frac{168-160}{5}\right)$ $= P(Z > 1.6)$ $= 0.0548$ <p style="text-align: right;"><b>awrt 0.0548</b></p>	M1 A1 A1 (3)
(b)	$P(X < w) = P\left(Z < \frac{w-160}{5}\right)$ $\frac{w-160}{5} = -2.3263$ $w = 148.37$ <p style="text-align: right;"><b>awrt 148</b></p>	M1 B1 A1 (3)
(c)	$\frac{160 - \mu}{\sigma} = 2.3263$ $\frac{152 - \mu}{\sigma} = -1.2816$ $160 - \mu = 2.3263\sigma$ $152 - \mu = -1.2816\sigma$ $8 = 3.6079\sigma$ $\sigma = 2.21\dots$ $\mu = 154.84\dots$ <p style="text-align: right;"><b>awrt 2.22</b> <b>awrt 155</b></p>	M1 B1 B1 M1 A1 A1 (6) [12]
<b>Notes</b>		
(a)	M1 for an attempt to standardize 168 with 160 and 5 i.e. $\pm\left(\frac{168-160}{5}\right)$ or implied by 1.6 1 <sup>st</sup> A1 for $P(Z > 1.6)$ or $P(Z < -1.6)$ ie $z = 1.6$ and a correct inequality or 1.6 on a shaded diagram <p style="text-align: center;"><b>Correct answer to (a) implies all 3 marks</b></p>	
(b)	M1 for attempting $\pm\left(\frac{w-160}{5}\right) =$ recognizable $z$ value ( $ z  > 1$ ) B1 for $z = \pm 2.3263$ or better. Should be $z = \dots$ or implied so: $1 - 2.3263 = \frac{w-160}{5}$ is M0B0 A1 for awrt 148. This may be scored for other $z$ values so M1B0A1 is possible <p style="text-align: center;"><b>For awrt 148 only with no working seen award M1B0A1</b></p>	
(c)	M1 for attempting to standardize 160 or 152 with $\mu$ and $\sigma$ (allow $\pm$ ) <u>and</u> equate to $z$ value ( $ z  > 1$ ) 1 <sup>st</sup> B1 for awrt $\pm 2.33$ or $\pm 2.32$ seen 2 <sup>nd</sup> B1 for awrt $\pm 1.28$ seen 2 <sup>nd</sup> M1 for attempt to solve their two linear equations in $\mu$ and $\sigma$ leading to equation in just one variable 1 <sup>st</sup> A1 for $\sigma =$ awrt 2.22 . Award when 1 <sup>st</sup> seen 2 <sup>nd</sup> A1 for $\mu =$ awrt 155. Correct answer only for part (c) can score all 6 marks. NB $\sigma = 2.21$ commonly comes from $z = 2.34$ and usually scores M1B0B1M1A0A1 <p style="text-align: center;"><b>The A marks in (c) require both M marks to have been earned</b></p>	



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481

Email [publications@linneydirect.com](mailto:publications@linneydirect.com)

Order Code UA026664 January 2011

For more information on Edexcel qualifications, please visit [www.edexcel.com/quals](http://www.edexcel.com/quals)

Edexcel Limited. Registered in England and Wales no.4496750  
Registered Office: One90 High Holborn, London, WC1V 7BH

# Mark Scheme (Results)

June 2011

GCE Statistics S1 (6683) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025 or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:  
<http://www.edexcel.com/Aboutus/contact-us/>

June 2011

Publications Code UA028837

All the material in this publication is copyright

© Edexcel Ltd 2011

## EDEXCEL GCE MATHEMATICS

### General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
- ft – follow through
- the symbol  $\checkmark$  will be used for correct ft
- cao – correct answer only
- cso - correct solution only. There must be no errors in this part of the question to obtain this mark
- isw – ignore subsequent working
- awrt – answers which round to
- SC: special case
- oe – or equivalent (and appropriate)
- dep – dependent
- indep – independent
- dp decimal places
- sf significant figures
- \* The answer is printed on the paper
- $\square$  The second mark is dependent on gaining the first mark

**June 2011**  
**Statistics S1 6683**  
**Mark Scheme**

Question Number	Scheme	Marks
<b>1.</b>		
<b>(a)</b>	$S_{yy} = 4305 - \frac{181^2}{8}$ $= \underline{209.875}$ <p style="text-align: right;"><b>(awrt 210)</b></p>	<p>M1</p> <p>A1</p> <p style="text-align: right;">(2)</p>
<b>(b)</b>	$r = \frac{(-)23726.25}{\sqrt{3535237.5 \times 209.875}}$ $= \underline{-0.87104\dots}$ <p style="text-align: right;"><b>(awrt -0.871)</b></p>	<p>M1</p> <p>A1</p> <p style="text-align: right;">(2)</p>
<b>(c)</b>	Higher towns have lower temperature or temp. decreases as height increases	<p>B1</p> <p style="text-align: right;">(1)</p>
<b>(d)</b>	$S_{hh} = 3.5352375$ <p style="text-align: right;"><b>(awrt 3.54)</b> (condone 3.53)</p>	<p>B1</p> <p style="text-align: right;">(1)</p>
<b>(e)</b>	$r = \underline{-0.87104\dots}$ <p style="text-align: right;"><b>(awrt -0.871)</b></p>	<p>B1ft</p> <p style="text-align: right;">(1)</p> <p style="text-align: right;"><b>(7 marks)</b></p>
<b>Notes</b>		
<b>(a)</b>	M1 for a correct expression. Allow one slip e.g. 4350 for 4305	
<b>(b)</b>	<p>M1 for a correct expression for <math>r</math>, follow through their answer to (a). Condone no “_”</p> <p>Allow M1 for <math>\pm 0.87</math> with no working. (<math>-0.871</math> is M1A1)</p>	
<b>(c)</b>	<p>B1 Must mention <u>temperature</u> (o.e.) and <u>height</u> (above sea level) and interpret the relationship between them. Must be a correct <u>and sensible</u> comment.</p> <p>e.g. "As temperature increases the height of the sea decreases" is B0. BUT simply stating "As temperature increases the height decreases" is B1 although "As height increases the temperature decreases" would be better. Treat mention of 0.87... as ISW</p> <p>"strong negative correlation between height and temp" is B0 (no interpretation)</p> <p>" as <math>x</math> increases <math>y</math> decreases" is B0 (no mention of height and temperature)</p>	


Question Number	Scheme	Marks
(d)	B1 accept awrt 3.54 and condone 3.53 (i.e truncation)	
(e)	B1ft for awrt $-0.871$ or ft their final answer to part (b) to the same accuracy (or 3 sf) provided $-1 < r < 1$ Answer to part (e) must be a number "it's the same" is B0	
2. (a)	$\frac{23-\mu}{5} = "1.40" \quad (\text{o.e})$ $\frac{\mu=16}{16.0}$	awrt $\pm 1.40$ B1 M1A1ft (or awrt A1 (4)
(b)	<u>0.4192</u>	B1 (1)
<b>Notes</b>		
(a)	B1 for awrt $\pm 1.40$ or better seen anywhere. Condone 1.4 instead of 1.40 M1 for attempting to standardise with 23 and 5 and $\mu$ , accept $\pm$ e.g. $\frac{23-\mu}{25} = 1.40$ can score B1M0 (since using 25 not 5 for standardising) $\frac{23-\mu}{5} = 0.9192$ can score B0M1 (since have correct standardisation) Can accept equivalent equations e.g. $23-\mu = 5 \times "1.40"$ 1 <sup>st</sup> A1ft for standardised expression = to a z value ( $ z  > 1$ ). Signs must be compatible. Follow through their z e.g. $\frac{23-\mu}{5} = \text{their } z \text{ where } z > 1$ or $\frac{\mu-23}{5} = \text{their } z \text{ where } z < -1$ 2 <sup>nd</sup> A1 for 16 or awrt 16.0 if they are using a more accurate z <b>Correct answer only scores 4/4 but if any working is seen apply scheme</b>	
(b)	B1 for 0.4192 (but accept 3sf accuracy if 0.9192 – 0.5 is seen)	

Question Number	Scheme	Marks
<p><b>3.</b></p> <p><b>(a)</b></p>	$[F(3) = F(2) + P(Y=3) = (0.5 + 0.3)]$ $a = \underline{0.1}$ $d = \underline{0.8}$ $b = F(2) - a = 0.5 - 0.1 \quad \underline{\text{or}} \quad a + b = 0.5$ $c = 1 - F(3) \quad \underline{\text{or}} \quad 1 - (a + b + 0.3) \quad \underline{\text{or}} \quad a + b + c = 0.7$ $b = \underline{0.4}$ $c = \underline{0.2}$	<p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1</p> <p>(5)</p>
<p><b>(b)</b></p>	$P(3Y + 2 \geq 8) = P(Y \geq 2) \quad \underline{\text{or}} \quad 1 - P(Y \leq 1)$ $= b + 0.3 + c \quad \underline{\text{or}} \quad 1 - a = \underline{0.9}$	<p>M1</p> <p>A1ft</p> <p>(2)</p> <p><b>7</b></p>
<b>Notes</b>		
<p><b>(a)</b></p> <p>1<sup>st</sup> B1</p> <p>2<sup>nd</sup> B1</p> <p>M1</p> <p>1<sup>st</sup> A1</p> <p>2<sup>nd</sup> A1</p> <p><b>(b)</b></p> <p>M1</p> <p>A1ft</p>	<p><b>Correct answers with no (or irrelevant) working score full marks</b></p> <p>for <math>a = 0.1</math></p> <p>for <math>F(3) = 0.8</math> or <math>d = 0.8</math></p> <p>for a method for <math>b</math> or <math>c</math>. E.g. sight of <math>a + b = 0.5</math> or <math>a + b + c = 0.7</math></p> <p>If their values satisfy one of these equations then score M1 provided their values are genuine probabilities (i.e. <math>0 &lt; p &lt; 1</math>)</p> <p>This M1 may be implied by a correct answer for <math>b</math> or <math>c</math></p> <p>for <math>b</math> or <math>P(2) = 0.4</math></p> <p>for <math>c</math> or <math>P(3) = 0.2</math></p> <p>for rearranging to <math>P(Y \geq 2)</math> or <math>1 - P(Y \leq 1)</math> or selecting cases <math>Y = 2, 3</math> and <math>4</math> for <math>0.3 +</math> their <math>b +</math> their <math>c</math> or <math>1 -</math> their <math>a</math>, provided final answer <math>&lt; 1</math> and their values are probabilities.</p>	

Question Number	Scheme	Marks
<p><b>4.</b></p> <p><b>(a)</b></p> <p><b>(b)</b></p>	$(z = \pm) \frac{15 - 16.12}{1.6} (= -0.70)$ $P(Z < -0.70) = 1 - 0.7580$ $= \underline{0.2420} \quad \text{(awrt 0.242)}$ <p>[P(T &lt; t) = 0.30 implies] <math>z = \frac{t - 16.12}{1.6} = -0.5244</math></p> $\frac{t - 16.12}{1.6} = -0.5244 \Rightarrow t = 16.12 - 1.6 \times "0.5244"$ $t = \text{awrt } \underline{15.28} \text{ (allow awrt 15.28/9)}$	<p>M1</p> <p>M1</p> <p>A1</p> <p>(3)</p> <p>M1 A1</p> <p>M1</p> <p>A1</p> <p>(4)</p> <p>7</p>
<b>Notes</b>		
<p><b>(a)</b></p> <p><b>(b)</b></p>	<p style="text-align: center;"><b>Allow slips e.g. 16.2 for 16.12 for 1<sup>st</sup> M1 in (a) and (b)</b></p> <p>1<sup>st</sup> M1 for standardising expression with 15, 16.12 and 1.6 - allow <math>\pm</math></p> <p>2<sup>nd</sup> M1 for 1 - a probability (&gt; 0.5) from tables or calculator based on their standardised value</p> <p style="text-align: center;"><b>Correct answer only scores 3/3</b></p> <p style="text-align: center;"><b>In part (b) they can use any letter or symbol instead of t</b></p> <p>1<sup>st</sup> M1 for standardising with t (o.e.), 16.12 and 1.6, allow <math>\pm</math>, and setting equal to a z value</p> <p>1<sup>st</sup> A1 for an equation with <math>z = \pm 0.5244</math> or better</p> <p>e.g. <math>\frac{t - 16.12}{1.6} = \pm 0.52</math> (or 0.525) scores M1 (but A0)</p> <p>2<sup>nd</sup> M1 for solving <u>their</u> linear equation as far as <math>t = a \pm b \times 1.6</math>. Not dependent on 1<sup>st</sup> M1</p> <p>e.g. solving <math>\frac{t - 16.12}{1.6} = 0.3</math> to give <math>t = 16.12 + 1.6 \times 0.3</math> scores this M1</p> <p>Allow <math>\frac{t - 16.12}{1.6^2} = 0.3</math> to give <math>t = 16.12 + 1.6^2 \times 0.3</math> to score M1 too</p> <p>2<sup>nd</sup> A1 dependent on both M marks. Allow awrt 15.28 or awrt 15.29</p> <p>Condone awrt 15.3 if a correct expression for <math>t = \dots</math> is seen.</p> <p><b>Answers with no working:</b></p> <p>15.28 is M1A1M1A1, 15.29 is M1A0M1A1, 15.3 is M1A0M1A0</p>	

Question Number	Scheme	Marks
<p><b>5.</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p style="text-align: right;"><u>10.5</u></p> <p><math>(Q_2 =) (15.5 +) \frac{\frac{1}{2} \times 30 - 14}{8} \times 3</math> or <math>\frac{\frac{1}{2} \times 31 - 14}{8} \times 3</math>  <math>= \underline{15.875}</math> or <math>\underline{16.0625}</math></p> <p><math>\bar{x} = \frac{477.5}{30} = \underline{15.9}</math> (15.916<sup>8</sup>) [Accept <math>\frac{191}{12}</math> or <math>15\frac{11}{12}</math>]</p> <p><math>\sigma = \sqrt{\frac{8603.75}{30} - \bar{x}^2} = \underline{5.78}</math> (accept <math>s = 5.88</math>)</p> <p>Since <u>mean and median are similar (or equal or very close)</u> a normal distribution may be suitable. [Allow mean or median close to <u>mode/modal class</u>]</p> <p><math>Q_3 - Q_2 (= 8) &gt; (4.5 =) Q_2 - Q_1</math>  Therefore <u>positive skew</u></p>	<p>B1 (1)</p> <p>M1 A1 (2)</p> <p>M1, A1 M1A1ft, A1 (5)</p> <p>B1 (1)</p> <p>M1 A1 (2)</p> <p><b>(11 marks)</b></p>
<b>Notes</b>		
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p><b>In parts (a) to (c) a correct answer with no working scores full marks for that value.</b></p> <p>B1 for 10.5 which may be in the table</p> <p>M1 for a correct ratio and times 3, ignore the lower boundary for this mark  A1 for awrt 15.9 (if <math>n = 30</math> used) or awrt 16.1 (if <math>n+1 = 31</math> is used)</p> <p>1<sup>st</sup> M1 for attempt at <math>\sum fx</math> (this may be seen in the table as <math>fx: 10, 73.5, 70, 136, 82, 106</math>  [condone 1 slip] or awrt 500) and use of <math>\frac{\sum fx}{\sum f}</math> or a correct expression for mean.</p> <p>1<sup>st</sup> A1 for awrt 15.9</p> <p>2<sup>nd</sup> M1 for an attempt at <math>\sigma</math> or <math>\sigma^2</math>, can ft their mean, condone mis-labelling <math>\sigma^2 = \sqrt{\dots}</math> etc  Allow use of their <math>\sum fx^2</math> (awrt 9000)</p> <p>2<sup>nd</sup> A1ft for a correct expression including square root, ft their mean but not their <math>\sum fx^2</math>.  No label or correct label is OK but wrong label (e.g. <math>\sigma^2 = \sqrt{\dots}</math>) is A0</p> <p>3<sup>rd</sup> A1 for awrt 5.78, allow <math>s =</math> awrt 5.88. <b>SC</b> Allow M1A1A0 for awrt 5.79 if <math>\bar{x}</math> correct</p> <p>B1 for a reason implying or stating symmetry. "Time is continuous" or "evenly distributed" is B0</p>	

Question Number	Scheme	Marks
(e)	M1 for a clear reason or comparison, values not essential but comparison implying they have been found is required. A1 for stating "positive skew". Condone just "positive" but "positive correlation" is A0 <b>Do not allow arguments based on mean and median since this part relates to a different set of data.</b>	
6.		
(a)	$P(J \cup K) = 1 - 0.7$ or $0.1 + 0.15 + 0.05 = \underline{0.3}$	B1 (1)
(b)	$P(K) = 0.05 + 0.15$ or " $0.3$ " $- 0.25 + 0.15$ or " $0.3$ " $= 0.25 + P(K) - 0.15$ May be seen on Venn diagram $= \underline{0.2}$	M1 A1 (2)
(c)	$[P(K   J)] = \frac{P(K \cap J)}{P(J)}$ $= \frac{0.15}{0.25}$ $= \underline{\frac{3}{5} \text{ or } 0.6}$	M1 A1 A1 (3)
(d)	$P(J) \times P(K) = 0.25 \times 0.2 (= 0.05)$ , $P(J \cap K) = 0.15$ <u>or</u> $P(K   J) = 0.6$ , $P(K) = 0.2$ <u>or</u> may see $P(J/K) = 0.75$ and $P(J) = 0.25$ not equal therefore not independent	M1 A1ft (2)
(e)	Not independent so confirms the teacher's suspicion <u>or</u> they are linked (This requires a statement about independence in (d) or in (e))	B1ft (1) <b>(9 marks)</b>

Question Number	Scheme	Marks
<b>Notes</b>		
<p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p> <p><b>(e)</b></p>	<p>M1 for a complete method, follow through their 0.3, leading to a linear equation for <math>P(K)</math></p> <p>NB You may see this Venn diagram.                      A correct diagram (Venn or table) implies M1 in (b)                      Need not include box or 0.7  <b>Correct answer only is 2/2</b></p>	
	<p><b>In parts (c) and (d) they must have defined A and B</b></p> <p>M1 for a correct expression (including ratio) in symbols.                      1<sup>st</sup> A1 for a correct ratio of probabilities (if this is seen the M1 is awarded by implication)                      Must be in (c). Condone no LHS but wrong LHS (e.g. <math>P(K)</math> or <math>P(J/K)</math>) is M0A0                      2<sup>nd</sup> A1 for correct answer as printed only. <b>Correct answer only 3/3</b></p>	
	<p><b>Mark (d) and (e) together</b></p>	
	<p>M1 for a correct comparison of known probabilities for an independence test - ft their values. E.g. <math>P(J) \times P(K)</math> with <math>P(J \cap K)</math> <u>or</u> <math>P(K J)</math> with <math>P(K)</math> [Must have expressions]                      The values of these probabilities should be given unless they are in the question or stated elsewhere.</p>	
	<p>A1ft for correct calculations and correct comment for their probabilities</p>	
<p>B1ft ft their conclusion on independence so not independent confirms teacher...independent contradicts teacher.  <b>Methods leading to negative probabilities should score M0</b></p>		

Question Number	Scheme	Marks
<p>7.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	$(S_{fh} =) 25291 - \frac{186 \times 1085}{8}$ $= 64.75 \quad (\text{accept } 64.8)$ $b = \frac{64.75}{39.5}, \quad = 1.6392\dots \quad (\text{awrt } 1.6)$ $a = \frac{1085}{8} - b \times \frac{186}{8}, \quad = 97.512\dots \quad (\text{awrt } 97.5)$ $h = 97.5 + 1.64f$ $h = 97.5 + 1.64 \times 25, \quad = 138 \sim 139 \quad (\text{final answer in } [138, 139])$ <p>Should be reliable, since 25 cm (or <math>f</math> or footlength) is within the range of the data</p> <p>Line is for children – a different equation would apply to adults</p> <p>or Children are still growing, height will increase more than foot length</p>	<p>M1</p> <p>A1</p> <p>(2)</p> <p>M1, A1</p> <p>M1, A1</p> <p>A1ft (dep on M1M1)</p> <p>(5)</p> <p>M1, A1</p> <p>(2)</p> <p>B1, B1</p> <p>(2)</p> <p>B1</p> <p>(1)</p> <p><b>12</b></p>
<b>Notes</b>		
	<p style="text-align: center;"><b>[NB <math>r = 0.871</math> so do not confuse this with question 1]</b></p> <p>(a) M1 for attempting a correct expression [allow a copying slip e.g. 25921]</p> <p>(b) 1<sup>st</sup> M1 for a correct expression for <math>b</math>, ft their part (a) but not <math>S_{fh} = 25291</math>  1<sup>st</sup> A1 for awrt 1.6  2<sup>nd</sup> M1 for use of <math>a = \bar{h} - b \times \bar{f}</math>, ft their value for <math>b</math>. Must use <math>\bar{h}</math> and <math>\bar{f}</math> not values from table.  2<sup>nd</sup> A1 for awrt 97.5 [NB <math>a = 135 - 1.63 \times 23 = 97.51</math> but M0A0 since not using <math>\bar{h}</math> and <math>\bar{f}</math> ]  3<sup>rd</sup> A1ft for an equation for <math>h</math> and <math>f</math> with <u>their</u> coefficients to 3sf. <b>Dependent on both Ms</b>  Must be 3sf not awrt. Give this mark if seen in (c). Equation must be in <math>h</math> and <math>f</math> not <math>y</math> and <math>x</math>.</p> <p>(c) M1 for using <u>their</u> equation and <math>f = 25</math> to find <math>h</math>  A1 for their final answer in [138, 139]. Can give if they have 137.7... but round to 138</p> <p>(d) 1<sup>st</sup> B1 for suggesting it <u>is</u> reliable  2<sup>nd</sup> B1 for mentioning that 25 cm is within range of data. “interpolation” or “not extrapol” B1  Use of “it” or a comment that height is in range is B0 but apply ISW</p> <p>(e) B1 for some comment that states a difference between children and teachers(adults)  Must mention <u>teacher/adults</u> and <u>children</u>  e.g. ".teacher is not in same age group as the children", "equation is for children not adults"  “children and adults are different populations”  "teacher will be taller" is B0 since no mention of children.  “equation is <u>only</u> valid for children” is OK since “only” implies not suitable for adults</p> <p><u>Or</u> Reference to different growth rates</p>	

Question Number	Scheme	Marks
<b>8.</b>		
(a)	$1 = p + (0.25 + 0.25 + 0.2 + 0.2), \Rightarrow p = \frac{1}{10} \text{ or } 0.1$	M1, A1 (2)
(b)	$E(S) = \frac{1}{4} + 2 \times \frac{1}{4} + 4 \times \frac{1}{5} + 5 \times \frac{1}{5}, \text{ (or equiv. in decimals)} = \underline{2.55}$	M1, A1 (2)
(c)	$E(S^2) = \frac{1}{4} + \frac{2^2}{4} + \frac{4^2}{5} + \frac{5^2}{5} \text{ or } 0.25 + 1 + 3.2 + 5 = \underline{9.45} (*)$	M1, A1cso (2)
(d)	$\text{Var}(S) = 9.45 - (E(S))^2, = \underline{2.9475} \text{ or } \frac{1179}{400} \text{ (accept awrt 2.95)}$	M1, A1 (2)
(e)	$P(5 \text{ and } 5) = \left(\frac{1}{5}\right)^2, = \underline{\frac{1}{25} \text{ or } 0.04}$	M1, A1 (2)
(f)	$P(4, 4, 2) = \left(\frac{1}{5}\right)^2 \times \frac{1}{4} \times 3 \text{ (} = 0.03 \text{ or } \frac{3}{100} \text{)}$ $P(4, 4, 4) = \left(\frac{1}{5}\right)^3 \text{ (} = 0.008 \text{ or } \frac{1}{125} \text{)}$ $P(\text{Tom wins in 3 spins}) = \underline{0.038}$	M1, M1 B1 A1 (4)
(g)	$P(\bar{5} \cap 5 \cap 5) + P(5 \cap \bar{5} \cap 5) = \frac{4}{5} \times \left(\frac{1}{5}\right)^2 \times 2 = \underline{0.064} \text{ or } \frac{8}{125}$	M1, M1, A1 (3)
<b>17</b>		

**Notes**

(a)	M1 for clear attempt to use sum of probabilities = 1 (fractions or decimals) Ans only 2/2
(b)	M1 for at least 2 correct terms ( $\neq 0$ ) of the expression. 2.55 with no working scores M1A1 <b>Any division by <math>k</math> (usually 5) in (b) or (c) or (d) scores M0</b>
(c)	M1 for at least 3 correct, non-zero terms of the expression seen, allow decimals. A1cso for the full expression (with 9.45) seen. Must be cso but can ignore wrong $p$ .
(d)	M1 for a correct expression (9.45 seen), can fit their $E(S)$ . May see $\sum (x - "2.55")^2 \times P(X = x)$ A1 accept awrt 2.95 Answer only can score M1 for correct fit and A1 for awrt 2.95 <b>Answer only in (e) and (f) is full marks, in (g) is no marks</b>
(e)	M1 for $\left(\frac{1}{5}\right)^2$ Condone $P(5) \times P(5) = 0.25 \times 0.25$ . [Beware 0.4 is A0]
(f)	1 <sup>st</sup> M1 for $\left(\frac{1}{5}\right)^2 \times \frac{1}{4}$ or 0.01 seen 2 <sup>nd</sup> M1 for multiplying a $p^2q$ probability by 3 ( $p, q \in (0,1)$ ). B1 for $(0.2)^3$ or better seen
(g)	1 <sup>st</sup> M1 for $\frac{4}{5} \times \left(\frac{1}{5}\right)^2$ or all cases considered and correct attempt at probabilities. 2 <sup>nd</sup> M1 for multiplying a $p^2(1-p)$ probability by 2. <b>Beware <math>(0.4)^3 = 0.064</math> is M0M0A0</b>



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UA028837 June 2011

For more information on Edexcel qualifications, please visit  
[www.edexcel.com/quals](http://www.edexcel.com/quals)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual



Llywodraeth Cynulliad Cymru  
Welsh Assembly Government



# Mark Scheme (Results)

January 2012

GCE Statistics S1 (6683) Paper 1

Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at [www.edexcel.com](http://www.edexcel.com).

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our **Ask The Expert** email service helpful.

Ask The Expert can be accessed online at the following link:

<http://www.edexcel.com/Aboutus/contact-us/>

January 2012

Publications Code UA030899

All the material in this publication is copyright

© Pearson Education Ltd 2012

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

**EDEXCEL GCE MATHEMATICS****General Instructions for Marking**

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\checkmark$  will be used for correct ft
  - cao – correct answer only
  - cso – correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - $\square$  The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

**General Principals for Core Mathematics Marking**

(But note that specific mark schemes may sometimes override these general principles).

**Method mark for solving 3 term quadratic:**1. Factorisation

$$(x^2 + bx + c) = (x + p)(x + q), \text{ where } |pq| = |c|, \text{ leading to } x = \dots$$

$$(ax^2 + bx + c) = (mx + p)(nx + q), \text{ where } |pq| = |c| \text{ and } |mn| = |a|, \text{ leading to } x = \dots$$

2. Formula

Attempt to use correct formula (with values for  $a$ ,  $b$  and  $c$ ), leading to  $x = \dots$

3. Completing the square

$$\text{Solving } x^2 + bx + c = 0 : \quad \left(x \pm \frac{b}{2}\right)^2 \pm q \pm c, \quad q \neq 0, \quad \text{leading to } x = \dots$$

**Method marks for differentiation and integration:**1. Differentiation

Power of at least one term decreased by 1. ( $x^n \rightarrow x^{n-1}$ )

2. Integration

Power of at least one term increased by 1. ( $x^n \rightarrow x^{n+1}$ )

**Use of a formula**

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:

Method mark for quoting a correct formula and attempting to use it, even if there are mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

**January 2012**  
**6683 Statistics S1**  
**Mark Scheme**

Question Number	Scheme	Marks
<b>1 (a)</b>	14, 5	M1 A1  (2)
<b>(b)</b>	$21 + 45 + 3 = 69$	M1 A1  (2)  <b>Total 4</b>
<b>NOTES</b>		
<b>(a)</b>	M1 for $2 \times 7$ or 14 or $5 \times 1$ or 5  A1 for both 14 and 5	
<b>(b)</b>	M1 for $21 + 45 + (0 < \text{frequency} < 9)$  A1 for 69 only.  69 no working, award M1A1      Incorrect answer with no working M0A0	

Question Number	Scheme	Marks
<p><b>2 (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p>	<p>(<i>R</i> and <i>S</i> are mutually) exclusive.</p> <p><math>\frac{2}{3} = \frac{1}{4} + P(B) - P(A \cap B)</math>                      use of Addition Rule</p> <p><math>\frac{2}{3} = \frac{1}{4} + P(B) - \frac{1}{4} \times P(B)</math>                      use of independence</p> <p><math>\frac{5}{12} = \frac{3}{4} P(B)</math></p> <p><math>P(B) = \frac{5}{9}</math></p> <p><math>P(A' \cap B) = \frac{3}{4} \times \frac{5}{9} = \frac{15}{36} = \frac{5}{12}</math></p> <p><math>P(B' A) = \frac{(1 - (b)) \times 0.25}{0.25}</math>    or <math>P(B')</math> or <math>\frac{1}{\frac{1}{4}}</math></p> <p><math>= \frac{4}{9}</math></p>	<p>B1 (1)</p> <p>M1</p> <p>M1 A1</p> <p>A1 (4)</p> <p>M1A1ft (2)</p> <p>M1</p> <p>A1 (2)</p> <p><b>Total 9</b></p>
<p><b>NOTES</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	<p>B1 for '(mutually) exclusive' or 'cannot occur at the same time' seen or equivalent. 'Intersection is zero' or 'no overlaps' without further explanation is B0.</p> <p>M1 for use of Addition Formula, including an intersection, with at least one probability substituted. Intersection must be explicitly considered for this mark.</p> <p>Accept <math>\frac{2}{3} = \frac{1}{4} + P(B) - 0</math> for M1.</p> <p>M1 for <math>P(A \cap B) = \frac{1}{4} P(B)</math></p> <p>A1 for completely correct equation or equivalent.</p> <p>A1 for <math>\frac{5}{9}</math> or exact equivalent..</p> <p>Venn Diagram with 2 overlapping closed curves and correct values possibly without <math>\frac{1}{3}</math>, award M1M1A1.</p> <p>M1 for <math>\frac{3}{4}</math> x 'their <math>P(B)</math>' or 'their <math>P(B) - P(A \cap B)</math>' or <math>P(A \cup B) - P(B) = \frac{2}{3} - \frac{1}{4}</math></p> <p>Or <math>P(A' \cap B) = P(A') + \text{'their } P(B) - P(A' \cup B) = \frac{3}{4} + \frac{5}{9} - \frac{8}{9}</math></p> <p>A1 for <math>\frac{5}{12}</math> or follow through from their method. Accept exact equivalent.</p> <p>Correct answer only with no working M1A1 but must be clearly labelled (c).</p>	

(d) M1 for using  $1 - \text{their } P(B)$  or  $(P(A \cup B) - P(A))/P(A)$  or  $(P(A) - P(A \cap B))/P(A)$  with a correct attempt at the numerator and denominator. If mutually exclusive is


assumed then the last option gives  $\frac{1}{4}$  for M1.

A1 for  $\frac{4}{9}$  or exact equivalent.

For part (c) follow through their stated values; **do not** follow through incorrectly labelled regions on a Venn Diagram.

Throughout the question we require probabilities between 0 and 1 for method marks.

Venn Diagram:




Question Number	Scheme	Marks
<p><b>3 (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p> <p><b>(e)</b></p>	$\frac{5}{21} + \frac{2k}{21} + \frac{7}{21} + \frac{k}{21} = 1$ $\frac{12 + 3k}{21} = 1$ $k = 3 \quad * \text{ AG}$ <p style="text-align: right;">required for both methods</p> $\frac{11}{21}$ $E(X) = 2 \times \frac{5}{21} + 3 \times \frac{6}{21} + 4 \times \frac{7}{21} + 6 \times \frac{1}{7}$ $= 3 \frac{11}{21} \text{ or } \frac{74}{21} \text{ or awrt } 3.52$ $E(X^2) = 2^2 \times \frac{5}{21} + 3^2 \times \frac{6}{21} + 4^2 \times \frac{7}{21} + 6^2 \times \frac{1}{7}$ $= 14$ $\text{Var}(X) = 14 - \left(3 \frac{11}{21}\right)^2$ $= 1 \frac{257}{441} \text{ or } \frac{698}{441} \text{ or awrt } 1.6$ $\text{Var}(7X - 5) = 7^2 \text{Var}(X)$ $= 77 \frac{5}{9} \text{ or } \frac{698}{9} \text{ or awrt } 77.6$	<p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>(2)</p> <p>(1)</p> <p>(2)</p> <p>(2)</p> <p>(4)</p> <p><b>Total 11</b></p>
<p><b>NOTES</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>M1 Award for verification. Sub in k=3 and show <math>\sum xP(X = x) = 1</math>. Require at least three correct terms seen or line 2 of scheme. A1 Correct solution only including verification.</p> <p>B1 Award for exact equivalent.</p> <p>M1 At least two correct terms required for method, follow through 'their k' for method. Correct answer only, award M1 A1.</p> <p>M1 At least two correct terms required for method. M0 if probability is squared. Correct answer only, award M1 A1. Accept exact equivalent of 14 for A1.</p> <p>M1 for use of correct formula in both. 1.6 can be implied by correct final answer. Working needs to be clearly labelled to award first method mark without second stage of calculation. If a new table for values of <math>7X - 5</math> is used, so <math>Y = 7X - 5</math> <math>E(Y^2) = \frac{9751}{21}</math> ; <math>\text{Var}(Y) = 77 \frac{5}{9}</math> or <math>\frac{698}{9}</math> or awrt 77.6 Award M1A1; M1A1 If any attempt to divide by 4 seen as part of working award M0 for that part.</p>	

Question Number	Scheme	Marks
<p><b>4 (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p> <p><b>(e)</b></p>	<p>60</p> <p><math>Q_1 = 46</math> <math>Q_2 = 56</math> <math>Q_3 = 64</math></p> <p>mean = 55.48.... or <math>\frac{2497}{45}</math> awrt 55.5</p> <p><math>sd = \sqrt{\frac{143369}{45} - \left(\frac{2497}{45}\right)^2}</math> = 10.342... (s = 10.459..) anything which rounds to 10.3 (or s = 10.5)</p> <p>Mean &lt; median &lt; mode or <math>Q_2 - Q_1 &gt; Q_3 - Q_2</math> with or without their numbers or median closer to upper quartile (than lower quartile) or (mean-median)/sd &lt; 0; negative skew;</p> <p>mean = <math>(55 - 5) \times 0.9</math> = 45 sd = <math>10 \times 0.9</math> = 9</p>	<p>B1 (1)</p> <p>B1 B1 B1 (3)</p> <p>B1 awrt 55.5</p> <p>M1</p> <p>A1 (3)</p> <p>B1</p> <p>B1dep (2)</p> <p>M1 A1 M1 A1 (4)</p> <p><b>Total 13</b></p>
<p><b>NOTES</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p>B1 60 only</p> <p>Award each B1 for correct answer only in this order.</p> <p>M1 for use of correct formula, including square root. Correct answers with no working B1M1A1.</p> <p>B1 any correct comparison of a pair of mean, median and mode using their values. B1 for 'negative skew' or allow (almost) symmetrical dependent upon correct reason.</p> <p>M1 for <math>(55 \text{ or } 55.5 - 5) \times 0.9</math> A1 for the correct answer only. M1 for <math>(10 \text{ or } 10.3 \text{ or } 10.5) \times 0.9</math> A1 for the correct answer only.</p>	

Question Number	Scheme	Marks
<b>5 (a)</b>	$S_{tt} = 2688 - \frac{158^2}{10} = 191.6$ $S_{tw} = 1760.62 - \frac{158 \times 111.75}{10} = -5.03$	awrt 192 awrt -5.03 (3)
<b>(b)</b>	$r = \frac{-5.03}{\sqrt{191.6 \times 0.16}} = -0.908469\dots$	awrt -0.908(5) (2)
<b>(c)</b>	$b = \frac{-5.03}{191.6} = -0.0263$ $a = 11.175 + 0.0263 \times 15.8$ $= 11.59$ $w = 11.6 - 0.0263t$	awrt -0.026 M1 A1 M1 A1 (4)
<b>(d)</b>	The explanatory variable is the age of each coin. This is because the age is set and the weight varies.	B1 B1 (2)
<b>(e) (i)</b> <b>(ii)</b>	awrt 11.5 Decrease (in weight of coin of 0.1052 g) = 0.1 or -0.1 or increase of -0.1	awrt(-0.1) B1 B1 (2)
<b>(f)</b>	Decrease; removing the fake will result in a better linear fit so $r$ will be closer to -1	B1;B1 (2)
<b>NOTES</b>		<b>Total 15</b>
<b>(a)</b> <b>(b)</b> <b>(c)</b> <b>(d)</b> <b>(e)</b> <b>(f)</b>	M1 for correct attempt at either method, A1 awrt 192 A1 awrt -5.03 M1 for correct attempt at use of formula, square root required. A1 awrt -0.908(5) M1 require 'their -5.03' as numerator and /their 191.6' as denominator. A1 awrt -0.026 M1 for use of correct formula with $b$ or 'their $b$ '; require -- or + and values in the correct place. A1 for equation as written with values awrt 3 sf. with $w$ and $t$ . Accept fractional answers that are accurate to 3sf when evaluated as decimals B1 for 'Age' or $t$ or 'years' B1 for 'you use age / $t$ to predict $w$ ' or 'you can control $t$ / age' or 'weight depends on age' or similar B1 awrt 11.5 B1 awrt -0.1 but 'decrease of -0.1' is B0. B1 for Decrease only but 'mod $r$ increases' explicitly stated in words or symbols award B1. B1 accept 'stronger correlation' or 'increase in correlation' or 'better linear fit' or ' $r$ closer to -1' or 'points are closer to a straight line' or 'point is an outlier' or equivalent	

<b>Special Case 1</b>	<p>Attempt to calculate <math>S_{tw}</math></p> $\sum tw = 1669.62, \sum t = 153, \sum w = 91.75 \text{ or } S_{tw} = 1660.62 - \frac{153 \times 91.75}{9} \text{ or awrt } 101$ <p>or <math>S_{tw} &gt; 0</math> with some calculation  “Increase”</p>	<p>B1  B1  (2)</p>
<b>Special Case 2</b>	<p>Attempt to calculate <math>S_{ww}</math></p> $\sum w^2 = 1248.96625 - 400 = 848.96625 \text{ or awrt } 849 \text{ or } S_{ww} = 848.96625 - \frac{91.75^2}{9}$ <p>or awrt -86.4 or <math>S_{ww} &lt; 0</math></p>	<p>B2  (2)</p>
<b>Special Case 3</b>	<p>Argument based on standard deviation.</p> <p>e.g. <math>\sigma_w \approx 0.126</math> and <math>\bar{w} = 11.175</math> so fake coin is over 69 sds away from the mean  ‘(very) unlikely’ or ‘impossible’</p>	<p>B1  B1  (2)</p>

Question Number	Scheme	Marks
<p><b>6 (a)</b></p>	<div style="display: flex; align-items: flex-start;"> <div style="flex: 1;">  </div> <div style="flex: 2; padding-left: 10px;"> <p>3 closed curves and 25 in correct place</p> <p>15,10,5</p> <p>15,3,20</p> <p>Labels R, S, C and box</p> </div> </div>	<p>M1</p> <p>A1</p> <p>A1</p> <p>B1</p>
<p><b>(b)</b></p>	<p>All values/100 or equivalent fractions award accuracy marks.</p> <p>7/100 or 0.07</p> <p>M1 for ('their 7' in diagram or here)/100</p>	<p>(4)</p> <p>M1 A1</p>
<p><b>(c)</b></p>	<p><math>(3+5)/100 = 2/25</math> or 0.08</p>	<p>(2)</p> <p>M1A1</p>
<p><b>(d)</b></p>	<p><math>(25+15+10+5)/100 = 11/20</math> or 0.55</p>	<p>(2)</p> <p>M1 A1</p>
<p><b>(e)</b></p>	<p><math display="block">P(S \cap C'   R) = \frac{P(S \cap C' \cap R)}{P(R)}</math></p> <p>Require denominator to be 'their 65' or 'their <math>\frac{65}{100}</math>'</p> <p><math display="block">= \frac{15}{65}</math></p> <p>require 'their 15' and correct denominator of 65</p> <p><math display="block">= \frac{3}{13}</math> or exact equivalents.</p>	<p>(3)</p> <p>M1</p> <p>A1</p> <p>A1</p>
<p><b>NOTES</b></p>		<p><b>Total 13</b></p>
<p><b>(b)</b></p>	<p>M1 for 'their 7'/100 seen.</p> <p>A1 Correct answer only</p> <p>In parts (c) and (d) we require "/100" for methods to be awarded. Also check their values and award correct method if they follow from their Venn Diagram.</p>	
<p><b>(c)</b></p>	<p>M1 For ('their 3'+ 'their 5')/100. <math>\frac{8}{48}</math> award M0.</p>	
<p><b>(d)</b></p>	<p>A1 Correct answer only or equivalent.</p>	
<p><b>(d)</b></p>	<p>M1 Accept sum of their 4 values from the Venn diagram /100.</p>	
<p><b>(d)</b></p>	<p>A1 Correct answer only or equivalent</p>	
<p><b>(e)</b></p>	<p>M1 Attempt to use correct formula for conditional probability.</p>	
<p><b>(e)</b></p>	<p>Award for correct formula and a denominator of 'their 65' or 'their 65/100'.</p>	
<p><b>(e)</b></p>	<p>A1 for 'their 15'/65 only.</p>	
<p><b>(e)</b></p>	<p>A1 for exact equivalent answers, including 15/65.</p>	
<p><b>(e)</b></p>	<p>In all parts correct answers with no working award full marks.</p>	

Question Number	Scheme	Marks
<p><b>7 (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p>	$P(W < 224) = P\left(z < \frac{224 - 232}{5}\right)$ $= P(z < -1.6)$ $= 1 - 0.9452$ $= 0.0548$ $0.5 - 0.2 = 0.3$ $\frac{w - 232}{5} = 0.5244$ $w = 234.622$ $0.2 \times (1 - 0.2)$ $2 \times 0.8 \times (1 - 0.8) = 0.32$	<p>M1</p> <p>M1</p> <p>A1</p> <p>(3)</p> <p>M1</p> <p>0.3 or 0.7 seen</p> <p>M1</p> <p>0.5244 seen</p> <p>B1; M1</p> <p>A1</p> <p>(4)</p> <p>M1</p> <p>M1 A1</p> <p>(3)</p> <p><b>Total 10</b></p>
<p><b>NOTES</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	<p>M1 for standardising with 232 and 5. (i.e. not <math>5^2</math> or <math>\sqrt{5}</math>). Accept <math>\pm \frac{w - 232}{5}</math>.</p> <p>M1 for finding (1 - a probability &gt; 0.5)</p> <p>A1 awrt 0.0548</p> <p>M1 Can be implied by use of <math>\pm 0.5244</math> or <math>\pm (0.52 \text{ to } 0.53)</math></p> <p>B1 for <math>\pm 0.5244</math> only.</p> <p>Second M1 standardise with 232 and 5 and equate to <math>z</math> value of (0.52 to 0.53) or (0.84 to 0.85)</p> <p>1 - <math>z</math> used award second M0.</p> <p>Require consistent signs i.e. <math>\frac{232 - w}{5} = -0.5244</math> or negative <math>z</math> value for M1.</p> <p>A1 dependent upon second M mark for awrt 235 but see note below.</p> <p>Common errors involving probabilities and not <math>z</math> values:</p> <p><math>P(Z &lt; 0.2) = 0.5793</math> used instead of <math>z</math> value gives awrt 235 but award M0B0M0A0</p> <p><math>P(Z &lt; 0.8) = 0.7881</math> used instead of <math>z</math> value award M0B0M0A0.</p> <p>M1B0M0A0 for 0.6179, M1B0M0A0 for 0.7580</p> <p>M1 for 0.16 seen</p> <p>M1 for '<math>2 \times p(1 - p)</math>'</p> <p>A1 0.32 correct answer only</p>	



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481  
Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)  
Order Code UA030899 January 2012

For more information on Edexcel qualifications, please visit  
[www.edexcel.com/quals](http://www.edexcel.com/quals)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual  




Llywodraeth Cynulliad Cymru  
Welsh Assembly Government





# Mark Scheme (Results)

Summer 2012

GCE Statistics S1  
(6683) Paper 1

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2012

Publications Code UA033137

All the material in this publication is copyright

© Pearson Education Ltd 2012

**Summer 2012**  
**6683 Statistics S1**  
**Mark Scheme**

**General Marking Guidance**

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## EDEXCEL GCE MATHEMATICS

### General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used if you are using the annotation facility on ePEN.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\checkmark$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - $\square$  The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

## General Principles for Mathematics Marking

(But note that specific mark schemes may sometimes override these general principles).

### Method mark for solving 3 term quadratic:

#### 1. Factorisation

$(x^2 + bx + c) = (x + p)(x + q)$ , where  $|pq| = |c|$ , leading to  $x = \dots$

$(ax^2 + bx + c) = (mx + p)(nx + q)$ , where  $|pq| = |c|$  and  $|mn| = |a|$ , leading to  $x = \dots$

#### 2. Formula

Attempt to use correct formula (with values for  $a$ ,  $b$  and  $c$ ), leading to  $x = \dots$

#### 3. Completing the square

Solving  $x^2 + bx + c = 0$  :  $(x \pm \frac{b}{2})^2 \pm q \pm c$ ,  $q \neq 0$ , leading to  $x = \dots$

### Method marks for differentiation and integration:

#### 1. Differentiation

Power of at least one term decreased by 1. ( $x^n \rightarrow x^{n-1}$ )

#### 2. Integration

Power of at least one term increased by 1. ( $x^n \rightarrow x^{n+1}$ )

### Use of a formula

Where a method involves using a formula that has been learnt, the advice given in recent examiners' reports is that the formula should be quoted first.

Normal marking procedure is as follows:


Method mark for quoting a correct formula and attempting to use it, even if there are mistakes in the substitution of values.

Where the formula is not quoted, the method mark can be gained by implication from correct working with values, but may be lost if there is any mistake in the working.

**Summer 2012**  
**6683 Statistics S1**  
**Mark Scheme**

Question	Scheme	Marks																
<b>1.</b>	<table border="1" style="margin: auto; border-collapse: collapse;"> <tr> <td style="padding: 5px;"><math>x</math></td> <td style="padding: 5px;"><math>-1</math></td> <td style="padding: 5px;"><math>0</math></td> <td style="padding: 5px;"><math>1</math></td> <td style="padding: 5px;"><math>2</math></td> </tr> <tr> <td style="padding: 5px;"><math>P(X=x)</math></td> <td style="padding: 5px;"><math>4k</math></td> <td style="padding: 5px;"><math>k</math></td> <td style="padding: 5px;"><math>0</math></td> <td style="padding: 5px;"><math>k</math></td> </tr> </table>	$x$	$-1$	$0$	$1$	$2$	$P(X=x)$	$4k$	$k$	$0$	$k$	M1						
	$x$	$-1$	$0$	$1$	$2$													
	$P(X=x)$	$4k$	$k$	$0$	$k$													
	(a)	$4k + k + (0) + k = 1$ (Allow verify approach) $6k = 1 \Rightarrow k = \frac{1}{6}$ (*)	A1 A1cso (3)															
(b)	$[E(X)] = -4k + (0 + 0) + 2k$ or $-2k$ or $-1 \times \frac{4}{6} + 2 \times \frac{1}{6}$ $= -\frac{1}{3}$ (or $-0.\overline{3}$ )	M1 A1 (2)																
(c)	$[E(X^2)] = (-1)^2 \times 4k + (0 + 0) + 2^2 k$ or $4k + 4k$ or $(-1)^2 \times \frac{4}{6} + 2^2 \times \frac{1}{6}$ (o.e.) $= \frac{4}{3}$ (*)	M1 A1cso (2)																
(d)	$[\text{Var}(X)] = \frac{4}{3} - \left(-\frac{1}{3}\right)^2$ or $8k - 4k^2 = \left[\frac{11}{9}\right]$ $\text{Var}(1-3X) = (-3)^2 \text{Var}(X)$ or $9\text{Var}(X)$ $= 11$	<table style="margin-left: auto; margin-right: auto;"> <tr> <td style="padding: 5px;"><math>Y = 1 - 3X : 4</math></td> <td style="padding: 5px;"><math>1</math></td> <td style="padding: 5px;"><math>-2</math></td> <td style="padding: 5px;"><math>-5</math></td> </tr> <tr> <td style="padding: 5px;">Prob:</td> <td style="padding: 5px;"><math>4k</math></td> <td style="padding: 5px;"><math>k</math></td> <td style="padding: 5px;"><math>0</math></td> </tr> <tr> <td colspan="4" style="text-align: center; padding: 5px;">And <math>E(Y) = 12k</math></td> </tr> <tr> <td colspan="4" style="padding: 5px;"><math>E(Y^2) = 90k</math> and <math>\text{Var}(Y) = 90k - 144k^2</math></td> </tr> </table>	$Y = 1 - 3X : 4$	$1$	$-2$	$-5$	Prob:	$4k$	$k$	$0$	And $E(Y) = 12k$				$E(Y^2) = 90k$ and $\text{Var}(Y) = 90k - 144k^2$			
$Y = 1 - 3X : 4$	$1$	$-2$	$-5$															
Prob:	$4k$	$k$	$0$															
And $E(Y) = 12k$																		
$E(Y^2) = 90k$ and $\text{Var}(Y) = 90k - 144k^2$																		
<b>Notes</b>		<b>[10]</b>																
<b>Verify</b>	<p>(a) M1 for attempt at <math>P(X=x)</math> with at least 2 correct. Do not give for 4, 1, etc but <math>\frac{4}{6}, \frac{1}{6}</math> are OK                      1<sup>st</sup> A1 for at least <math>4k + k + k = 1</math> seen. Allow <math>\frac{4}{6} + \frac{1}{6} + \frac{1}{6} = 1</math> [Must see = 1]                      2<sup>nd</sup> A1cso provided previous 2 marks are scored and no incorrect working seen                      It's not essential to see <math>P(X = -1) = 4k</math> etc but if wrongly assigned probabilities such as <math>P(X = 2) = 4k</math> and <math>P(X = -1) = k</math> are seen then the final A1 is lost.                      To score final A1cso there must be a comment such as "therefore <math>k = \frac{1}{6}</math>"</p> <p style="text-align: center;"><b>Division by 4 (or any other n) in (b), (c) or (d) is M0. Do not apply ISW</b></p> <p>(b) M1 for a full correct expression for <math>E(X)</math>, ft their <u>probabilities</u>. Allow in terms of <math>k</math>.                      A1 for <math>-\frac{1}{3}</math> or exact equivalent only. Just <math>-\frac{1}{3}</math> scores M1A1</p> <p>(c) M1 for evidence of both non-zero terms seen. May be simplified but 2 terms needed.                      A1cso for M1 seen leading to <math>\frac{4}{3}</math> or any exact equivalent. Condone <math>-1^2 \times 4k</math> but not <math>-4k</math></p> <p>(d) 1<sup>st</sup> M1 for correct attempt at <math>\text{Var}(X)</math> - follow through their <math>E(X)</math> and allow in terms of <math>k</math>                      Award if a correct formula is seen and some correct substitution made.                      2<sup>nd</sup> M1 for correct use of <math>\text{Var}(aX+b)</math>. Condone <math>-3^2 \text{Var}(X)</math> if it eventually yields <math>9\text{Var}(X)</math>                      A1cao for 11 only</p>																	

Question	Scheme	Marks
<p>2.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	$[S_{xy} =] 23070 - \frac{477 \times 480}{12} [= 3990]$ $r = \frac{"3990"}{\sqrt{5606.25 \times 4244}}$ $= 0.81799\dots$ <p style="text-align: right;"><b>awrt 0.818</b></p> <p><b>0.818</b></p> <p>Positive correlation <u>or</u> value of <math>r</math> is close to 1 <u>or</u> value of <math>r &gt; 0</math> (NOT "high/ strong correlation")</p> <p>So there <u>is support</u> for the bank's claim <u>or</u> "increase in unemployment is accompanied by increase in house repossessions"</p>	<p>B1</p> <p>M1</p> <p>A1 (3)</p> <p>B1ft (1)</p> <p>B1</p> <p>B1 (2)</p> <p style="text-align: right;"><b>[6]</b></p>
<b>Notes</b>		
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(c) SC</p>	<p><b>Marks for part (a) must be seen in (a), do not award if only seen in (b)</b></p> <p>B1 for a correct expression for <math>S_{xy}</math></p> <p>M1 for correct attempt at <math>r</math> f.t. their 3990 but <math>\frac{23070}{\sqrt{5606.25 \times 4244}}</math> is M0</p> <p>A1 for awrt 0.818 If an answer of 0.82 only is seen then B1M1A0 can be given</p> <p>B1ft for awrt 0.818 or f.t. their answer to part (a) for <math> r  &lt; 1</math>. Allow 2sf or 1sf follow through Answer in (b) must be correct or match one of their answers in (a). Must be a number.</p> <p>1<sup>st</sup> B1 for a reason of positive correlation (allow even if <math>r &gt; 1</math>) "positive skew" or "positive gradient" is B0 but 2<sup>nd</sup> B1 is still possible</p> <p>2<sup>nd</sup> B1 for a comment that suggest this supports the claim. Marks in (c) are independent but first B1 requires some idea of <u>positive</u> correlation</p> <p>If <math> r  &lt; 0.2</math> allow this alternative to the mark scheme:</p> <p>1<sup>st</sup> B1 for saying there is no or little correlation</p> <p>2<sup>nd</sup> B1 for a comment that says this does <u>not</u> support the bank's claim</p>	

Question	Scheme	Marks
<p>3. (a)</p>	 <p>(b) Points (appear to) lie close to a (straight) line <u>or</u> “strong /high correlation”</p> <p>(c) <math>\sum p = 93</math> and <math>\sum t = 34</math> (may be seen in table)  <math>S_{pt} = 694 - \frac{93 \times 34}{6} = [167]</math> <u>or</u> <math>S_{pp} = 1967 - \frac{93^2}{6} = [525.5]</math>  <math>S_{pt} = 167</math> ; <math>S_{pp} = \text{awrt } 526</math></p> <p>(d) <math>b = \left[ \frac{S_{pt}}{S_{pp}} \right] = \frac{167}{525.5} = [0.31779\dots]</math> (check their answer if expression not seen)  <math>a = \frac{34}{6} - 0.31779\dots \times \frac{93}{6} = 5.666\dots - 0.31779\dots \times 15.5 = , 0.74088\dots</math> awrt 0.74  <math>t = 0.741 + 0.318p</math> (Accept <math>a = \frac{2336}{3153}</math> and <math>b = \frac{334}{1051}</math> in their equation)</p> <p>(e) <math>(\bar{p}, \bar{t}) = (15.5, 5.7)</math> plotted on the graph (not wholly outside the circle)            Correct line plotted as per overlay. For <math>p = 5</math>; <math>2 &lt; t &lt; 3</math> <u>and</u> for <math>p = 30</math>; <math>10 &lt; t &lt; 11</math>            Their line must stretch roughly as far as the points and go through the <math>(\bar{p}, \bar{t})</math> circle</p> <p>(f) <math>t = 0.741 + 0.318 \times 16 = 5.825\dots</math> awrt 5.8</p>	<p>Use overlay</p> <p>B1 B1</p> <p>(2)</p> <p>B1 (1)</p> <p>M1 M1 A1; A1 (4)</p> <p>B1ft</p> <p>M1, A1</p> <p>A1 (4)</p> <p>B1 B1 (2)</p> <p>M1 A1 (2)</p> <p>[15]</p>
<b>Notes</b>		
<p>(a)</p> <p>(c)</p> <p>(d)</p> <p>(f)</p>	<p>B2 for all 6 data points plotted correctly. B1 for any 5 correct. Points not wholly outside the circles.</p> <p>1<sup>st</sup> M1 for attempting <math>\sum p</math> and <math>\sum t</math>. Allow <math>80 &lt; \sum p &lt; 100</math> and <math>30 &lt; \sum t &lt; 40</math>            2<sup>nd</sup> M1 for one correct expression for <math>S_{pt}</math> or <math>S_{pp}</math>, f.t. their <math>\sum p</math> and <math>\sum t</math>. 1<sup>st</sup> A1 for <math>S_{pt}</math> 2<sup>nd</sup> for <math>S_{pp}</math></p> <p>B1ft for correct expression for the gradient, f.t. their 167 and 525.5 from (c)            M1 for correct use of <math>a = \bar{t} - b\bar{p}</math> f.t. their values. Condone 5.6 for <math>\bar{t}</math>            1<sup>st</sup> A1 for awrt 0.74 NB use of 526 gives 0.745566... and gets A0            2<sup>nd</sup> A1 for a correct equation for <math>t</math> in terms of <math>p</math> with <math>a</math> and <math>b</math> awrt 3sf An equn in <math>y</math> or <math>x</math> is A0</p> <p>M1 for clear use of their line (equation or on graph) and <math>p = 16</math> to estimate <math>t</math>.            This may be an expression or lines marked on the diagram            A1 for awrt 5.8, even if their line is not fully correct. Accept “<math>t &gt; 5.8</math>”(oe). Answer only 2/2</p>	

Question	Scheme	Marks
<p>4. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p><math>B, W</math> or <math>T, W</math> [ accept <math>B \cup T, W</math> or <math>B \cap T, W</math> ] [Condone <math>P(B), P(W)</math> etc]            Since there is no <u>overlap</u> between the events or cannot happen together (o.e.)            (Accept comment in context e.g. “no one walks and takes the train”)</p> <p>e.g. <math>P(B) = \frac{9}{25}, P(T) = \frac{8}{25}, P(B \cap T) = \frac{5}{25}</math>  <math>P(B \cap T) \neq P(B) \times P(T)</math> [<math>0.2 \neq 0.36 \times 0.32 = 0.1152</math> o.e.]            So <math>B</math> and <math>T</math> are <u>not</u> independent</p> <p><math>[P(W) =] \frac{7}{25}</math> or 0.28</p> <p><math>[P(B \cap T) =] \frac{5}{25}</math> or <math>\frac{1}{5}</math> or 0.2</p> <p><math>[P(T   B) =] \frac{P(T \cap B)}{P(B)} = \frac{\text{"(d)"}}{(5+4)/25}</math>  <math>= \frac{5}{9}</math> or 0.5<del>6</del></p>	<p>B1 B1 (2)</p> <p>M1 M1 A1cso (3)</p> <p>B1 (1)</p> <p>B1 (1)</p> <p>M1 A1 (2)</p> <p style="text-align: right;"><b>[9]</b></p>
<b>Notes</b>		
	<p>(a) 1<sup>st</sup> B1 for a suitable pair. Do not accept universally exclusive pairs such as <math>B</math> and <math>B'</math> etc            2<sup>nd</sup> B1 for any <u>correct</u> statement. Accept use of symbols e.g.: <math>B \cap W = \emptyset</math> or <math>P(T \cap W) = 0</math> etc            But <math>T \cap W = 0</math> is B0 (since it is not a correct statement)</p> <p>(b) 1<sup>st</sup> M1 for an attempt at all required probabilities with labels for a suitable test (allow one error).            Accept use of <math>A</math> and <math>B</math> as long as they can be identified as <math>B</math> and <math>T</math> by correct probabilities            Must be probabilities not integers such as 5, 9, 8 etc for both these M marks            2<sup>nd</sup> M1 for <math>P(B) \times P(T)</math> evaluated (correct for <u>their</u> probabilities)            or <math>P(B \cap T) \neq P(B) \times P(T)</math> stated or implied in symbols or using their probabilities.            or <math>P(B   T) \neq P(B)</math> or <math>P(T   B) \neq P(T)</math> stated or implied in symbols or using their probabilities.            A1 for a conclusion of <u>not</u> independent. Requires all probabilities used to be correct and seen.            This A mark is dependent on both Ms</p> <p>NB <math>P(B   T) = \frac{5}{8}</math> &amp; <math>P(B) = \frac{9}{25}</math> or <math>P(T   B) = \frac{5}{9}</math> &amp; <math>P(T) = \frac{8}{25}</math> seen, followed by a correct conclusion scores 3/3</p> <p>(e) M1 for a correct ratio of probabilities e.g. <math>\frac{5/25}{(5+4)/25}</math> or <math>\frac{5}{5+4}</math> or            A correct ratio expression and at least one correct (or correct f.t.) probability substituted.            A1 for <math>\frac{5}{9}</math> with no incorrect working seen but <math>\frac{5}{9}</math> following from <math>P(B   T)</math> is 0/2. <math>\frac{5}{9}</math> alone is 2/2</p>	

Question	Scheme	Marks
5. (a)	One large square = $\frac{450}{22.5}$ <u>or</u> one small square = $\frac{450}{562.5}$ (o.e. e.g. $\frac{562.5}{450}$ ) One large square = 20 cars <u>or</u> one small square = 0.8 cars <u>or</u> 1 car = 1.25 squares No. > 35 mph is: $4.5 \times 20$ <u>or</u> $112.5 \times 0.8$ (or equivalent e.g. using fd) $= \underline{90}$ (cars)	M1 A1 dM1 A1 (4)
(b)	$[\bar{x}] = \frac{30 \times 12.5 + 240 \times 25 + 90 \times 32.5 + 30 \times 37.5 + 60 \times 42.5}{450} \left[ = \frac{12975}{450} \right]$ $= 28.83... \text{ or } \frac{173}{6} \text{ awrt } \underline{28.8}$	M1 M1 A1 (3)
(c)	$[Q_2 =] 20 + \frac{195}{240} \times 10$ (o.e.) [Allow use of $(n + 1)$ giving 195.5 instead of 195] $= 28.125$ [Use of $(n + 1)$ gives 28.145...] <b>awrt <u>28.1</u></b>	M1 A1 (2)
(d)	$Q_2 < \bar{x}$ So <u>positive skew</u>	[Condone $Q_2 \approx \bar{x}$ ] [so (almost) <u>symmetric</u> ] B1ft dB1ft (2)
(e)	[If chose <u>skew</u> in (d)] <b>median (<math>Q_2</math>)</b> Since the data is skewed or median not affected by extreme values	[If chose <u>symmetric</u> in (d)] <b>mean (<math>\bar{x}</math>)</b> Since it uses all the data B1 dB1 (2)

[13]

**Notes**

(a)	1 <sup>st</sup> M1 for attempt to count squares (accept "22.5" in [22, 23] and "562.5" in [550, 575]) <b>and</b> use 450 to obtain a measure of scale. [If using fd must use 450 to obtain scale factor] 1 <sup>st</sup> A1 for a correct calc. for 20 or 0.8 or 1.25 etc [ May be fd = 4 to 1 large sq. or 0.8 to 1 small sq. May be on the diagram.] 2 <sup>nd</sup> dM1 dep on 1 <sup>st</sup> M1 for correctly counting squares for > 35 mph and forming suitable expr' 2 <sup>nd</sup> A1 for 90 with no incorrect working seen. e.g. $\frac{4.5}{22.5} \times 450$ scores M1A1M1 and A1 when = 90 is seen. Answer only is 4/4
(b)	1 <sup>st</sup> M1 for clear, sensible use of mid-points at least 3 of (12.5, 25, 32.5, 37.5, 42.5) seen 2 <sup>nd</sup> M1 for an expression for $\bar{x}$ (at least 3 correct terms on num' and a compatible denominator) Follow through their frequencies. You may see these fractions: $\frac{16218.75}{562.5}$ (small squares), $\frac{12975}{450}$ (frequencies), $\frac{648.75}{22.5}$ (large squares) A1 for awrt 28.8 (answer only is 3/3)
(c)	M1 for a full expression for median (using their frequencies). May see e.g. $25 + \frac{75}{120} \times 5$ etc Do not accept boundaries of 19.5 or 20.5, these are M0A0 A1 for awrt 28.1 (answer only is 2/2) [For use of $(n + 1)$ accept 28.15 but not 28.2]
(d)	1 <sup>st</sup> B1ft for a correct statement about their $Q_2$ and $\bar{x}$ [Condone $Q_2 \approx \bar{x}$ only if $ Q_2 - \bar{x}  < 1$ ] Do not accept an argument based on the shape of the graph alone. 2 <sup>nd</sup> dB1ft dependent on 1 <sup>st</sup> B1 for a <u>compatible</u> description of skewness. F.t. their values If $Q_1 = 23.4$ and $Q_3 = 33.7 \sim 33.8$ are seen allow comparison of quartiles for 1 <sup>st</sup> B1 in (d)
<b>Quartiles</b>	(e) 1 <sup>st</sup> B1 for a correct choice based on their skewness comment in (d). If no choice made in (d) only $Q_2$ 2 <sup>nd</sup> dB1 for a suitable compatible comment

Question	Scheme	Marks
<p>6. (a)</p> <p>(b)</p> <p>(c)</p>	$[z =] \pm \left( \frac{150 - 162}{7.5} \right)$ $[z =] - 1.6$ $[P(F > 150) = P(Z > -1.6) =] = 0.9452(0071\dots)$ $z = \pm 0.2533 \text{ (or better seen)}$ $(\pm) \frac{s - 162}{7.5} = 0.2533(47\dots)$ $s = 163.9$ $z = \pm 1.2816 \text{ (or better seen)}$ $\frac{162 - \mu}{9} = -1.2815515\dots$ $\mu = 173.533\dots$	<p>M1</p> <p>A1</p> <p>A1 (3) <b>awrt 0.945</b></p> <p>B1</p> <p>M1</p> <p>A1 (3) <b>awrt 164</b></p> <p>B1</p> <p>M1</p> <p>A1</p> <p>A1 (4) <b>awrt 174</b></p> <p style="text-align: right;"><b>[10]</b></p>
<b>Notes</b>		
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p><b>NB</b></p>	<p>M1 for attempting to standardise with 150, 162 and 7.5. Accept <math>\pm</math>                  Allow use of symmetry and therefore 174 instead of 150                  1<sup>st</sup> A1 for <math>-1.6</math> seen. Allow 1.6 seen if 174 used or awrt 0.945 is seen. Sight of 0.945(2) is A1.                  2<sup>nd</sup> A1 for awrt 0.945 Do not apply ISW, if 0.9452 is followed by <math>1 - 0.9452</math> then award A0                  Correct answer only 3/3</p> <p>B1 for <math>(z =) \pm 0.2533</math> (or better) seen.                  Giving <math>z = \pm 0.25</math> or <math>\pm 0.253</math> scores B0 here but may get M1A1                  M1 for standardising with <math>s</math> (o.e.), 162 and 7.5, allow <math>\pm</math>, and setting equal to a <math>z</math> value                  Only allow <math>0.24 \leq z \leq 0.26</math> Condone e.g. 160 for 162 etc                  A1 for awrt 164 (Correct answer only scores B0M1A1)</p> <p>B1 for <math>(z =) \pm 1.2816</math> (or better) seen. Allow awrt <math>\pm 1.28</math> if B0 scored in (b) for <math>z =</math> awrt <math>\pm 0.25</math>                  M1 for attempting to standardise with 162, 9 and <math>\mu</math>, and setting equal to a <math>z</math> value where  <math>1.26 &lt;  z  &lt; 1.31</math>. Allow <math>\pm</math> here so signs don't have to be compatible.                  1<sup>st</sup> A1 for a correct equation <u>with</u> compatible signs and <math>1.26 &lt;  z  &lt; 1.31</math>                  2<sup>nd</sup> A1 for awrt 174 (Correct answer only scores B0M1A1A1). <b>Dependent on 1<sup>st</sup> A1</b></p> <p>An equation <math>\frac{162 - \mu}{9} = 1.2816</math> leading to an answer of <math>\mu = 174</math> is A0A0 <u>unless</u> there is clear                  correct working such as: <math>\frac{162 - x}{9} = 1.2816 \Rightarrow x = \dots \therefore \mu = 162 + (162 - x) = 174</math> then award A1A1</p> <p>A common error is: <math>\frac{162 - \mu}{9} = 1.2816</math> followed by <math>\mu = 162 + 9 \times 1.2816 =</math> awrt 174 It gets                  A0A0</p>	

Question	Scheme	Marks
<p>7. (a)</p> <div style="text-align: center;"> </div> <p>(b) <math>P(\text{Exactly one defect}) = 0.03 \times 0.3 + 0.97 \times 0.02</math> or <math>P(PS \cup Split) - 2P(PS \cap Split)</math>  <math>= [0.009 + 0.0194 = ]</math> <b>0.0284</b></p> <p>(c) <math>P(\text{No defects}) = (1 - 0.03) \times (1 - 0.02) \times (1 - 0.05)</math> (or better)  <math>= 0.90307</math> <b>awrt 0.903</b></p> <p>(d) <math>P(\text{Exactly one defect}) = (b) \times (1 - 0.05) + (1 - 0.03) \times (1 - 0.02) \times 0.05</math>  <math>= "0.0284" \times 0.95 + 0.97 \times 0.98 \times 0.05</math>  <math>= [0.02698 + 0.04753] = 0.07451</math> <b>awrt 0.0745</b></p>	<p>Shape B1</p> <p>Labels &amp; 0.03 B1</p> <p>Labels &amp; 0.7, 0.02 B1</p> <p>(3)</p> <p>M1A1ft A1 cao (3)</p> <p>M1 A1 cao (2)</p> <p>M1 M1 A1ft A1 cao (4)</p> <p><b>[12]</b></p>	
<b>Notes</b>		
<p>(a)</p> <p>(b)</p> <p><b>MR</b></p> <p>(c)</p> <p>(d)</p> <p><b>MR</b></p>	<p><b>Allow MR of 0.2 for 0.02 or 0.3 for 0.03 on tree diagram to score all M and A1ft marks only</b></p> <p>1<sup>st</sup> B1 for 2 branch then 4 branch shape                  2<sup>nd</sup> dB1 dep. on 1<sup>st</sup> B1 for labels showing stitching (accept letters) and 0.03 value correctly placed                  3<sup>rd</sup> dB1 dep. on 1<sup>st</sup> B1 for labels showing splitting and 0.7 and 0.02 correctly placed                  [probabilities shown in brackets are <u>not</u> required and any such values given can be ignored in (a)]</p> <p>M1 for <math>0.03 \times p + 0.02 \times q</math> where <math>p</math> and <math>q</math> follow from their tree diagram. Extra terms is M0                  1<sup>st</sup> A1ft for a fully correct expression. Accept <math>1 - 0.7</math> for 0.3 and <math>1 - 0.03</math> for 0.97                  Follow through 0.2 and 0.3 MR only</p> <p>0.2 for 0.02 <math>\rightarrow 0.203</math> or 0.3 for 0.03 <math>\rightarrow 0.104</math> or both <math>\rightarrow 0.23</math> should score M1A1A0                  2<sup>nd</sup> A1 cao for 0.0284 only (or exact equivalent such as <math>\frac{71}{2500}</math>)</p> <p><b>Do not allow 0.5 as MR of 0.05 so no M or A marks in (c) or (d)</b></p> <p>M1 for <math>(\text{their } 0.97) \times (\text{their } 0.98) \times (1 - 0.05)</math> (or better) f.t. values from their tree diagram                  A1 cao for awrt 0.903</p> <p>1<sup>st</sup> M1 for one correct triple (or correct ft from their tree) of:  <math>[0.03 \times 0.3 \times (1 - 0.05)] + [0.97 \times 0.02 \times (1 - 0.05)] + [0.97 \times 0.98 \times 0.05]</math></p> <p>2<sup>nd</sup> M1 for two correct triples or correct ft from their tree and adding <u>or</u> their (b) <math>\times (1 - 0.05)</math>                  1<sup>st</sup> A1ft for a fully correct expression or f.t. their (b) and 0.2 or 0.3 MR only</p> <p>0.2 for 0.02 <math>\rightarrow 0.23165</math> or 0.3 for 0.03 <math>\rightarrow 0.1331</math> or both <math>\rightarrow 0.2465</math> (or awrt 3sf) scores M1M1A1A0                  2<sup>nd</sup> A1 cao for awrt 0.0745</p>	



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UA033137 Summer 2012

For more information on Edexcel qualifications, please visit our website  
[www.edexcel.com](http://www.edexcel.com)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual  




Llywodraeth Cynulliad Cymru  
Welsh Assembly Government



Rewarding Learning



# Mark Scheme (Results)

January 2013

GCE Maths – Statistics S1 (6683/01)

## Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk) for our BTEC qualifications.

Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

If you have any subject specific questions about this specification that require the help of a subject specialist, you can speak directly to the subject team at Pearson.

Their contact details can be found on this link: [www.edexcel.com/teachingservices](http://www.edexcel.com/teachingservices).

You can also use our online Ask the Expert service at [www.edexcel.com/ask](http://www.edexcel.com/ask). You will need an Edexcel username and password to access this service.

### **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2013

Publications Code UA034849

All the material in this publication is copyright

© Pearson Education Ltd 2013

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.
- Unless indicated in the mark scheme a correct answer with no working should gain full marks for that part of the question.

## EDEXCEL GCE MATHEMATICS

### General Instructions for Marking

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

In some instances, the mark distributions (e.g. M1, B1 and A1) printed on the candidate's response may differ from the final mark scheme.

### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes and can be used.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\checkmark$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - $\square$  The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but incorrect answers should never be awarded A marks.
  5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.

6. If a candidate makes more than one attempt at any question:
  - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
  
7. Ignore wrong working or incorrect statements following a correct answer.
  
8. The maximum mark allocation for each question/part question(item) is set out in the marking grid and you should allocate a score of '0' or '1' for each mark, or "trait", as shown:

	0	1
aM		•
aA	•	
bM1		•
bA1	•	
bB	•	
bM2		•
bA2		•

**January 2013  
6683 Statistics S1  
Mark Scheme**

Question Number	Scheme	Marks
<b>1.</b>		
(a)	$(S_{tt}) = 8702 - \frac{258^2}{10} \quad \text{or} \quad (S_{gt}) = 1550.2 - \frac{258 \times 63.6}{10}$ $(S_{tt} =) 2045.6, \quad (S_{gt} =) -90.68 \quad \text{awrt (2046),} \quad \text{awrt } -90.7$	M1 A1, A1 (3)
(b)	$r = \frac{-90.68}{\sqrt{2045.6 \times 7.864}} = -0.714956... \quad \text{awrt } -0.715$	M1 A1 (2)
(c)	Positive e.g. high $v$ corresponds to low $t$ and low $t$ corresponds to high $g$ so expect high $v$ to corresponds to high $g$ <u>or</u> expect more revision to result in a better grade	B1 B1 (2)
<b>Notes</b>		
(a)	M1 for at least one correct expression 1 <sup>st</sup> A1 for $S_{tt} =$ awrt 2046 (Condone $S_{xx} = \dots$ or even $S_{yy} = \dots$ ) 2 <sup>nd</sup> A1 for $S_{gt} =$ awrt -90.7 (Condone $S_{xy} = \dots$ )	
(b)	M1 for attempt at correct formula. Must have their $S_{tt}$ , $S_{gt}$ and given $S_{gg}$ in the correct places. Condone missing “-” Award M1A0 for awrt -0.71 with no expression seen M0 for $\frac{1550.2}{\sqrt{8702 \times 7.864}}$ Correct answer only is 2/2	
(c)	1 <sup>st</sup> B1 for saying "positive". Ignore mention of skew. 2 <sup>nd</sup> B1 for suitable reason that mentions at least $v$ and $g$ and supports positive correlation. e.g. “the less <u>revision</u> done the lower the <u>grade</u> ” is B1 “should do better with more <u>revision</u> ” is B0 since does not mention <u>grades</u> “both coefficients are similar” or two sketches of negative correlation with labelled axes is B1 since $v$ , $t$ and $g$ are implied Allow use of letters $v$ and $g$ Allow equivalent terms e.g. “study” instead of “revision” or “score” instead of “grade”	
<b>7</b>		


Question Number	Scheme	Marks								
<p><b>2.</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p>	<p>F(3) = 1 gives <math>\frac{3^3 + k}{40} = 1</math>                      So <math>k = \underline{13}</math></p> <p><math>P(X = 1) = \frac{14}{40}</math> or 0.35 (o.e.)                      Use of <math>P(X = 2) = F(2) - F(1)</math> or <math>P(X = 3) = F(3) - F(2)</math>  <math>P(X = 2) = \frac{7}{40}</math> or 0.175, <math>P(X = 3) = \frac{19}{40}</math> or 0.475</p> <p><math>\text{Var}(4X - 5) = 4^2 \text{Var}(X)</math>                      So <math>\text{Var}(4X - 5) = \frac{259}{20}</math> or 12.95</p>	<p>M1 A1cso (2)</p> <p>B1 M1 A1, A1 (4)</p> <p>M1 A1 (2)</p> <p><b>8</b></p>								
<b>Notes</b>										
<p>(a)</p> <p><b>Verify</b></p> <p>(b)</p> <p>(c)</p>	<p>M1 for use of <math>F(3) = 1</math> Attempt at <math>\frac{3^3 + k}{40} = 1</math> must be seen  <math>27 + k = 40</math> without reference to <math>F(3) = 1</math> is M0                      A1cso for no incorrect working seen and M1 scored.</p> <p>Allow M1 for <math>\frac{3^3 + 13}{40} = 1</math> but the A1 requires an <u>explicit</u> comment such as “so <math>k = 13</math>”</p> <p>If a table such as this is seen then award B1M1A1A1. Ignore labels on 2<sup>nd</sup> row</p> <table border="1" data-bbox="480 1205 1150 1330"> <tr> <td></td> <td style="text-align: center;">1</td> <td style="text-align: center;">2</td> <td style="text-align: center;">3</td> </tr> <tr> <td></td> <td style="text-align: center;"><math>\frac{7}{20}</math> or 0.35</td> <td style="text-align: center;"><math>\frac{7}{40}</math> or 0.175</td> <td style="text-align: center;"><math>\frac{19}{40}</math> or 0.475</td> </tr> </table> <p>Otherwise apply the following:</p> <p>B1 for <math>\frac{14}{40}</math> or 0.35 or any exact equivalent. Can be labelled <math>F(1)</math>, <math>P(X = 1)</math> or <math>p(x)</math> and associated with <math>x = 1</math> or given in a table but must have a label.</p> <p>M1 for clear method showing how to obtain <math>P(X = \dots)</math> from <math>F(x)</math>                      M1 can be implied if either <math>P(X = 2)</math> or <math>P(X = 3)</math> is correct</p> <p>1<sup>st</sup> A1 for <math>P(X = 2) = \frac{7}{40}</math> or 0.175 or exact equivalent</p> <p>2<sup>nd</sup> A1 for <math>P(X = 3) = \frac{19}{40}</math> or 0.475 or exact equivalent</p> <p>M1 for correct use of the variance formula (<math>4^2 \text{Var}(X)</math> alone secures M1)                      A value for <math>\text{Var}(X)</math> is not required for this M1</p> <p>A1 for any exact equivalent to 12.95 Correct answer only is 2/2</p>		1	2	3		$\frac{7}{20}$ or 0.35	$\frac{7}{40}$ or 0.175	$\frac{19}{40}$ or 0.475	
	1	2	3							
	$\frac{7}{20}$ or 0.35	$\frac{7}{40}$ or 0.175	$\frac{19}{40}$ or 0.475							

Question Number	Scheme	Marks
<p><b>3.</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p><math>\sum t = 140</math> (or <math>\bar{t} = 17.5</math>) and <math>\sum m = 32</math> (or <math>\bar{m} = 4</math>)</p> <p><math>(S_{tm}) = 469.5 - \frac{"140" \times "32"}{8}</math></p> <p><math>(S_{tm} =) -90.5</math></p> <p><math>b = \frac{S_{tm}}{S_{tt}} = \frac{-90.5}{354}</math></p> <p><math>b = -0.255649\dots</math> (allow <math>\frac{181}{708}</math>)      <math>-0.25</math> or awrt <math>-0.26</math></p> <p><math>a = \frac{"32"}{8} - b \times \frac{"140"}{8}</math></p> <p>So equation of the line is <b><math>m = 8.47 - 0.256t</math></b> (allow <math>m = \frac{11999}{1416} - \frac{181}{708}t</math>)</p> <p><math>(8.47 - 0.256 \times 10 =) 5.9\dots</math>      awrt <b><u>5.9</u></b></p> <p>Should be reliable since 10 is in the range (of the data )</p>	<p>B1 B1</p> <p>M1</p> <p>A1cso</p> <p>(4)</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>(4)</p> <p>B1</p> <p>(1)</p> <p>B1</p> <p>(1)</p> <p><b>10</b></p>
<b>Notes</b>		
	<p>(a) 1<sup>st</sup> B1 for 140 seen in correct context or correctly labelled                  2<sup>nd</sup> B1 for 32 seen in correct context or correctly labelled.                  (allow a fully correct expression – not “...+...+...”)                  4480 used correctly is B1B1                  M1 for attempting a correct expression. Follow through their 140 and their 32                  You may see attempt at <math>\sum (t - \bar{t})(m - \bar{m})</math>. This must have all the products seen.                  A1cso requires a correct expression seen and no incorrect working leading to <math>-90.5</math></p> <p>(b) 1<sup>st</sup> M1 for a correct expression for <math>b</math>. Follow through their <math>S_{tm}</math>. Condone missing “–”                  1<sup>st</sup> A1 for awrt <math>-0.26</math> or condone <math>-0.25</math>                  2<sup>nd</sup> M1 for a correct method for <math>a</math>. Follow through their sums from part (a) and their value of <math>b</math>                  2<sup>nd</sup> A1 for a correct equation for <math>m</math> and <math>t</math> with <math>a =</math> awrt <math>8.47</math> and <math>b =</math> awrt <math>-0.256</math>                  Must be an equation in <math>m</math> and <math>t</math>, use of <math>x</math> or <math>y</math> scores A0 here.</p> <p>(c) B1 for awrt <math>5.9</math> Accept 6 if the correct expression (awrt <math>8.47 - 10 \times</math> awrt <math>0.256</math>) is seen</p> <p>(d) B1 for suggesting it is reliable and mentioning 10 within the range (of the data.)                  or suggesting it is reliable since <u>interpolating</u> or <u>not extrapolating</u></p> <p>NB “it is reliable since <u>it</u> is in the range” is B0 since “<u>it</u>” is not explicit enough                  Condone extra non-relevant comments but penalise contradictory comments.                  e.g. “near the extreme so <u>not reliable</u> but not extrapolated so reliable” is B0 since <u>contradicts</u>                  “reliable since 10 is within the range (of temps) <u>and 5.9 within range of times</u>” is B1 since <u>irrelevant</u></p>	

Question Number	Scheme	Marks
<p><b>4.</b></p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p><b>S.C.</b></p>	$\frac{127-100}{15}$ <p>So <math>P(L &gt; 127) = P(Z &gt; 1.8)</math> or <math>1 - P(Z &lt; 1.8)</math> o.e.  <math>= 1 - 0.9641 = \underline{\mathbf{0.0359}}</math> (awrt <b>0.0359</b>)</p> $\frac{d-100}{15} = -1.2816 \quad (\text{Calculator gives } -1.2815515\dots)$ $d = 80.776 \quad (\text{awrt } \underline{\mathbf{80.8}})$ <p>Require <math>P(L &gt; 133   L &gt; 127)</math></p> $= \frac{P(L > 133)}{P(L > 127)} = \frac{P(Z > 2.2)}{P(L > 127)}$ $= \frac{1-0.9861}{1-0.9641} = \frac{0.0139}{0.0359}$ $= 0.3871\dots = \text{awrt } \underline{\mathbf{0.39}}$ <p>An attempt at <math>P(L &lt; 133   L &gt; 127)</math> that leads to awrt 0.61 (M0M1A0A0)</p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>(3)</p> <p>M1, B1</p> <p>A1</p> <p>(3)</p> <p>M1</p> <p>dM1</p> <p>A1</p> <p>A1</p> <p>(4)</p> <p><b>10</b></p>
<b>Notes</b>		
<p>(a)</p> <p>(b)</p> <p><b>Calc</b></p> <p>(c)</p> <p><b>ALT</b></p>	<p>M1 for attempting to standardise with 127, 100 and 15. Allow <math>\pm</math></p> <p>1<sup>st</sup> A1 for <math>Z &gt; 1.8</math>. Allow a diagram but must have 1.8 and correct area indicated. Must have the Z so <math>P(L &gt; 127)</math> with or without a diagram is insufficient. May be implied by 0.0359</p> <p>2<sup>nd</sup> A1 for awrt 0.0359 (calc. gives 0.035930266...). Correct ans only 3/3. M1A0A1 not poss.</p> <p>M1 for an attempt to standardise with 100 and 15 and set <math>= \pm</math> any z value (<math> z  &gt; 1</math>)</p> <p>B1 for <math>z = \pm 1.2816</math> (or better) seen anywhere [May be implied by 80.776(72...) or better seen]</p> <p>A1 for awrt 80.8 (can be scored for using 1.28 but then they get M1B0A1)</p> <p>The 80.8 must follow from correct working.</p> <p>If answer is awrt 80.8 <b>and</b> awrt 80.777 or 80.776... or better seen then award M1B1A1</p> <p>If answer is awrt 80.8 or 80.77 then award M1B0A1 (unless of course <math>z = 1.2816</math> is seen)</p> <p>1<sup>st</sup> M1 for clear indication of correct conditional probability or attempt at correct ratio</p> <p>So clear attempt at <math>\frac{P(L &gt; 133)}{P(L &gt; 127)}</math> is sufficient for the 1<sup>st</sup> M1</p> <p>2<sup>nd</sup> dM1 dependent on 1<sup>st</sup> M1 for <math>P(L &gt; 133)</math> leading to <math>P(Z &gt; 2.2)</math>.</p> <p>1<sup>st</sup> A1 for 0.0139 or better seen coming from <math>P(Z &gt; 2.20)</math>. Dependent on both Ms</p> <p>2<sup>nd</sup> A1 for awrt 0.39. Both Ms required</p> <p>If they assume Alice did not check that the phone was working you may see:  <math>[P(L &lt; 127).0] + P(L &gt; 127).P(L &gt; 133   L &gt; 127)</math> Provided the <u>conditional probability</u> is seen as part of this calculation the 1<sup>st</sup> M1 can be scored and their final answer will be 0.0139(4/4)</p> <p>An answer of 0.0139 without sight of the conditional probability is 0/4.</p>	

Question Number	Scheme	Marks
<p>5. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p>	<p>Width = 4 (cm)                      Area of 14 cm<sup>2</sup> represents frequency 28 and area of 4h represents 18                      Or <math>\frac{4h}{18} = \frac{14}{28}</math> (o.e.) <span style="float: right;"><math>h = \underline{2.25}</math> (cm)</span></p> <p><math>m = (240) + \frac{10}{22} \times 80</math> (o.e.)  <math>= 276.36... \left(\frac{3040}{11}\right)</math> <span style="float: right;">((£)<u>276</u> ≤ m &lt; (£)276.5)</span></p> <p><math>\sum fy = 31600</math> leading to <math>\bar{y} = 316</math></p> <p><math>\sigma_y = \sqrt{\frac{12452800}{100} - (\bar{y})^2} = 157.07...</math> (awrt <u>157</u>) Allow s = 157.86...</p> <p>Skewness = 0.764... (awrt <u>0.76</u> or <u>0.75</u>)                      [If n+1 used in (b) and m = £278 accept awrt 0.73 or 0.72]  <u>Positive</u> skew</p> <p><math>z = \pm \frac{80}{150}</math></p> <p>P(240 &lt; X &lt; 400) = <u>0.40 ~ 0.41</u></p> <p>(e) suggests a reasonable fit for this range BUT                      (d) since skew it will not be a good fit overall</p>	<p>B1 M1 A1 (3)</p> <p>M1 A1 (2)</p> <p>M1A1 M1A1 (4)</p> <p>B1 B1ft (2)</p> <p>M1 A1 (2)</p> <p>B2/1/0 (2)</p> <p><b>15</b></p>
<b>Notes</b>		
	<p>(a) B1 for width (ignore units)                      M1 for clear method using area and frequency <u>or</u> their width × their height = 9                      e.g. seeing both fd of 0.7 and 0.225 (may see fd in the table) [Must use correct interval]</p> <p>(b) M1 for <math>\frac{10}{22} \times 80</math> or <math>\frac{10.5}{22} \times 80</math> (o.e.). Allow use of (n + 1) leading to £278.18... or [278, 278.5]                      A1 Do not award if incorrect end-point seen but answer only is 2/2</p> <p>(c) 1<sup>st</sup> M1 attempt at <math>\sum fy</math> with at least 3 correct products or ans. that rounds to 30 000 (to 1 sf) &amp;/100                      2<sup>nd</sup> M1 for correct expression including <math>\sqrt{\quad}</math>. Follow through <math>\bar{y}</math>. Need <math>\sum fy^2</math> correct but condone a minor transcription error e.g. 12458200.</p> <p>(d) 1<sup>st</sup> B1 for awrt 0.76/0.75 for m = £276 or awrt 0.73/0.72 for m = £278                      2<sup>nd</sup> B1ft for a correct description of their skew based on their measure <u>or</u> if no measure given based on their values of mean and median. (correlation is B0)</p> <p>(e) M1 for an attempt to standardise using the 320 and 150 and either 240 or 400 (implied by 0.53)                      A1 for answer in range [0.40, 0.41] (tables gives 0.4038, calculator 0.40619...) Ans only 2/2</p> <p>(f) For B2 we need 2 comments that make reference to each of part (e) and part (d)                      One comment should suggest it is <u>not</u> good since <u>skew</u>. The other it <u>is</u> since matches <u>range in (e)</u>                      1<sup>st</sup> B1 for one relevant comment                      2<sup>nd</sup> B1 for both comments NB Do not use B0B1</p>	

Question Number	Scheme	Marks							
<p><b>6. (a)</b></p> <table border="1" data-bbox="272 275 638 398"> <tr> <td><math>b</math></td> <td>1</td> <td>3</td> <td>5</td> </tr> <tr> <td><math>P(B = b)</math></td> <td><math>\frac{1}{3}</math></td> <td><math>\frac{1}{3}</math></td> <td><math>\frac{1}{3}</math></td> </tr> </table> <p><b>(b)</b> Discrete Uniform {distribution}</p> <p><b>(c)</b> <math>[E(B) = ]</math> 3 (by symmetry)</p> <p><b>(d)</b> <math>[E(R) = ]</math> <math>2 \times \frac{2}{3} + 4 \times \frac{1}{6} + 6 \times \frac{1}{6}</math> <math>= \underline{3}</math></p> <p><b>(e)</b> <math>[E(R^2) = ]</math> <math>2^2 \times \frac{2}{3} + 4^2 \times \frac{1}{6} + 6^2 \times \frac{1}{6}</math> <math>\left[ = \frac{34}{3} \right]</math> <math>[\text{Var}(R) = ]</math> <math>\frac{34}{3} - 3^2 = \frac{7}{3}</math> (or any exact equivalent. NB 2.33 is A0)</p> <p><b>(f)</b> Coin lands on <b>2</b>, choose <b>blue</b> die; coin lands on <b>5</b> choose <b>red</b> die <math>P(\text{Avisha wins}) = \frac{1}{2} \times \left( \frac{1}{3} + \frac{1}{3} \right) + \frac{1}{2} \times \frac{1}{6}</math> <math>= \frac{5}{12}</math> (allow awrt 0.417)</p>	$b$	1	3	5	$P(B = b)$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$	<div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;">                 Also allow <math>b</math> values 1,1,3,3,5,5 and probabilities all <math>\frac{1}{6}</math> </div> <p>B1 B1 (2)</p> <p>B1 (1)</p> <p>B1 (1)</p> <p>M1 A1 (2)</p> <p>M1</p> <p>dM1, A1 (3)</p> <p>B2/1/0 M1 A1 (4)</p> <p style="text-align: right;"><b>13</b></p>
$b$	1	3	5						
$P(B = b)$	$\frac{1}{3}$	$\frac{1}{3}$	$\frac{1}{3}$						
<b>Notes</b>									
	<p><b>(a)</b> 1<sup>st</sup> B1 for correctly identifying values of <math>b</math> as 1, 3, 5 or 1,1,3,3,5,5 2<sup>nd</sup> B1 for probabilities all <math>= \frac{1}{3}</math> or exact equivalent (or of course 6 cases of <math>\frac{1}{6}</math>) Any correct probability distribution or probability function is 2/2. Must be in part (a)</p> <p><b>(b)</b> B1 for "Discrete Uniform". Both words required.</p> <p><b>(c)</b> B1 for answer of 3 o.e. Accept <math>E(X) = 3</math></p> <p><b>(d)</b> M1 for an attempt at correct formula. At least 2 correct products seen. If later divide by <math>n (\neq 1)</math> M0 A1 for an answer of 3. Correct answer only scores both marks.</p> <p><b>(e)</b> 1<sup>st</sup> M1 for a correct attempt at <math>E(R^2)</math>. At least 2 correct products seen. Condone <math>\text{Var}(R) = \text{etc}</math> May be implied by sight of <math>\frac{34}{3}</math> or 11.3 or better. 2<sup>nd</sup> dM1 Dep. on 1<sup>st</sup> M1 for clear attempt at <math>E(R^2) - [E(R)]^2</math> Must see their values <u>used</u>. NB <math>\text{Var}(R) = E(R^2) - [E(R)]^2 = \frac{34}{3} - 3^2</math> is M1M0A0 since do not <u>use</u> their <math>[E(R)]^2</math></p> <p><b>(f)</b> B2/1/0 Both correct B1B1, one correct B1B0. Do not use B0B1 [e.g. always red or RR is B1B0] NB Allow other descriptions of the die e.g. 1<sup>st</sup> or fair for blue, 2<sup>nd</sup> for red if they are clear. M1 for evaluating correct probabilities i.e. only <math>\frac{1}{3}, \frac{1}{12}</math> seen <u>or</u> if incorrect choice made: M1 for an answer of : if choose RR (<math>\frac{1}{4}</math>), if choose BB (<math>\frac{1}{3}</math>), if choose RB (<math>\frac{1}{6}</math>) NB <math>\frac{5}{12}</math> as answer scores M1A1. Need to see choices of die stated for B marks.</p>								

Question Number	Scheme	Marks
<p>7.</p> <p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p>	<p><math>P(A \cup B) = 0.35 + 0.45 - 0.13</math> <u>or</u> <math>0.22 + 0.13 + 0.32</math>  <math>= \underline{0.67}</math></p> <p><math>P(A'   B') = \frac{P(A' \cap B')}{P(B')}</math> <u>or</u> <math>\frac{0.33}{0.55}</math>  <math>= \frac{3}{5}</math> <u>or</u> <math>0.6</math></p> <p><math>P(B \cap C) = 0.45 \times 0.2</math>  <math>= \underline{0.09}</math></p>  <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p>Allow 1<sup>st</sup> B1 for 3 intersecting circles in a box with zeros in the regions for <math>A \cap C</math>              Do not accept "blank" for zero</p> </div> <p><math>P(B \cup C)' = 0.22 + \underline{0.22}</math> <u>or</u> <math>1 - [0.56]</math> <u>or</u> <math>1 - [0.13 + 0.23 + 0.09 + 0.11]</math> o.e.  <math>= \underline{0.44}</math></p>	<p>M1 A1 (2)</p> <p>M1 A1 (2)</p> <p>M1 A1 (2)</p> <p>B1 B1ft B1 B1 (4)</p> <p>M1 A1 (2)</p> <p style="text-align: right;"><b>12</b></p>
<b>Notes</b>		
<p>NB May see Venn diagram for <math>A</math> and <math>B</math> only used for (a) and (b) but M marks are awarded for <u>correct expressions only</u>. No fit from an incorrect diagram for M marks.</p> <p>(a) M1 for attempt to use the addition rule. Correct substitution i.e. correct expression seen              A1 for 0.67 only. Correct answer only scores 2/2</p> <p>(b) M1 for a correct ratio of probabilities or a correct formula and at least one correct prob              For a correct formula allow "1 – their (a)" instead of 0.33 but not for correct ratio case.              Do not award for assuming independence i.e. <math>\frac{P(A' \cap B')}{P(B')} = \frac{0.65 \times 0.55}{0.55}</math> is M0. M0 if num &gt; denom              A1 for 3/5 or any exact equivalent.</p> <p>(c) M1 for correct expression. Need correct values for <math>P(B)</math> and <math>P(C)</math> seen.              A1 for 0.09 or any exact equivalent. Correct answer only is 2/2</p> <p>(d) <b>No labels <math>A, B, C</math> in (d) loses 1<sup>st</sup> B1 but can score the other 3 by implication</b>              B1 for box with <math>B</math> intersecting <math>A</math> and <math>C</math> but <math>C</math> not intersecting <math>A</math>. No box is B0              B1ft for 0.13 and their 0.09 in correct places. [ft <math>P(B \cap C)</math> from (c)]              B1 for any 2 of 0.22, <u>0.22</u>, 0.11 and 0.23 correct              B1 for all 4 values correct</p> <p>(e) M1 for a correct expression or follow through from their Venn diagram              NB <math>P(B') \times P(C') = 0.55 \times 0.8</math> is OK. Do not fit "blank" for zero and M0 for negative probs.              A1 for 0.44 only. Correct answer only is 2/2</p>		

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467  
Fax 01623 450481  
Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)  
Order Code UA034849 January 2013

For more information on Edexcel qualifications, please visit our website  
[www.edexcel.com](http://www.edexcel.com)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual  




Llywodraeth Cynulliad Cymru  
Welsh Assembly Government





Mark Scheme (Results)

Summer 2013

GCE Statistics 1 (6683/01R)

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2013

Publications Code UA036996

All the material in this publication is copyright

© Pearson Education Ltd 2013

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

**EDEXCEL GCE MATHEMATICS****General Instructions for Marking**

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes:

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\checkmark$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - $\square$  The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
  5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
  6. If a candidate makes more than one attempt at any question:
    - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
    - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
  7. Ignore wrong working or incorrect statements following a correct answer.
  8. In some instances, the mark distributions (e.g. M1, B1 and A1) printed on the candidate's response may differ from the final mark scheme

Question	Scheme	Marks
<p><b>1. (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p>	$b = \frac{18.35}{312.1} [= 0.058795\dots]$ $a = 5.8 - "0.058795\dots" \times 4.8$ <p style="text-align: right;"><b><u>a = awrt 5.52</u></b></p> <p>So <b><u>y = 5.52 + 0.0588x</u></b></p> $\frac{e}{10} = "5.52" + "0.0588" \times \left( \frac{g-60}{4} \right)$ $4e = 220.71 + 0.588(g-60)$ <p style="text-align: right;"><b><u>e = 46 + 0.15g</u></b></p> $e = "46" + "0.15" \times 100$ <p style="text-align: right;"><b>= <u>61</u></b></p>	<p>M1</p> <p>M1</p> <p>A1</p> <p>A1 (4)</p> <p>M1</p> <p>dM1</p> <p>A1A1 (4)</p> <p>M1</p> <p>A1 (2)</p> <p><b>[10]</b></p>
<b>Notes</b>		
<p><b>(a)</b></p> <p><b>(b)</b></p> <p><b>ALT</b></p> <p><b>(c)</b></p>	<p>1<sup>st</sup> M1 for a correct expression for <math>b</math></p> <p>2<sup>nd</sup> M1 for a correct expression for <math>a</math> – ft their value of <math>b</math></p> <p>1<sup>st</sup> A1 for <math>a = \text{awrt } 5.52</math></p> <p>2<sup>nd</sup> A1 for a correct equation in <math>y</math> and <math>x</math> with <math>a</math> and <math>b</math> correct to awrt 3 sf</p> <p>1<sup>st</sup> M1 for substitutions into <u>their</u> equation to get an equation in <math>e</math> and <math>g</math>. Need <math>y = \frac{e}{10}</math> and <math>x = \frac{g-60}{4}</math></p> <p>2<sup>nd</sup> dM1 Dep. on 1<sup>st</sup> M1 for an attempt to simplify (at least removing fractions). Allow one slip</p> <p>1<sup>st</sup> A1 for an equation <math>e = \text{awrt } 46 \pm \dots</math></p> <p>2<sup>nd</sup> A1 for an equation <math>e = \dots + \text{awrt } 0.15g</math></p> <p>1<sup>st</sup> M1 for use of <math>d = \frac{10 \times \text{"their } b\text{"}}{4}</math> or sight of 0.15 used as gradient</p> <p>2<sup>nd</sup> dM1 Dep. on 1<sup>st</sup> M1 for use of <math>\bar{e} = 10 \times \text{"their } \bar{y}\text{"}</math> or sight of 58 and use of <math>\bar{g} = 4 \times \text{"their } \bar{x}\text{"} + 60</math> or sight of 79.2 and use of these values to find <math>c</math> in <math>c = \bar{e} - d\bar{g}</math></p> <p>M1 for substituting <math>g = 100</math> into their new equation (or <math>x = 10</math> and then attempting to <math>\times</math> ans.by 10)</p> <p>A1 for awrt 61</p>	

Question	Scheme			Marks								
<p><b>2. (a)</b></p>	<table border="1"> <tr> <td><math>x</math></td> <td><b>1</b></td> <td><b>2</b></td> <td><b>3</b></td> </tr> <tr> <td><math>P(X = x)</math></td> <td><b>0.4</b></td> <td>0.25</td> <td>0.35</td> </tr> </table>	$x$	<b>1</b>	<b>2</b>	<b>3</b>	$P(X = x)$	<b>0.4</b>	0.25	0.35			
	$x$	<b>1</b>	<b>2</b>	<b>3</b>								
$P(X = x)$	<b>0.4</b>	0.25	0.35									
$P(X = 2) = F(2) - F(1)$ (o.e.)		$P(X = 2) = \underline{\underline{0.25}}$ $P(X = 3) = \underline{\underline{0.35}}$	M1 A1 A1 (3)									
<p><b>(b)</b></p>	$[F(1.8) = P(X \leq 1.8) = P(X \leq 1) = ] \underline{\underline{0.4}}$			B1 (1)								
<b>Notes</b>												
<p><b>(a)</b></p>	M1 for $P(X = 1) = 0.4$ <b>and</b> evidence of a correct method for finding $P(X = 2)$ or $P(X = 3)$ . Implied by correct ans. 1 <sup>st</sup> A1 for $P(X = 2) = 0.25$ 2 <sup>nd</sup> A1 for $P(X = 3) = 0.35$											
<p><b>(b)</b></p>	B1 for 0.4											

Question	Scheme	Marks
<p><b>3. (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p> <p><b>(e)</b></p>	<p>Width = <math>2 \times 1.5 = \underline{\mathbf{3 \text{ (cm)}}}</math>                      Area = <math>8 \times 1.5 = 12 \text{ cm}^2</math> Frequency = 24 so <math>1 \text{ cm}^2 = 2 \text{ plants}</math> (o.e.)                      Frequency of 12 corresponds to area of 6 so height = <math>\underline{\mathbf{2 \text{ (cm)}}}</math></p> <p><math>[Q_2 =] (5+) \frac{19}{24} \times 5</math>      <b>or</b>    (use of <math>(n+1)</math>) <math>(5+) \frac{19.5}{24} \times 5</math>                      = 8.9583...      <b>awrt 8.96</b>      <b>or</b>      9.0625...    awrt 9.06</p> <p><math>[\bar{x} =] \frac{755}{70}</math> or <b>awrt 10.8</b></p> <p><math>[\sigma_x =] \sqrt{\frac{12037.5}{70} - \bar{x}^2} = \sqrt{55.6326...}</math>                      = <b>awrt 7.46</b>    (Accept <math>s =</math> awrt 7.51)</p> <p><math>\bar{x} &gt; Q_2</math>                      So <u>positive skew</u></p> <p><math>\bar{x} + \sigma \approx 18.3</math> so number of plants is e.g. <math>\frac{(25 - "18.3")}{10} \times 12 (+4)</math> (o.e.)                      = 12.04 so <b>12</b> plants</p>	<p>B1 M1 A1    (3)</p> <p>M1 A1    (2)</p> <p>B1 M1A1ft A1    (4)</p> <p>B1ft dB1    (2)</p> <p>M1 A1    (2)</p> <p><b>[13]</b></p>
<b>Notes</b>		
<p><b>(a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p> <p><b>(d)</b></p> <p><b>ALT</b></p> <p><b>(e)</b></p>	<p>M1 for forming a relationship between area and no. of plants or their width <math>\times</math> their height = 6                      A1 for height of 2 (cm). Make sure the 2 refers to height and not plants!</p> <p>M1 for a suitable fraction <math>\times 5</math> (ignore end points)                      A1 for awrt 8.96 (or <math>\frac{215}{24}</math> or <math>8\frac{23}{24}</math>) <u>or</u> 9.06 (or <math>\frac{145}{16}</math> or <math>9\frac{1}{16}</math>) if using <math>(n+1)</math></p> <p>B1 for a correct mean. Accept exact fraction or awrt 10.8                      M1 for a correct expression for <math>\sigma</math> or <math>\sigma^2</math>. Condone mixed up labelling- ft their mean                      A1ft for a correct expression – ft their mean but must have square root                      A1 for awrt 7.46 (use of <math>s =</math> awrt 7.51). Condone correct working and answer called variance.</p> <p>1<sup>st</sup> B1ft for a correct comparison of their <math>\bar{x}</math> and their <math>Q_2</math></p> <p>Allow use of a formula for skewness that involves <math>(\bar{x} - Q_2)</math> or use of quartiles but must have correct values                      NB <math>Q_1 = 5.31</math>, <math>Q_3 = 14.46</math> (awrt 14.5), <math>Q_3 - Q_2 \approx 5.5</math>, <math>Q_2 - Q_1 \approx 3.7/6</math>                      2<sup>nd</sup> dB1 Dependent on a suitable reason for concluding “positive skew”. “correlation” is B0</p> <p>M1 for a suitable expression involving some interpolation (condone missing 4 so accept awrt 8)                      Condone use of end points of 25.5 and 14.5 in their interpolation expressions.                      A1 for 12 (condone awrt 12). Answer only 2/2</p>	

Question	Scheme	Marks
<p>4. (a)</p>	$[P(M < 145) = ] P\left(Z < \frac{145-150}{10}\right)$ $= P(Z < -0.5) \text{ or } P(Z > 0.5)$ $= \text{awrt } \underline{\underline{0.309}}$	<p>M1 A1 A1 (3)</p>
	<b>Notes</b>	
<p>(b)</p>	$[P(B > 115) = 0.15 \Rightarrow] \frac{115-100}{d} = 1.0364$ $\underline{\underline{d = 14.5}}$ <p>(Calc gives 1.036433...) (Calc gives 14.4727...)</p>	<p>M1B1A1 A1 (4)</p>
<p>(c)</p>	$[P(X > \mu + 15   X > \mu - 15) = ] \frac{P(X > \mu + 15)}{P(X > \mu - 15)}$ $= \frac{0.35}{1-0.35}$ $= \underline{\underline{\frac{7}{13}}} \text{ or } \underline{\underline{\text{awrt } 0.538}}$	<p>M1 A1 A1 (3)</p> <p>[10]</p>
	<b>Notes</b>	
	<b>Condone poor use of notation if a correct line appears later.</b>	
<p>(a)</p>	<p>M1 for standardising with 145, 150 and 10. Allow <math>\pm</math> and use of symmetry so 155 instead of 145 1<sup>st</sup> A1 for <math>P(Z &lt; -0.5)</math> or <math>P(Z &gt; 0.5)</math> i.e. a z value of <math>\pm 0.5</math> and a correct region indicated 2<sup>nd</sup> A1 for awrt 0.309 Answer only is 3/3</p>	
<p>(b)</p>	<p>M1 for <math>\pm \frac{115-100}{d} = z</math> where <math> z  &gt; 1</math> Condone MR of <math>\mu = 150</math> instead of 100 for M1B1 only B1 for a standardised expression = <math>\pm 1.0364</math> (do not allow for use of <math>1 - 1.0364</math>) 1<sup>st</sup> A1 for <math>z = \text{awrt } 1.04</math> and compatible signs i.e. a correct equation with <math>z = \text{awrt } 1.04</math> 2<sup>nd</sup> A1 for awrt 14.5 (allow awrt 14.4 if <math>z = \text{awrt } 1.04</math> is seen)</p>	
<p>Calc</p>	<p>Answer only of awrt 14.473 scores M1B1A1A1 Answer only of awrt 14.48 scores M1B0A1A1</p>	
<p>(c)</p>	<p>M1 for a correct ratio expression need <math>P(X &gt; \mu + 15)</math> on numerator. Allow use of a value for <math>\mu</math> May be implied by next line. NB <math>\frac{0.35 \times 0.65}{0.65} = \frac{0.2275}{0.65}</math> is M0</p>	
<p>1<sup>st</sup> A1</p>	<p>for a correct ratio of probabilities</p>	
<p>2<sup>nd</sup> A1</p>	<p>for awrt 0.538 or <math>\frac{7}{13}</math> (o.e.). Allow 0.5385 provided 2<sup>nd</sup> A1 is scored.</p>	



Question	Scheme	Marks
<p><b>6. (a)</b></p> <p><b>(b)</b></p> <p><b>(c)</b></p>	<p><math>[P(B) = 0.4, P(A) = p + 0.1 \text{ so } 0.4 \times (p + 0.1) = 0.1 \text{ or } 0.4 \times P(A) = 0.1</math></p> $p = \frac{1}{4} - 0.1 \quad \quad \quad \underline{p = 0.15}$ <p><math>\frac{5}{11} = \left[ \frac{P(B \cap C)}{P(C)} = \right] \frac{0.2}{0.2 + q} \quad \text{or} \quad \frac{5}{11} = \frac{0.2}{P(C)}</math></p> $11 \times 0.2 = 5 \times (0.2 + q)$ $r = 0.6 - (p + q) \quad \text{i.e. } \underline{r = 0.21} \quad \quad \quad \underline{q = 0.24}$ <p><math>\left[ \frac{P((A \cup C) \cap B)}{P(B)} \right] = \frac{0.3}{0.4}</math></p> $= \underline{0.75}$	<p>M1</p> <p>M1A1 (3)</p> <p>M1</p> <p>dM1</p> <p>A1</p> <p>A1ft (4)</p> <p>M1</p> <p>A1 (2)</p> <p><b>[9]</b></p>
<b>Notes</b>		
	<p><b>(a)</b> 1<sup>st</sup> M1 for using independence in an attempt to form an equation in <math>p</math> or <math>P(A)</math>                  2<sup>nd</sup> M1 for a correct attempt to solve their linear equation leading to <math>p = \dots</math>                  A1 for 0.15 or exact equivalent</p> <p><b>(b)</b> 1<sup>st</sup> M1 for a clear attempt to use <math>P(B/C)</math> to form an equation for <math>q</math> or <math>P(C)</math>. Assuming indep M0                  2<sup>nd</sup> dM1 Dep. on 1<sup>st</sup> M1 for correctly simplifying to a linear equation in <math>q</math> or <math>P(C)</math>                  e.g. accept <math>11 \times 0.2 = 5 \times 0.2 + q</math> or <math>5P(C) = 2.2</math>                  1<sup>st</sup> A1 for <math>q = 0.24</math> or exact equivalent                  2<sup>nd</sup> A1ft for <math>0.6 - \text{their } (p + q)</math> Dependent on 1<sup>st</sup> M1 in (b) only.</p> <p><b>(c)</b> M1 for a correct ratio expression and one correct value (num &lt; denom) <u>or</u> a fully correct ratio.                  Allow <math>\frac{P(A \cup C \cap B)}{P(B)}</math> with one probability correct but only if num &lt; denom.                  A numerator of <math>P(A \cup C) \times P(B)</math> scores M0                  A1 for 0.75 or an exact equivalent</p>	

Question	Scheme	Marks
7. (a)	$E(S) = 0+1 \times 0.2+2 \times 0.1+4 \times 0.3+5 \times 0.2 = [0.2 + 0.2 + 1.2 + 1.0]$ <b>2.6</b>	M1 A1 (2)
(b)	$E(S^2) = 0+1 \times 0.2+2^2 \times 0.1+4^2 \times 0.3+5^2 \times 0.2$ or $0.2 + 0.4 + 4.8 + 5$ <b>10.4 (*)</b>	M1 A1cso (2)
(c)	$\text{Var}(S) = 10.4 - ("2.6")^2$ <b>3.64</b> or $\frac{91}{25}$ (o.e.)	M1 A1 (2)
(d)(i)	$5E(S) - 3 = 5 \times "2.6" - 3,$ = <b>10</b>	M1, A1
(ii)	$5^2 \text{Var}(S) = 25 \times 3.64,$ = <b>91</b>	M1, A1 (4)
(e)	$5S - 3 > S + 3 \Rightarrow 4S > 6$ or $S > 1.5,$ so need $P(S \geq 2)$ $P(S \geq 2) =$ <b>0.6</b>	M1, A1 A1 (3)
(f)	$P(S_1 = 1) \times P(S_2 \leq 4), = 0.2 \times 0.8 = 0.16$ (*)	M1, A1cso(2)
(g)	$P(S_1 = 2) \times P(S_2 \leq 2) = 0.1 \times 0.5 = 0.05$ $P(S_1 = 4) \times P(S_2 \leq 1) = 0.3 \times 0.4 = 0.12$ Full method – all cases listed $P(S_1 = 5) \times P(S_2 = 0) = 0.2 \times 0.2 = 0.04$ all correct products $P(S_1 = 0) \times P(S_2 = \text{any value}) = 0.2 \times 1 = 0.20$ <b>0.57</b>	M1 A1 A1 (3) <b>[18]</b>

Notes		
(a)	M1 for an attempt at $\sum xP(X = x),$ at least 2 non-zero terms seen. Correct answer 2/2 A1 for 2.6 or any exact equivalent	
(b)	M1 for a correct attempt, at least 3 non-zero terms seen A1cso for 10.4 provided M1 is scored and no incorrect working seen	
(c)	M1 for $10.4 - \mu^2,$ ft their $\mu.$ Must see their value of $\mu$ squared (A1 for 3.64 or any exact equiv.)	
(d)(i)	M1 for a correct expression using their 2.6 (A1 for 10)	
(ii)	M1 for $25 \times \text{Var}(S)-$ ft their $\text{Var}(S)$ (A1 for 91)	
(e)	M1 for solving the inequality as far as $pS > q$ where one of $p$ or $q$ are correct 1 <sup>st</sup> A1 for $P(S \geq 2)$ 2 <sup>nd</sup> A1 for 0.6 (provided $S > 1.5$ was obtained). Ans only of 0.6 scores 3/3	
<b>A table showing all 25 cases can only score M1 in (g) if the correct cases are indicated.</b>		
(f)	M1 for using independence (so multiplying) and attempting $P(S_2 \leq 4)$ e.g. $0.2 \times (0.2 + 0.2 + 0.1 + 0.3)$ or $0.04 + 0.04 + 0.02 + 0.06$ score M1 BUT $\frac{4}{25}$ (not from $0.2 \times 0.8$ ) is M0A0 A1cso for a fully correct explanation leading to 0.16. Must come from $0.2 \times 0.8$ not $\frac{4}{25}$	
(g)	M1 for all cases for $S_1$ or all 15 cases for $X$ 1 <sup>st</sup> A1 for all correct probability products for $S_1$ or $X$ 2 <sup>nd</sup> A1 for 0.57 Correct answer scores 3/3. Probabilities out of 25 score A0A0	

Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UA036996 Summer 2013

For more information on Edexcel qualifications, please visit our website  
[www.edexcel.com](http://www.edexcel.com)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual  




Llywodraeth Cynulliad Cymru  
Welsh Assembly Government



Mark Scheme (Results)

Summer 2013

GCE Statistics 1 (6683/01)

## Edexcel and BTEC Qualifications

Edexcel and BTEC qualifications come from Pearson, the world's leading learning company. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information, please visit our website at [www.edexcel.com](http://www.edexcel.com).

Our website subject pages hold useful resources, support material and live feeds from our subject advisors giving you access to a portal of information. If you have any subject specific questions about this specification that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

[www.edexcel.com/contactus](http://www.edexcel.com/contactus)

## **Pearson: helping people progress, everywhere**

Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

Summer 2013

Publications Code UA036993

All the material in this publication is copyright

© Pearson Education Ltd 2013

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.


**EDEXCEL GCE MATHEMATICS****General Instructions for Marking**

1. The total number of marks for the paper is 75.
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M** marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
  - **A** marks: accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B** marks are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.
3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes:

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\surd$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - dep – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper
  - $\square$  The second mark is dependent on gaining the first mark
4. All A marks are 'correct answer only' (cao.), unless shown, for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.
  5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
  6. If a candidate makes more than one attempt at any question:
    - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
    - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
  7. Ignore wrong working or incorrect statements following a correct answer.
  8. In some instances, the mark distributions (e.g. M1, B1 and A1) printed on the candidate's response may differ from the final mark scheme.

Question	Scheme	Marks
<p>1. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p>	$(S_{th}) = 64980 - \frac{7150 \times 110}{9} = -22408.9 \dots$ $(S_{hh}) = 7171500 - \frac{7150^2}{9} = 1491222.2 \dots$ $r = \frac{-22408.9}{\sqrt{1491222 \times 371.56}} = -0.95200068 \dots$ <p>Yes as <math>r</math> is close to <math>-1</math> (if <math>-1 &lt; r &lt; -0.5</math>) <u>or</u> Yes as <math>r</math> is close to <math>1</math> (if <math>1 &gt; r &gt; 0.5</math>)                      [ If <math>-0.5 \leq r \leq 0.5</math> allow "no since <math>r</math> is close to 0" ] [ If <math> r  &gt; 1</math> award B0 ]</p> $b = \frac{-22408.9}{1491222.2} = -0.015027 \dots \quad \left( \text{allow } \frac{-56}{3725} \right)$ $a = \frac{110}{9} - \text{"their } b \text{"} \times \frac{7150}{9} = (12.2 - -0.015 \times 794.4), = 24.1604 \dots \text{ so } t = \mathbf{24.2 - 0.015h}$ <p>0.015 is the <u>drop</u> in temp, (in <math>^{\circ}\text{C}</math>), for every 1(m) <u>increase</u> in height above sea level.</p> $\text{Change} = ("24.2 - 0.015" \times 500) - ("24.2 - 0.015" \times 1000) \text{ or } 500 \times "0.015"$ $= \pm 7.5 \quad (\text{awrt } \pm 7.5) \quad (\text{only ft a value } < 100)$	<p>M1 A1</p> <p>A1</p> <p>(3)</p> <p>M1A1</p> <p>(2)</p> <p>B1ft</p> <p>(1)</p> <p>M1 A1</p> <p>M1, A1</p> <p>(4)</p> <p>B1</p> <p>(1)</p> <p>M1</p> <p>A1ft</p> <p>(2)</p> <p><b>(13 marks)</b></p>
<b>Notes</b>		
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p>	<p>M1 for at least one correct expression (condone transcription error)</p> <p>1<sup>st</sup> A1 for <math>S_{hh} = \text{awrt } 1\,490\,000</math> or <math>S_{th} = \text{awrt } -22\,400</math> (Condone <math>S_{xx}</math> or <math>S_{xy} = \dots</math> or even <math>S_{yy} = \dots</math>)</p> <p>2<sup>nd</sup> A1 for <math>S_{th} = -22\,400</math> <u>and</u> <math>S_{hh} = 1\,490\,000</math> only. [This mark is assessing correct rounding]                      (Allow no labels but mis-labelling <math>S_{th}</math> as <math>S_{hh}</math> etc loses the final A1)</p> <p>M1 for attempt at correct formula. Allow minor transcription errors of 2 or 3 digits.                      Must have their <math>S_{hh}</math>, <math>S_{th}</math> and given <math>S_{tt}</math> (3sf or better) in the correct places. Condone missing “-”</p> <p>Award M1A0 for awrt <math>-0.95</math> with no expression seen. M0 for <math>\frac{64980}{\sqrt{7171500 \times 7.864}}</math></p> <p>B1ft must comment on supporting <b>and</b> state: <u>high/strong/clear</u> (negative or positive) <u>correlation</u>                      “points lie close to a straight line” is B0 since there is no evidence of this.</p> <p>1<sup>st</sup> M1 for a correct expression for <math>b</math>. Follow through their <math>S_{hh}</math> &amp; <math>S_{th}</math>. Condone missing “-”</p> <p>1<sup>st</sup> A1 for awrt <math>-0.015</math> or allow exact fraction from rounded values.</p> <p>2<sup>nd</sup> M1 for a correct method for <math>a</math>. Follow through their value of <math>b</math></p> <p>2<sup>nd</sup> A1 for a correct equation for <math>t</math> and <math>h</math> with <math>a = \text{awrt } 24.2</math> and <math>b = \text{awrt } -0.015</math> No fractions</p> <p>B1 Must mention <math>h</math> (or height) and <math>t</math> (or temperature) and their (1 sf) <u>value</u> of <math>b</math> in a correct comment</p> <p>M1 for a correct expression seen based on their equation. Allow transcription error of 1 digit.                      If answer is <math>500 \times</math> their <math>b</math> to 2sf and <math>&lt; 100</math> (M1A1), If answer is <math>500 \times</math> their <math>b</math> to 2sf and <math>\geq 100</math> (M1A0)</p>	

Question	Scheme	Marks
<p>2. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p>	<p>25 (allow any <math>x</math> where <math>24 &lt; x &lt; 26</math>)</p> <p><math>Q_2</math> (or median or <math>m</math>) = <b>51</b>  <b>IQR</b> = <math>63 - 46</math> ,= <b>17</b> (or <math>Q_3 - Q_1 = 17</math>)</p> <p>Outliers given by <math>46 - 1.5 \times 17 = 20.5</math> <u>or</u> <math>63 + 1.5 \times 17 = 88.5</math>  Outliers limits are <b>20.5</b> and <b>88.5</b></p>  <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p>Allow lower whisker to 20.5 and upper whisker to 88.5  Do <b>not</b> allow a mix of whiskers e.g 20.5 and 85  Do <b>not</b> allow both sets of whiskers</p> </div> <p><b>Medians:</b> Median for females lower than males  <b>IQR:</b> IQR for females smaller than males. Allow “lower/higher” but not “wider”  <b>Range:</b> Range of females is less than males  <b>Skewness:</b> Male and female marks are both positively skew  Ignore other statements about average, spread, mean, st. Dev, variation, outliers etc</p>	<p>B1 <b>(1)</b></p> <p>B1 M1, A1 <b>(3)</b></p> <p>M1 A1</p> <p>M1 A1ft</p> <p>B1 <b>(5)</b></p> <p>B1ft B1ft <b>(2)</b></p> <p><b>(11 marks)</b></p>
<b>Notes</b>		
<p>(b)</p> <p>(c)</p> <p>(d)</p>	<p><b>Mark (b) and (c) together BUT</b> must see clear statement that median (or <math>m</math> or <math>Q_2</math>) = 51 and IQR = 17</p> <p>M1 for 2 quartiles (at least one correct) and attempt to find the difference. Must see their 63 – their 46  A1 for 17 only. [Answer only of IQR= 17 scores M1A1]</p> <p><b>A fully correct box-plot (either version) with no supporting work scores 5/5. Otherwise:</b>  1<sup>st</sup> M1 for correct attempt to calc’ at least one limit for outliers, ft their quartiles or IQR  <u>or</u> award for sight of 20.5 or 88.5  1<sup>st</sup> A1 for identifying both limits of 20.5 and 88.5  2<sup>nd</sup> M1 for a box with an upper and a lower whisker(s) with at least 2 correct values (or correct ft) (condone no median marked) (condone 2 upper or 2 lower whiskers)  2<sup>nd</sup> A1ft for their 20.5 or 26 ,46, 51, 63 and 85 or their 88.5 in appropriate places and readable off their scale. Follow through their 20.5 and their 88.5 only, other values need to be correct  If there are 2 upper or 2 lower whiskers A0  B1 for only 2 outliers appropriately marked at 14 and 90 Do not award if whiskers go beyond these values.  <b>Apply <math>\pm 0.5</math> square accuracy for diagram</b>  A box plot <u>not</u> on the graph paper can only score the 1<sup>st</sup> M1A1</p> <p><b>In (d) ft from their diagrams (if no diagram then use their values)</b>  1<sup>st</sup> B1ft for one correct comment comparing median, IQR , range or skewness  2<sup>nd</sup> B1ft for a second correct comment comparing median, IQR, range or skewness  Do not allow contradictory statements</p>	

Question	Scheme	Marks	
3. (a)	$\frac{35+75}{200} = 0.55$	M1 A1 (2)	
(b)	$\frac{200-2}{200} = 0.99$	M1 A1 (2)	
(c)	$[P(W C)] = \frac{P(W \cap C)}{P(C)} = \frac{30/200}{80/200} = \frac{30}{80} = 0.375$	M1 A1 (2)	
(d)		<p>Allow diagrams with intersections between <math>F</math>, <math>C</math> and <math>H</math> provided these are marked with 0.</p> <p>If their diagram indicates extra empty regions do not treat a blank as 0.</p>	M1 B1 for 9, 1 B1 for 77,33 B1 for 64,16 (4)
(e)	$\frac{1+16+33}{200} = 0.25$	M1 A1 (2)	
<b>(12 marks)</b>			

**Notes**

**Correct answers only score full marks for each part  
If a probability is not in [0, 1] award M0**

- (a) M1 for denominator of 200 and attempt to add  $2 + 8$  or  $35 + 75$  or  $30 + 50$   
A1 for 0.55 or exact equivalent fraction e.g.  $\frac{11}{20}$
- (b) M1 for a fully correct expression ( e.g.  $1 - 0.01$  )  
A1 for 0.99 or an exact equivalent fraction
- (c) M1 for a correct ratio or a correct formula and at least one correct prob (i.e. a correct num or denom). BUT award M0 if num is  $P(W) \times P(C) = \frac{67}{200} \times \frac{80}{200}$  or if num > denom  
A1 for 0.375 or  $3/8$  or any exact equivalent.
- (d) M1 for a box and the 3 regions  $F$ ,  $C$  and  $H$  labelled or implied and single set  $B$  labelled. There should be no intersections between  $F$ ,  $C$  and  $H$  unless marked by zeros. They may have 3 circles for  $F$ ,  $C$  and  $B$  with  $H = F' \cap C'$  etc. Condone lack of zero in the given diagram.
- |                                                                                                                                                                                                                                                                                                                                                    |                                                                 |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------|
| <p><math>F</math> 1<sup>st</sup> B1 for the 9 and 1 <u>or</u> 0.045 and 0.005 (o.e.) in the correct regions</p> <p><math>H</math> 2<sup>nd</sup> B1 for the 77 and 33 <u>or</u> 0.385 and 0.165 (o.e.) in the correct regions</p> <p><math>C</math> 3<sup>rd</sup> B1 for the 64 and 16 <u>or</u> 0.32 and 0.08 (o.e.) in the correct regions.</p> | <p>May have <math>B</math> in 3 bits that are disconnected.</p> |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------|
- (e) M1 for a numerator made up of their  $1 +$  their  $16 +$  their  $33$  and a denom of 200 and num < 200  
Also allow sum of their probabilities (provided sum < 1)  
A1 for 0.25 or any exact equivalent

Question	Scheme	Marks
4. (a)	$\sum ft = 4837.5$ (allow 4838 or 4840) Mean = $\frac{"4837.5"}{200} = 24.1875$ awrt <b>24.2</b> or $\frac{387}{16}$ $\sigma = \sqrt{\frac{134281.25}{200} - \left(\frac{4837.5}{200}\right)^2}$ $= 9.293 \dots\dots$ (accept $s = 9.32$ ) awrt <b>9.29</b>	B1 M1 A1 M1 A1 (5)
(b)	$Q_2 = [20.5] + \frac{(100/100.5 - 62)}{88} \times 5 = 22.659\dots$ awrt <b>22.7</b>	M1 A1 (2)
(c)	$Q_1 = 10.5 + \frac{(50/50.25)}{62} \times 10 [= 18.56]$ (*) ( $n + 1$ gives 18.604...)	B1 cso (1)
(d)	$Q_3 = 25.5$ (Use of $n + 1$ gives 25.734...) IQR = 6.9 (Use of $n + 1$ gives 7.1)	B1 B1 ft (2)
(e)	The data is skewed (condone "negative skew")	B1 (1)
(f)	Mean decreases and st. dev. remains the same. [Must mention mean and st. dev.] (from(a)) The median and quartiles would decrease. [Must refer to median <u>and</u> at least $Q_1$ .] ((b)(c)) The IQR would remain unchanged (from (d))	B1 B1 B1 (3) (14 marks)

**Notes**

<b>Correct answers only score full marks in each part except (c)</b>	
(a)	B1 for 4837.5 or 4838 or 4840 seen. If no $\sum ft$ seen (or attempt at $\sum ft$ seen), B1 can be implied by a correct mean of awrt 24.2 1 <sup>st</sup> M1 for attempt at their $\frac{\sum ft}{\sum f}$ allow 1sf so $\sum f =$ awrt 200 and $\sum ft =$ awrt 5000. Or award M1 for a clear attempt at mean where at least 4 correct products of $\sum ft$ are seen 2 <sup>nd</sup> M1 for correct expression including square root seen. Follow through their mean. Allow a transcription error in 134281.25 but not an incorrect re-calculation.
(b)	M1 for a correct fraction $\times 5$ . Ignore end point but must be +. Allow use of $(n + 1)$ giving 100.5...
(c)	B1cso for a fully correct expression including end point. NB Answer is given. Allow use of $(n + 1)$ giving 50.25...but use of 50.5 scores B0
(d)	1 <sup>st</sup> B1 for 25.5 (or awrt 25.7 using $n + 1$ ) 2 <sup>nd</sup> B1ft for their $Q_3 -$ their $Q_1$ (or 18.6) (provided $> 0$ ) Accept awrt 2sf. Correct ans. only scores 2/2
(e)	B1 Must mention that the data is skewed or not symmetrical. Do not award for "outliers"
(f)	1 <sup>st</sup> B1 for one correct comment from the above. May refer to parts (a), (b), (c) or (d) 2 <sup>nd</sup> B1 for two correct comments from the above 3 <sup>rd</sup> B1 for all 3 correct comments from the above

Question	Scheme	Marks												
5. (a)	$3a + 2b = 0.7$ $a + 2a + 3a + 4b + 5b + 1.8 = 4.2$ or $6a + 9b = 2.4$ $5b = 1$ Attempt to solve $b = \underline{0.2}$ cao      B1 $a = \underline{0.1}$ cao      B1	(5)												
(b)	$E(X^2) = 1 \times 0.1 + 2^2 \times 0.1 + 3^2 \times 0.1 + 4^2 \times 0.2 + 5^2 \times 0.2 + 6^2 \times 0.3 (= 20.4)$ (*)	B1cso (1)												
(c)	$[\text{Var}(X) = ] 20.4 - 4.2^2 [= 2.76]$ $\text{Var}(5 - 3X) = 9 \text{Var}(X)$ $= \underline{24.84}$ or $\underline{24.8}$ (allow $\frac{621}{25}$ )      cao	M1 M1 A1 (3)												
(d)	$[5k = 1 \quad \text{so}] \quad k = \underline{0.2}$	B1 (1)												
(e)	$P(Y = 1) = 0.1$ e.g. $P(Y = 2) = F(2) - F(1) = 0.1$	B1 M1												
	<table border="1" style="display: inline-table; vertical-align: middle;"> <tr> <td>y</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> </tr> <tr> <td>P(Y = y)</td> <td>0.1</td> <td>0.1</td> <td>0.4</td> <td>0.2</td> <td>0.2</td> </tr> </table> Condone use of X(x) instead of Y(y) Ignore incorrect or no label if table fully correct	y	1	2	3	4	5	P(Y = y)	0.1	0.1	0.4	0.2	0.2	A1 (3)
y	1	2	3	4	5									
P(Y = y)	0.1	0.1	0.4	0.2	0.2									
(f)	$P(X = 1) \times P(Y = 1) = \underline{0.01}$	cao      M1, A1 (2)												

**Notes**

**Probabilities outside [0, 1] should be awarded M0**

(a)	1 <sup>st</sup> M1 for an attempt at a linear equation in $a$ and $b$ based on sum of probs. = 1 2 <sup>nd</sup> M1 for an attempt at a second linear equation in $a$ and $b$ based on $E(X) = 4.2$ Allow one slip. 3 <sup>rd</sup> M1 for an attempt to solve their 2 linear equations based on sum of probs and $E(X)$ . Must reduce to a linear equation in one variable. 1 <sup>st</sup> B1 for $b$ and 2 <sup>nd</sup> B1 for $a$ . Answers only score B1B1 only The 3 <sup>rd</sup> M1 may be implied if M2 is scored and both correct answers are given.
<b>ALT</b>	B1B1 for stating $b$ and $a$ . 1 <sup>st</sup> M1 for showing that sum of probs. = 1 2 <sup>nd</sup> M1 for showing that $E(X) = 4.2$ 3 <sup>rd</sup> M1 for an overall comment “(therefore) $a = \dots$ and $b = \dots$ ” No comment loses this mark.
(b)	B1cso for a fully correct expression (no incorrect work seen). E.g. allow $14 \times 0.1 + 41 \times 0.2 + 36 \times 0.3$ Or $0.1 + 0.4 + 0.9 + 3.2 + 5 + 10.8$ . Allow in a table (with 20.4) but without “+” explicitly seen.
(c)	1 <sup>st</sup> M1 for a correct expression for $\text{Var}(X)$ . Must see $-4.2^2$ 2 <sup>nd</sup> M1 for $(-3)^2 \text{Var}(X)$ or better, no need for a value. Accept $-3^2$ if it clearly is used as +9 later.
(e)	B1 for $P(Y = 1) = 0.1$ M1 for correct use of $F(y)$ to find one other prob. Can ft their $k$ if finding $P(Y = y)$ for $y > 2$ Can be implied by one other prob. correct or correct ft Look out for $P(3) = 3k - 0.2$ or $P(4) = P(5) = k$ . A1 for a fully correct probability distribution. Correct table only is 3/3
(f)	M1 for a correct expression or answer ft their $P(Y = 1)$ and their $P(X = 1)$ A1 for 0.01 or exact equivalent only Don't ISW here e.g. $0.1 \times 0.1 + 0.1 \times 0.1$ or $2 \times 0.1 \times 0.1$ are M0A0

Question	Scheme	Marks
<p>6. (a)</p> <p>(b)</p> <p>(c)</p>	<p>[Let <math>X</math> be the amount of beans in a tin. <math>P(X &lt; 200) = 0.1</math>]  <math>\frac{200 - \mu}{7.8} = -1.2816</math> [ calc gives 1.28155156...]  <math>\mu = 209.996\dots</math> awrt 210</p> <p><math>P(X &gt; 225) = P\left(Z &gt; \frac{225 - "210"}{7.8}\right)</math>  <math>= P(Z &gt; 1.92)</math> or <math>1 - P(Z &lt; 1.92)</math> (allow 1.93)  <math>= 1 - 0.9726 = 0.0274</math> (or better) [calc gives 0.0272037...]  <math>= 0.0274</math>  <math>=</math> awrt <b>2.7%</b> allow <b>0.027</b></p> <p>[Let <math>Y</math> be the new amount of beans in a tin]  <math>\frac{210 - 205}{\sigma} = 2.3263</math> or <math>\frac{200 - 205}{\sigma} = -2.3263</math> [ calc gives 2.3263478...]  <math>\sigma = \frac{5}{2.3263}</math>  <math>\sigma = 2.15</math> (2.14933...)</p>	<p>M1 B1 A1 <b>(3)</b></p> <p>M1 A1 A1 <b>(3)</b></p> <p>M1 B1 dM1 A1 <b>(4)</b> <b>(10 marks)</b></p>
<b>Notes</b>		
<p>(a)</p> <p>(b)</p> <p>(c)</p>	<p><b>Condone poor handling of notation if answers are correct but A marks must have correct working.</b></p> <p>M1 for an attempt to standardise (allow <math>\pm</math>) with 200 and 7.8 and set <math>= \pm</math> any <math>z</math> value (<math> z  &gt; 1</math>)  B1 for <math>z = \pm 1.2816</math> (or better used as a <math>z</math>) [May be implied by 209.996(102...) or better seen]  A1 for awrt 210 (can be scored for using 1.28 but then they get M1B0A1)  The 210 must follow from correct working – sign scores A0  If answer is awrt 210 <b>and</b> 209.996... or better seen then award M1B1A1  <math>z = 1.28</math> gives 209.984 and <math>z = 1.282</math> gives 209.9996 and both score M1B0A1  If answer is awrt 210 or awrt 209.996 then award M1B0A1 (unless of course <math>z = 1.2816</math> is seen)</p> <p>M1 for attempting to standardise with 225, their mean and 7.8 . Allow <math>\pm</math>  1<sup>st</sup> A1 for <math>Z &gt;</math> awrt 1.92/3. Allow a diagram but must have 1.92/3 and correct area indicated.  Must have the <math>Z</math> so <math>P(X &gt; 225)</math> with or without a diagram is not sufficient.  Award for <math>1 - 0.9726</math> or <math>1 - 0.9732</math>  2<sup>nd</sup> A1 for 2.7 % or better (calculator gives 2.72...) Allow awrt 0.027. Correct ans scores 3/3</p> <p>1<sup>st</sup> M1 for an attempt to standardise with 200 or 210, 205 and <math>\sigma</math> and set <math>= \pm</math> any <math>z</math> value (<math> z  &gt; 2</math>)  B1 for <math>z = 2.3263</math> (or better) <b>and</b> compatible signs.  If B0 in (a) for using a value in [1.28, 1.29) but not using 1.2816: allow awrt 2.33 here  2<sup>nd</sup> dM1 <b>Dependent on the first M1</b> for correctly rearranging to make <math>\sigma = \dots</math> May be implied  e.g. <math>\frac{5}{\sigma} = 2.32 \rightarrow \sigma = 2.16</math> (M1A0) BUT must have <math>\sigma &gt; 0</math>  A1 for awrt 2.15 . Must follow from correct working but a range of possible <math>z</math> values will do.  NB <math>2.320 &lt; z \leq 2.331</math> will give an answer of awrt 2.15</p>	



Further copies of this publication are available from  
Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467

Fax 01623 450481

Email [publication.orders@edexcel.com](mailto:publication.orders@edexcel.com)

Order Code UA036993 Summer 2013

For more information on Edexcel qualifications, please visit our website  
[www.edexcel.com](http://www.edexcel.com)

Pearson Education Limited. Registered company number 872828  
with its registered office at Edinburgh Gate, Harlow, Essex CM20 2JE

Ofqual



Llywodraeth Cynulliad Cymru  
Welsh Assembly Government



Rewarding Learning