



Mark Scheme (Results)

October 2016

Pearson Edexcel International A Level  
Mathematics

Statistics 1 (WST01)

## Edexcel and BTEC Qualifications

**Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body.** We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

Pearson: helping people progress, everywhere

**Pearson aspires to be the world's leading learning company.** Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all **kinds of people, wherever they are in the world. We've been involved in education for over 150 years**, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

October 2016

Publications Code WST01\_01\_1610\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2016

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero **marks if the candidate's response is not worthy of** credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the **application of the mark scheme to a candidate's response**, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

## PEARSON EDEXCEL IAL MATHEMATICS

### General Instructions for Marking

1. The total number of marks for the paper is 75
2. The Edexcel Mathematics mark schemes use the following types of marks:
  - **M marks: Method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.**
  - **A marks:** Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
  - **B marks** are unconditional accuracy marks (independent of M marks)
  - Marks should not be subdivided.

### 3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod – benefit of doubt
  - ft – follow through
  - the symbol  $\surd$  will be used for correct ft
  - cao – correct answer only
  - cso - correct solution only. There must be no errors in this part of the question to obtain this mark
  - isw – ignore subsequent working
  - awrt – answers which round to
  - SC: special case
  - oe – or equivalent (and appropriate)
  - **d... or dep** – dependent
  - indep – independent
  - dp decimal places
  - sf significant figures
  - \* The answer is printed on the paper or ag- answer given
  - $\square$  or d... **The second mark is dependent on gaining the first mark**
4. **All A marks are 'correct answer only' (cao.), unless shown,** for example, as A1 ft to indicate that previous wrong working is to be followed through. After a misread however, the subsequent A marks affected are treated as A ft, but manifestly absurd answers should never be awarded A marks.

5. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected.
6. If a candidate makes more than one attempt at any question:
  - If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
7. Ignore wrong working or incorrect statements following a correct answer.


## IAL Statistics 1 (WST01) – October 2016

Question Number	Scheme	Marks
1. (a)	[ $P(X > \mu - a)$ ] = <u>0.65</u>	B1 (1)
(b)	[ $P(\mu - a < X < \mu + a)$ ] = $1 - 2 \times 0.35$ or “0.65” – 0.35 or $0.15 + 0.15$ = <u>0.3</u>	M1 A1 (2)
(c)	[ $P(X < \mu + a   X > \mu - a)$ ] = $\frac{\text{"(b)"}}{\text{"(a)"}} = \frac{0.3}{0.65}$  $= \frac{6}{13}$ (Allow awrt 0.462)	M1 A1 (2) <b>[Total 5]</b>
<b>Notes</b>		
(a)	B1 for 0.65 NB you may see $P(Z < 0.35) = 0.6368$ which is of course B0	
(b)	M1 for a correct numerical expression, ft their answer to part (a) [M0 for a probability < 0] A1 for 0.3 (Answer only scores both marks)	
(c)	M1 for a correct ratio of probabilities or follow through their answers provided (b) < (a) A1 for $\frac{6}{13}$ or an exact equivalent and allow awrt 0.462	

Question Number	Scheme	Marks
2. (a)	$2a + 2b + 0.2 = 1$ (o.e.)	B1 (1)
(b)	$[E(X) = ] -2b - a + a + 2b + 3 \times 0.2$ $= \underline{0.6}$	M1 A1 (2)
(c)(i)	$[E(X^2) = 3.5 \Rightarrow ] 3.5 = (-2)^2 \times b + (-1)^2 \times a + a + 2^2 \times b + 3^2 \times \frac{1}{5}$ or $3.5 = 8b + 2a + 1.8$ or $8b + 2a = 1.7$ (o.e.)	M1
(ii)	Solving: $2a + 2b = 0.8$ and $2a + 8b = 1.7$ gives $6b = 0.9$ So $\underline{a = 0.25}$ and $\underline{b = 0.15}$	M1 A1A1 (4)
(d)	$[Var(X) = ] 3.5 - "0.6"{}^2$ $= 3.5 - 0.36 = \underline{3.14}$	M1 A1 (2)
(e)	$P(5 - 3X > 0) = P(5 > 3X) = P(X < 1.66\dots)$ i.e. $P(X = 1 \text{ or } -1 \text{ or } -2)$ $= \underline{0.65}$	M1 A1 A1ft (3)
(f)(i)	$[E(Y) = 5 - 3E(X) = 5 - 1.8] = \underline{3.2}$ (Allow any exact equivalent e.g. $\frac{16}{5}$ )	B1
(ii)	$[Var(Y) = ] (-3)^2 Var(X)$ $= [9 \times 3.14] = \underline{28.26}$ or $\underline{28.3}$ or $\frac{1413}{50}$	M1 A1 (3)
		<b>[Total 15]</b>

**Notes**


(a)	B1 for a correct equation based upon sum of probs. = 1												
(b)	M1 for an attempt at E(X) i.e. an expression with at least 4 correct terms A1 for 0.6 or any exact equivalent. Allow 2/2 for 0.6 only or 0.6 following no incorrect working												
(c)	1 <sup>st</sup> M1 for an attempt to form a second linear equ'n in $a$ and $b$ using $E(X^2)$ , $\geq 3$ correct terms 2 <sup>nd</sup> M1 for an attempt to solve their linear equ'ns...reducing to an equation in one variable. 1 <sup>st</sup> A1 for either $a$ or $b$ correct 2 <sup>nd</sup> A1 for both correct												
(d)	M1 for a correct expression for Var(X) ft their 0.6 A1 for 3.14 or any exact equivalent e.g. $\frac{157}{50}$												
(e)	M1 for attempt to solve the inequality leading to $X < 1.66\dots$ Allow $5 > 3X$ leading to $X < 0.6$ A1 for identifying the 3 correct values of $X$ required A1ft for 0.65 or ft their $2a + b$ ( $a \neq b$ )												
(f)(ii)	M1 for seeing $(-3)^2 Var(X)$ or better [ft their value of Var(X)] A1 for 28.26 or 28.3												
ALT(e,f)	<u>Distribution of Y</u>												
	<table border="1"> <tr> <td><math>y</math></td> <td>11</td> <td>8</td> <td>2</td> <td>-1</td> <td>-4</td> </tr> <tr> <td><math>P(Y=y)</math></td> <td>0.15</td> <td>0.25</td> <td>0.25</td> <td>0.15</td> <td>0.2</td> </tr> </table>	$y$	11	8	2	-1	-4	$P(Y=y)$	0.15	0.25	0.25	0.15	0.2
$y$	11	8	2	-1	-4								
$P(Y=y)$	0.15	0.25	0.25	0.15	0.2								
(f)(ii)	(1 <sup>st</sup> M1 in (e) for at least 3 correct $y$ values) (1 <sup>st</sup> A1 for 11, 8 and 2 with correct ft probs) M1 for correct expression for $E(Y^2) [= 38.5]$ and for $Var(Y) = "38.5" - 3.2^2$												

Question Number	Scheme	Marks
<p>3. (a)</p> <p>(b)(i)</p> <p>(ii)</p> <p>(iii)</p> <p>(c)</p>	<div style="display: flex; justify-content: space-around; align-items: center;">  </div> <p style="text-align: center;">or</p> $P(S) = \left[ \frac{12+23+13}{80} \right] = \frac{48}{80} \text{ or } \frac{3}{5} \text{ or } 0.6$ $P(S C) = \frac{P(S \cap C)}{P(C)} = \frac{\frac{12}{80}}{\frac{20}{80}} = \frac{12}{20} \text{ or } 0.6$ <p><math>P(S) = P(S C)</math> <u>or</u> <math>P(C) = 0.25</math>, <math>P(C \cap S) = 0.15</math> <u>and</u> <math>P(C) \times P(S) = 0.6 \times 0.25</math> so <math>S</math> and <math>C</math> <u>are</u> independent</p> <p>Need <math>P(S G) = \frac{13}{23}</math>  <math>P(S C) = 0.6 &gt; 0.565</math> so assistant selling <u>coats</u> has the better performance</p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1</p> <p>(4)</p> <p>B1ft</p> <p>(1)</p> <p>M1</p> <p>A1cso</p> <p>(2)</p> <p>B1ft</p> <p>dB1ft</p> <p>(2)</p> <p>M1A1</p> <p>A1</p> <p>(3)</p> <p><b>[Total 12]</b></p>
<b>Notes</b>		
<p>(a)</p> <p>(b)(i)</p> <p>(ii)</p> <p>(iii)</p> <p>(c)</p>	<p>1<sup>st</sup> B1 for 3 labelled circles with 12, 13 &amp; <math>n(C \cap G) = 0</math> marked or implied (e.g.RH diagram)</p> <p>2<sup>nd</sup> B1 for 8 and 10 correctly placed</p> <p>3<sup>rd</sup> B1 for 23 correctly placed</p> <p>4<sup>th</sup> B1 for box and 14</p> <div style="border: 1px solid black; padding: 5px; width: fit-content; margin: 10px auto;"> <p>May use probabilities not integers A blank space does not imply a zero</p> </div> <p>B1ft for 0.6 or any exact equivalent (single fraction) <u>or</u> ft their values (ft blank as 0)</p> <p>M1 for a correct conditional prob. Correct expression and one correct ft prob. Num &lt; Den</p> <p>A1cso for 0.6 which must come from a denominator of 20</p> <p>1<sup>st</sup> B1ft for a full reason. If not <math>P(S) = P(S C)</math> then <u>all</u> values must be stated, labelled and correct or correct ft from diagram. Correct not'n required so <math>P(S \cup C) = 0.15</math> is B0B0</p> <p>2<sup>nd</sup> dB1ft dep. on a correct reason for correct conclusion for their values</p> <p>M1 for attempt at <math>P(S G)</math> correct ratio of probabilities or numbers using their figs</p> <p>1<sup>st</sup> A1 for <math>\frac{13}{23}</math> (accept awrt 0.565) [Sight of <math>P(S G) = \frac{13}{23}</math> is M1A1]</p> <p>2<sup>nd</sup> A1 for a correct conclusion that chooses "coats" based on a correct comparison Allow incorrect <math>P(S C)</math> provided <math>&gt; 0.565</math> to score 2<sup>nd</sup> A1 and so all 3 marks Condone poor use of notation eg <math>S G</math> with no <math>P(\dots)</math>. Probabilities may be described in words. Condone comparison of <math>\frac{13}{23}</math> with 0.6 even if <math>\frac{13}{23}</math> is not labelled as <math>P(S G)</math></p>	



Question Number	Scheme	Marks
<p>4. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p>	<p><math>\sum f = 456</math> or <math>\bar{f} = 57</math> and <math>\sum f^2 = 26\ 072</math></p> <p><math>[S_{ff}] = 26072 - \frac{456^2}{8}</math> or <math>26072 - \frac{207936}{8}</math> or <math>26072 - 25992 = 80</math> (*)</p> <p><math>[S_{hh}] = 610\ 463 - \frac{2209^2}{8} = 502.875</math> awrt <b>503</b></p> <p><math>r = \frac{182}{\sqrt{80 \times "502.875"}} = 0.90739\dots</math> = awrt <b>0.907</b></p> <p><math>r</math> is close to 1 or “strong” (positive) correlation (idea of “strong” required) ... so <b>does</b> support the belief</p> <p><math>b = \frac{182}{80} = [2.275]</math>, <math>a = \frac{2209}{8} - b \times \frac{"456"}{8} = [276.125 - 129.675 = 146.45]</math></p> <p>So <b><math>h = 146 + 2.28f</math></b> [Accept <math>h = 146 + 2.27f</math>]</p> <p><math>P(-3 &lt; E &lt; 3)</math> or <math>2P(0 &lt; E &lt; 3)</math>  <math>= \{2\} P\left(0 &lt; Z &lt; \frac{3}{4}\right)</math>  <math>= 2(0.7734 - 0.5) = 0.5468 =</math> awrt <b>0.547</b></p>	<p>M1, A1</p> <p>A1 cso (3)</p> <p>M1, A1</p> <p>(2)</p> <p>M1, A1</p> <p>(2)</p> <p>B1ft</p> <p>(1)</p> <p>M1, M1</p> <p>A1A1</p> <p>(4)</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>(3)</p> <p><b>[Total 15]</b></p>
<b>Notes</b>		
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p>	<p>M1 for an attempt at <math>\sum f</math> and <math>\sum f^2</math> where <math>400 &lt; \sum f &lt; 500</math> and <math>\sum f^2 =</math> awrt 30 000</p> <p>1<sup>st</sup> A1 for <math>\sum f = 456</math> and <math>\sum f^2 = 26\ 072</math> (456 may be implied by 207936 or 25992)</p> <p>2<sup>nd</sup> A1cso for a correct expression leading to 80 with no incorrect working seen.</p> <p>Allow 3/3 for <math>\sum f^2 - 57^2 \times 8</math> or <math>\Sigma(f - 57)^2</math> if all 8 terms seen or <math>5^2 + 4^2 + 1^2 + 2^2 + 3^2 + 5^2</math> if leads to 80</p> <p>M1 for a correct expression for <math>S_{hh}</math></p> <p>A1 for awrt 503 (Answer only scores 2/2 and condone <math>\frac{4023}{8}</math>)</p> <p>M1 for a correct expression using their value for <math>S_{hh}</math> but 182 and 80 must be correct.</p> <p>A1 for awrt 0.907 (Answer only M1A1 awrt 0.91 with no expression M1A0)</p> <p>B1ft for a correct comment that uses their value of <math>r</math> as support e.g. “yes since <u>strong</u> correlation” For <math> r  &lt; 0.5</math> allow comment that does <u>not</u> support.</p> <p>1<sup>st</sup> M1 for a correct expression for <math>b</math> or awrt 2.28 (Allow 2.27 here as well)</p> <p>2<sup>nd</sup> M1 for a correct expression for <math>a</math> ft their value for <math>b</math> and their 456 (using <math>b = r</math> is M0)</p> <p>1<sup>st</sup> A1 for either awrt 146 or awrt 2.28 correctly placed (so M1M0A1A0 is possible)</p> <p>2<sup>nd</sup> A1 for a fully correct regression equation. Accept <math>h - 276 = 2.28(f - 57)</math> [awrt 3 sf] (Allow 2.27 instead of 2.28 for the value of <math>b</math>, since <math>b = 2.275</math>)[Use of <math>y</math> and <math>x</math> is 2<sup>nd</sup> A0]</p> <p>1<sup>st</sup> M1 for a correct probability statement (either of these two expressions)</p> <p>2<sup>nd</sup> M1 for standardising with 3, 0 and 4 (allow <math>\pm 0.75</math> as answer). Ignore <math>\times 2</math> and <math>0 &lt; \dots</math></p> <p>A1 for awrt 0.547 [NB M0M1A0 is common]</p>	

Question Number	Scheme	Marks								
<p>5. (a)</p> <p>(b)</p>	<p><math>[P(A &lt; 388) = 0.001 \Rightarrow] \frac{388 - \mu}{\sigma} = \pm z</math> (where <math> z  &gt; 2.25</math>)</p> <p>So <math>388 - \mu = -3.0902\sigma</math> (o.e.)</p> <p><math>[P(A &gt; 410) = 0.0197 \Rightarrow] \frac{410 - \mu}{\sigma} = \pm z</math> (where <math>1 &lt;  z  &lt; 2.5</math>)</p> <p>So <math>410 - \mu = 2.06\sigma</math> (o.e.)</p> <p>So <math>22 = 5.1502 \sigma</math> (calc gives 5.1502323...)</p> <p><math>\underline{\sigma} = 4.2716...</math> (awrt <b>4.27</b>) and <math>\underline{\mu} = 401.20...</math> (awrt <b>401</b>)</p> <p>[Let <math>X =</math> profit in £ ]</p> <table border="0" style="margin-left: auto; margin-right: auto;"> <tr> <td style="padding-right: 20px;"><math>x</math></td> <td style="padding-right: 20px;">-100</td> <td style="padding-right: 20px;">-0.30</td> <td>0.25</td> </tr> <tr> <td><math>P(X = x)</math></td> <td>0.001</td> <td>0.0197</td> <td><math>1 - (0.001 + 0.0197) = 0.9793</math></td> </tr> </table> <p><math>E(X) = -100 \times 0.001 - 0.30 \times 0.0197 + 0.25 \times 0.9793</math></p> <p style="text-align: right;"><math>= 0.138915...</math> (<b>£</b>) <b>0.14</b></p> <p style="text-align: center;">(allow any answer in the range 0.138...~0.141...)</p>	$x$	-100	-0.30	0.25	$P(X = x)$	0.001	0.0197	$1 - (0.001 + 0.0197) = 0.9793$	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1, A1</p> <p>(7)</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p>(4)</p> <p>[Total 11]</p>
	$x$	-100	-0.30	0.25						
$P(X = x)$	0.001	0.0197	$1 - (0.001 + 0.0197) = 0.9793$							
<b>Notes</b>										
<p>(a)</p> <p>NB</p> <p>(b)</p>	<p>1<sup>st</sup> M1 for attempt to standardise with 388 and set equal to <math>\pm</math> a <math>z</math> value where <math> z  &gt; 2.25</math></p> <p>1<sup>st</sup> A1 for a fully correct equation with <math>z = -3.0902</math> or better (calc gives 3.0902320...)</p> <p>2<sup>nd</sup> M1 for attempt to standardise with 410 and set equal to <math>\pm</math> a <math>z</math> value where <math>1 &lt;  z  &lt; 2.5</math></p> <p>2<sup>nd</sup> A1 for a fully correct equation with <math>z =</math> awrt 2.06 (calc gives 2.059984...)</p> <p>3<sup>rd</sup> M1 for solving their 2 linear equations in <math>\mu</math> and <math>\sigma</math> (i.e. reduce to an equation in one var')</p> <p>Corect processes used but allow 1 sign or numerical slip. Must see the equation in one variable unless the correct answers are obtained in which case this can be implied.</p> <p>3<sup>rd</sup> A1 for <math>\sigma =</math> awrt 4.27 (Allow 4.26 if 1<sup>st</sup> A0 for awrt -3.1)</p> <p>4<sup>th</sup> A1 for <math>\mu =</math> awrt 401</p> <p>Use of <math>\sigma^2</math> instead of <math>\sigma</math> in (a) will score 0/7</p> <p><b>For 3<sup>rd</sup> and 4<sup>th</sup> A marks, apply a 1 mark penalty (A0) the first occasion that an answer in fraction form occurs. So, allow A1 for 2<sup>nd</sup> answer if both fractions give the correct awrt answers.</b></p> <p>1<sup>st</sup> M1 for the 3 correct values of <math>x</math> (i.e. -100, -0.30, +0.25)</p> <p>2<sup>nd</sup> M1 for attempt at all 3 probabilities (correct expression for 0.9793)</p> <p>3<sup>rd</sup> M1 dep on 3 values of <math>X</math> and 3 probs. For an expression for <math>E(X)</math> using their values</p> <p>A1 for an answer of 0.14 or in the range (0.138...~0.141...)</p> <p><u>Not</u> awrt 0.14</p> <p>Accept 0.13 if correct expression is seen beforehand</p>									

Question Number	Scheme	Marks
<p>6. (a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p> <p>(g)</p>	<p><math>Q_1 = 117</math>    <math>Q_2 = 122</math>    <math>Q_3 = 125</math></p> <p><math>IQR = 125 - 117 = \underline{8}</math></p> <p>Upper limit: <math>125 + 1.5 \times 8 = 125 + 12 = \underline{137}</math>    Lower limit: <math>117 - 12 = \underline{105}</math>  Outliers are: <u>101</u>    <u>102</u>    and    <u>139</u></p>  <p><math>[\bar{x} = ] \underline{121}</math></p> <p><math>[\sigma_x = ] \sqrt{\frac{279709}{19} - \bar{x}^2} = \sqrt{14721.526... - 14641} = \sqrt{80.526...}</math> , = awrt <u>8.97</u></p> <p><math>\bar{x} + 2.7 \times \sigma_x = \text{awrt } \underline{145}</math>    <math>\bar{x} - 2.7 \times \sigma_x = \text{awrt } \underline{96.8}</math> (allow 97)</p> <p>Probably not suitable...since: data is skewed <u>or</u> (f) says no outliers, (c) says 3 <u>or</u> (a) says median = 122, (e) says mean = 121</p>	<p>B1B1B1 (3)</p> <p>B1ft (1)</p> <p>M1A1 A1ft (3)</p> <p>M1 A1ft B1 B1 (4)</p> <p>B1 (3)</p> <p>M1A1 (2)</p> <p>B1 (1)</p> <p><b>[Total 17]</b></p>
<b>Notes</b>		
<p>(a)</p> <p>(b)</p> <p>(c)</p> <p>(d)</p> <p>(e)</p> <p>(f)</p>	<p>B1B1B1 for all 3 correct, B1B1B0 for any 2 correct and B1B0B0 for only one correct</p> <p>B1ft for their greatest value from (a) – their smallest value from (a)</p> <p>M1 for a correct expression for either limit, ft their values  1<sup>st</sup> A1 for both 137 and 105 identified  2<sup>nd</sup> A1ft for correctly identifying <u>all</u> the outliers using <u>their</u> limits. Must have scored M1  This 2<sup>nd</sup> A1ft can be awarded if all the outliers are seen on the box plot in (d)</p> <p>M1 for a box and two whiskers (Must be on the grid, otherwise 0/4)  A1ft for <math>Q_1, Q_2</math> and <math>Q_3</math> all plotted correctly, ft their values from (a)  1<sup>st</sup> B1 for lower whisker plotted at 105 or 112 and outliers at 101 and 102  2<sup>nd</sup> B1 for upper whisker plotted at 137 or 132 and an outlier at 139.  (Whiskers must be compatible e.g. if lower whisker is at 105 upper must be at 137)</p> <p>M1 for a correct expression including square root, ft their mean  A1 for awrt 8.97 (Allow <math>s = 9.2195...</math> or awrt 9.22)</p> <p>M1 for an attempt to calculate one limit, ft their mean and s.d.  A1 for both limits correct allow awrt 145 and awrt 96.8 or 97</p>	

- (g) B1 for stating, or implying, normal not suitable **and** giving at least one supporting reason. A calculation or description of skewness is not required but if present must be correct for their values or their box plot. “Not normal since data is not continuous” is B0

Pearson Education Limited. Registered company number 872828  
with its registered office at 80 Strand, London WC2R 0RL