

Mark Scheme (Results) Summer 2009

GCE

GCE Mathematics (6691/01)

June 2009
6691 Statistics S3
Mark Scheme

Question Number	Scheme	Marks
Q1	<p>(a) Randomly select a number between 00 and 499 (001 and 500) select every 500th person</p> <p>(bi) <u>Quota</u> Advantage: <u>Representative</u> sample can be achieved (with small sample size) <u>Cheap</u> (costs kept to a minimum) not “quick“ Administration relatively <u>easy</u> Disadvantage Not possible to estimate sampling errors (due to lack of randomness) Not a random process Judgment of interviewer can affect choice of sample – <u>bias</u> Non-response not recorded Difficulties of defining controls e.g. social class</p> <p>(bii) <u>Systematic</u> Advantage: <u>Simple</u> or <u>easy</u> to use not “quick” or “cheap” or “efficient” It is suitable for large <u>samples</u> (not populations) Disadvantage Only random if the ordered list is (truly) random Requires a list of the population <u>or</u> must assign a number to each member of the pop.</p>	<p>B1 B1 (2)</p> <p>B1</p> <p>B1</p> <p>(2)</p> <p>B1</p> <p>B1 (2)</p> <p>[6]</p>
(a)	<p>1st B1 for idea of using random numbers to select the first from 1 - 500 (o.e.) 2nd B1 for selecting every 500th (name on the list)</p> <p style="text-align: center;">If they are clearly trying to carry out <u>stratified</u> sample then score B0B0</p>	
(b)	Score B1 for any one line	
(i)	<p>1st B1 for Quota advantage 2nd B1 for Quota disadvantage</p>	
(ii)	<p>3rd B1 for Systematic Advantage 4th B1 for Systematic Disadvantage</p>	

Question Number	Scheme	Marks
Q2	<p>(a) Limits are $20.1 \pm 1.96 \times 0.5$</p> <p style="text-align: center;"><u>(19.1, 21.1)</u></p> <p>(b) 98 % confidence limits are</p> $20.1 \pm 2.3263 \times \frac{0.5}{\sqrt{10}}$ <p style="text-align: center;"><u>(19.7, 20.5)</u></p> <p>(c) The growers claim is not correct Since 19.5 does not lie in the interval (19.7, 20.5)</p>	<p>M1 B1 A1cso (3)</p> <p>M1 B1 A1A1 (4)</p> <p>B1 dB1 (2) [9]</p>
	<p>(a) M1 for $20.1 \pm z \times 0.5$. Need 20.1 and 0.5 in correct places with no $\sqrt{10}$ B1 for $z = 1.96$ (or better) A1 for awrt 19.1 <u>and</u> awrt 21.1 but must have scored both M1 and B1 [Correct answer only scores 3/3]</p> <p>(b) M1 for $20.1 \pm z \times \frac{0.5}{\sqrt{10}}$, need to see 20.1, 0.5 and $\sqrt{10}$ in correct places B1 for $z = 2.3263$ (or better) 1st A1 for awrt 19.7 2nd A1 for awrt 20.5 [Correct answer only scores M1B0A1A1]</p> <p>(c) 1st B1 for rejection of the claim. Accept “unlikely” or “not correct” 2nd dB1 Dependent on scoring 1st B1 in this part for rejecting grower’s claim for an argument that supports this. Allow comment on <u>their</u> 98% CI from (b)</p>	

Question Number	Scheme	Marks																																																							
Q3 (a)	<table border="1"> <thead> <tr> <th></th> <th>A</th> <th>B</th> <th>C</th> <th>D</th> <th>E</th> <th>F</th> <th>G</th> <th>H</th> <th>I</th> <th>J</th> </tr> </thead> <tbody> <tr> <td>BMI</td> <td>1</td> <td>6</td> <td>3</td> <td>8</td> <td>4</td> <td>5</td> <td>7</td> <td>2</td> <td>9</td> <td>10</td> </tr> <tr> <td>or</td> <td>10</td> <td>5</td> <td>8</td> <td>3</td> <td>7</td> <td>6</td> <td>4</td> <td>9</td> <td>2</td> <td>1</td> </tr> <tr> <td>Finishing position</td> <td>3</td> <td>5</td> <td>1</td> <td>9</td> <td>6</td> <td>4</td> <td>10</td> <td>2</td> <td>7</td> <td>8</td> </tr> <tr> <td>d^2</td> <td>4</td> <td>1</td> <td>4</td> <td>1</td> <td>4</td> <td>1</td> <td>9</td> <td>0</td> <td>4</td> <td>4</td> </tr> </tbody> </table> <p> $\sum d^2 = 32$ (298) $r_s = 1 - \frac{6 \times 32}{10 \times 99}$ $= 0.80606\dots$ (-0.80606) accept $\pm \frac{133}{165}$ awrt ± 0.806 </p>		A	B	C	D	E	F	G	H	I	J	BMI	1	6	3	8	4	5	7	2	9	10	or	10	5	8	3	7	6	4	9	2	1	Finishing position	3	5	1	9	6	4	10	2	7	8	d^2	4	1	4	1	4	1	9	0	4	4	<p>M1</p> <p>M1</p> <p>M1 A1ft</p> <p>A1 (5)</p>
	A	B	C	D	E	F	G	H	I	J																																															
BMI	1	6	3	8	4	5	7	2	9	10																																															
or	10	5	8	3	7	6	4	9	2	1																																															
Finishing position	3	5	1	9	6	4	10	2	7	8																																															
d^2	4	1	4	1	4	1	9	0	4	4																																															
(b)	<p>$H_0 : \rho = 0, H_1 : \rho > 0,$</p> <p>Critical value is $(\pm)0.5636$</p> <p>$(0.806 > 0.5636$ therefore) in critical region/ reject H_0</p>	<p>B1 B1</p> <p>B1</p> <p>M1 A1ft</p>																																																							
(c)	<p>The lower the BMI the higher the position in the race./ support for doctors belief</p> <p>The position is already ranked OR Position is not Normally distributed</p>	<p>(5)</p> <p>B1 (1)</p> <p>[11]</p>																																																							
(a)	<p>1st M1 for attempt to rank BMI scores</p> <p>2nd M1 for attempt at $\sum d^2$ (<u>must</u> be using ranks)</p> <p>3rd M1 for use of the correct formula with their $\sum d^2$. If answer is not correct an expression is required.</p> <p>1st A1ft for a correct expression. ft their $\sum d^2$ but only if all 3 Ms are scored</p> <p>2nd A1 awrt ± 0.806 (but sign must be compatible with their $\sum d^2$)</p>	<p>No ranking can score 3rd M1 only</p>																																																							
(b)	<p>2nd B1 for $\rho > 0$ (or < 0 but must be one tail and consistent with their ranking)</p> <p>3rd B1 for critical value that is compatible with their H_1. If one-tail must be ± 0.5636 if two-tail must be ± 0.6485 [Condone wrong sign]</p> <p>M1 for a correct statement relating their r_s with their cv. e.g. “reject H_0”, “in critical region”, “significant result” May be implied by a correct comment</p> <p>A1ft for correct comment in context. Must mention low/high BMI and race/fitness <u>or</u> doctor’s belief. Comment should be <u>one</u>-tailed. Allow positive <u>correlation</u> between... but <u>NOT</u> ...positive <u>relationship</u>...</p>	<p>No H_1 assume one-tail for 3rd B1</p>																																																							
(c)	<p>B1 for a correct and relevant comment either based on the fact that the data was originally partially ordered <u>or</u> on the underlying normal assumption “Quicker” or “easier” score B0</p>																																																								

Question Number	Scheme	Marks
Q4	<p>$X \sim N(55, 3^2)$ therefore $\bar{X} \sim N\left(55, \frac{9}{8}\right)$</p> $P(\bar{X} > 57) = P\left(Z > \frac{57 - 55}{\sqrt{\frac{9}{8}}}\right) = P(Z > 1.8856\dots)$ $= 1 - 0.9706$ $= 0.0294$ <p style="text-align: right;"><u>0.0294~0.0297</u></p>	<p>B1 B1</p> <p>M1</p> <p>M1 A1</p> <p style="text-align: right;">[5]</p>
ALT	<p>1st B1 for $\bar{X} \sim$ normal and $\mu = 55$, may be implied but must be \bar{X}</p> <p>2nd B1 for $\text{Var}(\bar{X})$ or st. dev of \bar{X} e.g. $\bar{X} \sim N\left(55, \frac{9}{8}\right)$ or $\bar{X} \sim N\left(55, \left(\frac{3}{\sqrt{8}}\right)^2\right)$ for B1B1</p> <p>Condone use of X if they clearly mean \bar{X} so $X \sim N\left(55, \frac{9}{8}\right)$ is OK for B1B1</p> <p>1st M1 for an attempt to standardize with 57 and mean of 55 and their st. dev. $\neq 3$</p> <p>2nd M1 for 1 - tables value. Must be trying to find a probability < 0.5</p> <p>A1 for answers in the range 0.0294~0.0297</p> $\sum_{i=1}^8 X_i \sim N(8 \times 55, 8 \times 3^2)$ <p>1st B1 for $\sum X \sim$ normal and mean = 8×55</p> <p>2nd B1 for variance = 8×3^2</p> <p>1st M1 for attempt to standardise with 57×8, mean of 55×8 and their st dev $\neq 3$</p>	

Question Number	Scheme	Marks																		
Q5 (a)	$\lambda = \frac{0 \times 40 + 1 \times 33 + 2 \times 14 + 3 \times 8 + 4 \times 5}{100} = 1.05$	M1 A1 (2)																		
(b)	Using Expected frequency = $100 \times P(X = x) = 100 \times \frac{e^{-1.05} 1.05^x}{x!}$ gives $r = 36.743$ awrt 36.743 or 36.744 $s = 19.290$ 19.29 or awrt 19.290	M1 A1 A1 (3)																		
(c)	H_0 : Poisson distribution is a suitable model H_1 : Poisson distribution is not a suitable model	B1																		
	<table border="1"> <thead> <tr> <th>Number of goals</th> <th>Frequency</th> <th>Expected frequency</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>40</td> <td>34.994</td> </tr> <tr> <td>1</td> <td>33</td> <td>36.743</td> </tr> <tr> <td>2</td> <td>14</td> <td>19.290</td> </tr> <tr> <td>3</td> <td>8</td> <td>6.752</td> </tr> <tr> <td>≥ 4</td> <td>5</td> <td>2.221</td> </tr> </tbody> </table>	Number of goals	Frequency	Expected frequency	0	40	34.994	1	33	36.743	2	14	19.290	3	8	6.752	≥ 4	5	2.221	M1
Number of goals	Frequency	Expected frequency																		
0	40	34.994																		
1	33	36.743																		
2	14	19.290																		
3	8	6.752																		
≥ 4	5	2.221																		
	$\nu = 4 - 1 - 1 = 2$ CR : $\chi^2_2(0.05) > 5.991$	B1ft B1																		
	$\sum \frac{(O - E)^2}{E} = \frac{(40 - 34.9937)^2}{34.9937} + \dots + \frac{(13 - 8.972443)^2}{8.972443}$ <p style="text-align: center;">[=0.7161...+0.3813...+1.4508...+1.80789..] = 4.356. (ans in range 4.2 – 4.4)</p>	M1 A1																		
	Not in critical region Number of goals scored can follow a Poisson distribution / managers claim is justified	A1 ft (7)																		
		[12]																		
(a)	M1 for an attempt to find the mean- at least 2 terms on numerator seen Correct answer only will score both marks																			
(b)	M1 for use of correct formula (ft their mean). 1 st A1 for r , 2 nd A1 for s (19.29 OK)																			
(c)	1 st B1 Must have both hypotheses and mention Poisson at least once inclusion of their value for mean in hypotheses is B0 but condone in conclusion 1 st M1 for an attempt to pool ≥ 4 2 nd B1ft for $n - 1 - 1 = 2$ i.e realising that they must subtract 2 from their n 3 rd B1 for 5.991 only 2 nd M1 for an attempt at the test statistic, at least 2 correct expressions/values (to 3sf) 1 st A1 for answers in the range 4.2~4.4 2 nd A1 for correct comment in context based on their test statistic and their cv that mentions goals or manager. Dependent on 2 nd M1 Condone mention of Po(1.05) in conclusion Score A0 for inconsistencies e.g. “significant” followed by “manager’s claim is justified”																			

Question Number	Scheme	Marks
Q6 (a)	<p>$\mu_U \sim$ mean length of upper shore limpets, $\mu_L \sim$ mean length of lower shore limpets</p> <p>$H_0 : \mu_u = \mu_L$</p> <p>$H_1 : \mu_u < \mu_L$</p> <p style="text-align: right;">both</p> $\text{s.e.} = \sqrt{\frac{0.42^2}{120} + \frac{0.67^2}{150}}$ $= 0.0668$ $z = \frac{5.05 - 4.97}{0.0668} = (\pm)1.1975 \quad \text{awrt } \pm \underline{1.20}$ <p>Critical region is $z \geq 1.6449$, or probability = awrt (0.115 or 0.116) $z = \pm 1.6449$</p> <p>(1.1975 < 1.6449) therefore not in critical region / accept H_0/not significant (or $P(Z \geq 1.1975) = 0.1151$, $0.1151 > 0.05$ or z not in critical region)</p> <p>There is no evidence that the limpets on the upper shore are shorter than the limpets on the lower shore.</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>dM1 A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>(8)</p> <p>B1</p> <p>B1</p> <p>(2)</p> <p>[10]</p>
(a)	<p>1st B1 If μ_1, μ_2 used then it must be clear which refers to upper shore. Accept sensible choice of letters such as u and l.</p> <p>1st M1 Condone minor slips e.g. $\frac{0.67^2}{120}$ or $\frac{0.67}{150} + \frac{0.42^2}{120}$ etc i.e. swapped n or one sd and one variance but M0 for $\sqrt{\frac{0.67}{150} + \frac{0.42}{120}}$</p> <p>1st A1 can be scored for a fully correct expression. May be implied by awrt 1.20</p> <p>2nd dM1 is dependent upon the 1st M1 but can fit their se value if this mark is scored.</p> <p>2nd A1 for awrt (+) 1.20</p> <p>3rd M1 for a correct statement based on their z value and their cv. No cv is M0A0 If using probability they must compare their p (<0.5) with 0.05 (o.e) so can allow $0.884 < 0.95$ to score this 3rd M1 mark. May be implied by their contextual statement and M1A0 is possible.</p>	
(b)	<p>3rd A1 for a correct comment to accept null hypothesis that mentions <u>length of limpets</u> on the two <u>shores</u>.</p> <p>1st B1 for one correct statement. Accept "samples are independent"</p> <p>2nd B1 for both statements</p>	

Question Number	Scheme	Marks
Q7 (a)	<p>Estimate of Mean = $\frac{600.9}{5} = 120.18$</p> <p>Estimate of Variance = $\frac{1}{4} \left\{ 72216.31 - \frac{600.9^2}{5} \right\}$ or $\frac{0.148}{4} = 0.037$</p> <p>(b) $P(-0.05 < \mu - \hat{\mu} < 0.05) = 0.90$ or $P(-0.05 < \bar{X} - \mu < 0.05) = 0.90$ [\leq is OK]</p> $\frac{0.05}{\frac{0.2}{\sqrt{n}}} = 1.6449$ $n = \frac{1.6449^2 \times 0.2^2}{0.05^2}$ $n = 43.29\dots$ $n = 44$	<p>M1A1</p> <p>M1 A1ft A1 (5)</p> <p>B1</p> <p>M1 A1</p> <p>dM1</p> <p>A1</p> <p>A1 (6) [11]</p>
(a)	<p>1st M1 for an attempt at $\sum x$ (accept 600 to 1sf)</p> <p>1st A1 for $\frac{600.9}{5} =$ awrt 120 or awrt 120.2. No working give M1A1 for awrt 120.2</p> <p>2nd M1 for the use of a correct formula including a reasonable attempt at $\sum x^2$ (Accept 70 000 to 1sf) or $\sum (x - \bar{x})^2 = 0.15$ (to 2 dp)</p> <p>2nd A1ft for a correct expression with correct $\sum x^2$ but can fit their <u>mean</u> (for expression - no need to check values if it is incorrect)</p> <p>3rd A1 for 0.037 Correct answer with no working scores 3/3 for variance</p> <p>(b) B1 for a correct probability statement <u>or</u> “width of 90% CI = $0.05 \times 2 = 0.1$”</p> <p>1st M1 for $\frac{0.05}{\frac{0.2}{\sqrt{n}}} = z$ value <u>or</u> $2 \times \frac{0.2}{\sqrt{n}} \times z = 0.1$</p> <p>Condone 0.5 instead of 0.05 <u>or</u> missing 2 <u>or</u> 0.05 for 0.1 for M1</p> <p>1st A1 for a correct equation including 1.6449</p> <p>2nd dM1 Dependent upon 1st M1 for rearranging to get $n = \dots$ Must see “squaring”</p> <p>2nd A1 for $n =$ awrt 43.3</p> <p>3rd A1 for rounding up to get $n = 44$</p> <p>Using e.g. 1.645 instead of 1.6449 can score all the marks except the 1st A1</p>	<p>1st B1 may be implied by 1st A1 scored or correct equation.</p>

Question Number	Scheme	Marks
Q8 (a)	$E(4X - 3Y) = 4E(X) - 3E(Y)$ $= 4 \times 30 - 3 \times 20$ $= 60$	M1 A1 (2)
(b)	$\text{Var}(4X - 3Y) = 16 \text{Var}(X) + 9 \text{Var}(Y)$ $= 16 \times 9 + 9 \times 4$ $= 180$	16 or 9; adding M1; M1 A1 (3)
(c)	$E(B) = 80$ $\text{Var}(B) = 16$ $E(B - A) = 20$ $\text{Var}(B - A) = 196$ $P(B - A > 0) = P\left(Z > \frac{-20}{\sqrt{196}}\right) = [P(Z > -1.428\dots)]$ $= 0.923 \dots$	$E(B) - E(A)$ ft on 180 and 16 B1 B1 M1 A1ft stand. using their mean and var dM1 awrt 0.923 - 0.924 A1 (6) [11]
(a)	M1 for correct use of $E(aX + bY)$ formula	
(b)	1 st M1 for $16\text{Var}(X)$ or $9\text{Var}(Y)$ 2 nd M1 for <u>adding</u> variances Key points are the 16, 9 and +. Allow slip e.g using $\text{Var}(X)=4$ etc to score Ms	
(c)	1 st M1 for attempting $B - A$ and $E(B - A)$ or $A - B$ and $E(A - B)$ This mark may be implied by an attempt at a correct probability e.g. $P\left(Z > \frac{0 - (80 - 60)}{\sqrt{180 + 16}}\right)$. To be implied we must see the "0" 1 st A1ft for $\text{Var}(B - A)$ can ft their $\text{Var}(A) = 180$ and their $\text{Var}(B) = 16$ 2 nd dM1 Dependent upon the 1 st M1 in part (c). for attempting a correct probability i.e. $P(B - A > 0)$ or $P(A - B < 0)$ and standardising with their mean and variance. They must standardise properly with the 0 to score this mark 2 nd A1 for awrt 0.923 ~ 0.924	