



Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname

Forename(s)

Candidate signature

A-level PHYSICS

Paper 3

Section B Astrophysics

Thursday 14 June 2018

Morning

Time allowed: The total time for both sections of this paper is 2 hours. You are advised to spend approximately 50 minutes on this section.

Materials

For this paper you must have:

- a pencil and a ruler
- a scientific calculator
- a Data and Formulae Booklet.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each page or on blank pages.
- Do all rough work in this book. Cross through any work you do not want to be marked.
- Show all your working.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 35.
- You are expected to use a scientific calculator where appropriate.
- A Data and Formulae Booklet is provided as a loose insert.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
TOTAL	



J U N 1 8 7 4 0 8 3 B A 0 1

IB/M/Jun18/E9

7408/3BA

Section B

*Do not write
outside the
box*

Answer **all** questions in this section.

0 1

The Griffith Observatory in Los Angeles includes an astronomical refracting telescope (Griffith telescope) with an objective lens of diameter 305 mm and focal length 5.03 m

0 1 . 1

Calculate the wavelength of light for which the Griffith telescope has a minimum angular resolution of 1.8×10^{-6} rad

[2 marks]

wavelength = _____ m

0 1 . 2

The Griffith telescope is used to observe two point objects which subtend an angle of 1.8×10^{-6} rad at the unaided eye.

The typical human eye has a minimum angular resolution of approximately 3.2×10^{-4} rad

Calculate the focal length of the eyepiece lens so that an observer can just resolve the two objects when observing them through the Griffith telescope.

[3 marks]

focal length = _____ m



0 **1** **3**

The asteroid Apophis has a diameter of 325 m

It has been calculated that, in 2029, its distance of closest approach to the Earth's surface will be 3.0×10^4 km

The Griffith telescope may be used to view Apophis using the eyepiece calculated in question **01.2**

Deduce whether this telescope is suitable to obtain a detailed view of Apophis. Support your answer with a calculation.

[3 marks]

8

Turn over ►



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



0 4


0 2 . 1

Sketch, on the axes in **Figure 1**, the black-body radiation curve for a typical star.

[2 marks]

*Do not write
outside the
box*

Figure 1



0 2 . 2

Explain, with reference to the SI units involved, how the curve you have drawn can be used to determine the black-body temperature of the star.

[3 marks]

Question 2 continues on the next page

Turn over ►



0 2 . 3

Two stars, 61 Cygnus A and 61 Cygnus B, can be seen very close together in the constellation Cygnus. Early astronomers were unsure whether the two stars form a binary system, or simply appear in the same line of sight.

Table 1 shows some of the properties of the two stars.

Table 1

	Temperature / K	Radius / km	Apparent magnitude
61 Cygnus A	4500	4.7×10^5	5.2
61 Cygnus B	4100	4.1×10^5	6.1

Evaluate whether the data support the suggestion that the two stars form a binary system.

In your answer you should

- compare the two stars as seen by an observer on Earth
- support your evaluation with suitable calculations.

[6 marks]

*Do not write
outside the
box*



*Do not write
outside the
box*

0 2 . 4

What is the spectral class of 61 Cygnus A?
Tick (✓) the correct box.

[1 mark]

- A
- F
- G
- K

12

Turn over ►



0 3 . 1

Describe the links between galaxies, black holes and quasars.

[2 marks]

0 3 . 2

At a distance of 5.81×10^8 light year, Markarian-231 is the closest known quasar to the Earth. The red shift z of Markarian-231 is 0.0415

Use these data to estimate an age, in seconds, of the Universe.

[4 marks]

age = _____ s



0	3	.	3
---	---	---	---

A typical quasar is believed to be approximately the size of the solar system, with a power output similar to that of a thousand galaxies.

Estimate, with reference to the inverse-square law, how much further the most distant visible quasar is likely to be compared to the most distant visible galaxy.

[3 marks]

*Do not write
outside the
box*

9

Turn over for the next question

Turn over ►



*Do not write
outside the
box*

0	4
---	---

Evidence to support the Big Bang theory comes from cosmological microwave background radiation and the relative abundance of hydrogen and helium in the Universe.

0	4	.	1
---	---	---	---

Explain what is meant by cosmological microwave background radiation and how its existence supports the Big Bang theory.

[3 marks]



0 4 . 2

Explain how the relative abundance of hydrogen and helium supports the Big Bang theory.

[3 marks]

Do not write outside the box

6

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third party copyright material will be published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2018 AQA and its licensors. All rights reserved.

