

PH5

Question	Marking details		Marks Available
SECTION A			
1	(a)	<p>(i) $C = \frac{Q}{V}$ understood (1) [or by impl.] i.e Rearranging to give $V = Q/C$ or substitution of capacitance for C and charge for Q $V = 12.5(3)$ V (1)</p> <p>(ii) $C = \frac{\epsilon_0 A}{d}$ understood [simply quoting is not enough] (1) [substitution of all quantities except d] $d = 9.44 \times 10^{-4}$ m [accept 0.9 mm] (1)</p> <p>(b) $Q = Q_0 \exp\left(\frac{-t}{RC}\right)$ understood (1) [substitution] Taking logs correctly e.g. $\ln Q = \ln Q_0 - \frac{t}{RC}$ (1) Algebra e.g. $-1.9 = \frac{-t}{15 \times 10^6 \times 375 \times 10^{-12}}$ (1) $t = 0.01$ [0.007] s (1) [Use of $\log_{10} \rightarrow 0.47$: treat as calculator slip \rightarrow 3 marks] [Mysterious vanishing of minus sign \rightarrow slip]</p> <p>(c) [Dielectric (or water)] increases C or allows more Q to be stored [accept: store more energy or time constant increased] (1) so change in C or Q means fog or use coulometer to measure Q or use multi(meter) to measure C [or voltage] } (1) NB. 0 marks awarded for answers referring to conduction by water.</p>	<p>2</p> <p>2</p> <p>4</p> <p>2</p> <p>[10]</p>

Question	Marking details		Marks Available
SECTION A			
2	(a)	$B = \frac{\mu_0 I}{2\pi a}$ understood [or $B = 4.8 \times 10^{-7} \text{ T}$] (1) [not $\mu_0 n I$] either $5 \times 4.8 \times 10^{-7}$ or $B = \frac{4\pi \times 10^{-7} \times 1.5}{2\pi \times 0.125}$ (1)	2
	(b)	(i) $\sin \theta = 0^\circ$ or $\theta = 0^\circ$ or $\theta = 180, \pi$ etc (1) Travels along [parallel or opposite to] field lines (1) [NB: 2 nd mark implies first] “to the right” $\rightarrow 0$ “to the right parallel to field” $\rightarrow 1$ bod.	2
		(ii) $F = Bq \sin \theta$ understood (1) [or by impl.], i.e. $\theta = 90^\circ$ calculated [by using $q = 1e$] $\rightarrow 1$ mark $\theta = 30^\circ / 0.52$ radian (1)	2
	(c)	(i) Arrow anti-clockwise ✓	1
		(ii) $Bqv = \frac{mv^2}{r}$ [or $mr\omega^2$] [accept $r = \frac{mv}{Bq}$] (1) $m = 4 \times 1.66 \times 10^{-27} \text{ kg}$ and $q = 2e$ [e.c.f. on q] (1) $r = 76.08 \text{ km}$ (1) Allow ecf on $q = 1e$ i.e. $\rightarrow r = 157 \text{ km}$ [$\rightarrow 2/3$ marks]	3
			[10]

Question	Marking details		Marks Available
SECTION A			
3	(a)	<p>Either Flux changes (1) <u>hence emf induced</u> (1) [Because of RH rule or Faraday → 2nd mark, but not 1st mark] flux increases and decreases [implies 1st mark] [i.e. $\frac{d\Phi}{dt}$ alternates implied](1) NB. “Change in field” not 1st mark but others available]</p> <p>Or B-lines being cut (1) <u>hence emf induced</u> (1) [Because of RH rule or Faraday → 2nd mark, but not 1st mark] direction of cutting changing (1) [Not “magnet oscillating” accept “magnet changing direction [of motion]”]</p>	3
	(b)	<p>(i) $V_{\text{rms}} = \frac{V_0}{\sqrt{2}} = 0.5 \text{ V}$</p> <p>(ii) Rate of change of flux (linkage) = 0.707 [V] (1) from Faraday’s [or Neuman’s] law or $E = N \frac{d\Phi}{dt}$ [allow $E = \frac{\Phi}{t}$](1) [Independent mark – must be stated] For 1 turn = $\frac{0.707}{200} = 0.0035(35) \text{ Wb s}^{-1}$ (1) NB. 0.0025 Wb s^{-1} [from use of $V = 0.5 \text{ V}$] → 2 if 2nd mark awarded.</p>	1
	(c)	<p>Stating or implying that there is a magnetic field set up in the coil (1) Opposes motion / due to Lenz’s law (1) Detail given, e.g. loss (dissipation) of energy due to current or resistance, polarity of coil discussed [can imply 1st mark], work done against resistive force (1)</p>	3
			[10]

Question		Marking details	Marks Available
SECTION A			
4	(a)	<p>γ (1) Needs high penetration (1) [or to irradiate shielded side of metal, or because α and β not penetrating enough etc.] [NB. 2nd mark cannot be given if 1st mark not awarded]</p>	2
	(b)	<p>(i) $\lambda = \frac{\ln 2}{T_{\frac{1}{2}}}$ understood (1) $\lambda = 0.1[308] \text{ year}^{-1} / 4[.14] \times 10^{-9} \text{ s}^{-1}$ ((unit)) [accept Bq] (1) [allow ecf on \log_{10} used $\rightarrow 1.8 \times 10^{-9} \text{ s}^{-1} / 0.057 \text{ year}^{-1}$] [NB per year or per second]</p> <p>(ii) Attempt at using $A = \lambda N$ (1) [allow use of number of moles for N] $1 \text{ mg} = \frac{1}{60} \times 10^{-3} \text{ mol}$ or $N = 10^{19}$ (1) $A = 4.16 \times 10^{10} \text{ Bq}$ [or $1.31 \times 10^{18} \text{ year}^{-1}$] (1) [NB No unit penalty]</p> <p>(iii) $\frac{A}{A_0} = \frac{1}{4}$ (1) Or $2^n = \frac{A_0}{A}$ Or 2 half lives (implies above) (1) $n = 2$ (1) or $A = A_0 e^{-\lambda t}$ (1) $t = 10.6 \text{ year}$ (1) $t = 10.6 \text{ year}$ (1) [used] taking logs (1) $t = 10.6 \text{ year}$ (1)</p> <p>NB $3.3 \times 10^8 \rightarrow 2$ marks, i.e. answer quoted in seconds.</p>	2 3
			3 [10]

Question	Marking details	Marks Available
SECTION A		
5	<p>(a) Conservation of A and Z (1) ${}_{95}^{241}\text{Am} \rightarrow {}_{93}^{237}\text{Np} + {}_2^4\alpha$ (1) ${}_{95}^{241}\text{Am} \rightarrow {}_{96}^{241}\text{Np} + {}_1^0\alpha \rightarrow 1$ mark But not ${}_{95}^{241}\text{Am} \rightarrow {}_{93}^{237}\text{Np} + {}_0^0\alpha$</p> <p>(b) Attempt at LHS – RHS [= 0.00608 but allow slips] (1) Mass in u \times 931 (1) or $E = mc^2$ [with mass in kg] (1) = 5.66 MeV (1) ((unit)) or 9.06×10^{-12} J ((unit))</p> <p>(c) (i) <u>attempt</u> at total mass of p + n (1) [e.g. = $95 m_p + 146 m_n$] – 241.00471 (1) [1.95125] \times931 and \div 241 (1) or $E = mc^2$ and \div 241 = 7.5[378] MeV / nucleon (1) or 1.206×10^{-12} J/nucleon [Slips in total mass can get first 3 marks] NB mixing up number of protons and neutrons \rightarrow 7.27 MeV/nucleon</p> <p>(ii) Plot answer on graph e.c.f. $\pm \frac{1}{2}$ square [7.4 – 7.6 MeV/nucleon and 238-244 for nucleon number]</p>	<p>2</p> <p>3</p> <p>4</p> <p>1</p> <p>10</p>

Question	Marking details	Marks Available
SECTION A		
6.	<p>(a) Insert a voltmeter [V in a circle] on the diagram between front and back faces</p> <p>(b) <u>Electrons</u> feel force due to B-field [or Bqv or FLHR; accept BII] (1) Force towards rear face [accept electrons move to rear face or into the page] (1) <u>Leaving</u> / <u>hence</u> front positive (or shortage of electrons) (1)</p> <p>(c) $E = \frac{V}{d}$ (1) [or by impl.] = $\frac{8.5 \times 10^{-3}}{0.004} = 2.125 \text{ V m}^{-1}$ (1)</p> <p>(d) $Bqv = Eq$ (1) $v = \frac{I}{nAe}$ (rearrange) (1) $E = \frac{BI}{nAe}$ (1) [subst] $n = \frac{BI}{EAe}$ (1) = $5.15 \times 10^{21} \text{ m}^{-3}$ ((unit))</p> <div style="border: 1px dashed black; padding: 5px; display: inline-block; margin-left: 20px;"> <p>or $V_H = \frac{BI}{ntq}$ (1) → $n = 5.15 \times 10^{21} \text{ m}^{-3}$ ((unit)) (1) Max 2/4 for remembering equation</p> </div>	<p style="text-align: center;">1</p> <p style="text-align: center;">3</p> <p style="text-align: center;">2</p> <p style="text-align: center;">4</p> <p style="text-align: center;">10</p>

Question	Marking details	Marks Available
SECTION B		
7	<p>(a) Correct substitution into speed = $\frac{\text{distance}}{\text{time}}$ (1)</p> $\left[t = \frac{8 \times 10^8}{3 \times 10^8} \right] = 2.67 \text{ s (1) [Accept fraction } \frac{8}{3}]$ <p>(b) After travelling both ways extra distance is $\lambda / 2$ (1) Hence destructive interference or antiphase / completely out of phase(1)</p> <p>(c) use of $n\lambda = d \sin \theta$ e.g. $7 \times 640 = 815 \sin \theta$ (1) $d = 1.23 \times 10^{-5} \text{ m}$ (1) [accept $\frac{1}{81500}$] any 2 of $\theta_1 = 2.99$, $\theta_2 = 5.99$, $\theta_3 = 9.00$ (1) Sensible comment, e.g. true, nearly true or wrong [if qualified, e.g. separation increases slightly etc.] [e.c.f.](1) [1st mark required for 3rd mark to be awarded]</p> <p>(d) $N \times \frac{1}{2} mc^2 = \frac{3}{2} nRT$ or $\frac{1}{2} mc^2 = \frac{3}{2} kT$ (1) [or by impl.]</p> <p>Algebra $\overline{c^2} = \frac{3kT}{m}$ (1) [or by impl.]</p> $\sqrt{\overline{c^2}} = \sqrt{\frac{3 \times 1.38 \times 10^{-23} \times 300}{23 \times 1.66 \times 10^{-27}}} = [570.35 \text{ m s}^{-1}] \text{ (1)}$ <p>NB. Mixing up m/M and n/N with correct algebra $\rightarrow 1$.</p> <p>(e) Any 3 \times (1) from</p> <ul style="list-style-type: none"> • 0.97 GHz corresponds to Doppler shift [due to 570 m s^{-1}] / red shift / blue shift ✓ • Sodium atom moving towards laser we get resonant absorption / wavelength [or frequency or energy] is exactly right ✓ • \therefore slowing down is tuned or more probable etc ✓ • If atom moving away there is a shift <u>away from</u> resonance / absorption less probable ✓ <p>[NB “more strongly absorbed”, “Doppler-shifted up 0.97 GHz”, “Match the resonance frequency” are phrases in the passage.]</p>	<p>2</p> <p>2</p> <p>4</p> <p>3</p> <p>3</p>

Question	Marking details		Marks Available
SECTION B			
7	(f)	<p>Photon energy = $\frac{hc}{\lambda}$ or hf and $c = \frac{f}{\lambda}$ (1) [= 3.825×10^{-19} J]</p> <p>No. of photos/sec = power \div photon energy (1.93×10^{10}) (1)</p> <p>Momentum of 1 photon = $h / \lambda = 1.275 \times 10^{-27}$ kg ms⁻¹ (1) [indep. mark]</p> <p>Force = $1.93 \times 10^{10} \times 1.275 \times 10^{-27} \times \sin 30 = 1.23 \times 10^{-17}$ N (1)</p> <p>[Slip with nm / m \rightarrow allow ecf]</p> <p>Alternative Method:</p> <p>Force = $\frac{\text{Power}}{c}$ (1) [or by impl.] = 2.467×10^{-17} N (1)</p> <p>Force upwards (on particle) = Force down on light or reference to F = rate of change of momentum(1)</p> <p>= $2.467 \times 10^{-17} \times \sin 30^\circ = 123 \times 10^{-17}$ N (1)</p>	4
	(g)	<p>Good</p> <ul style="list-style-type: none"> • Lasts long time [accept: sustainable / renewable, lasts 000s years] • No nuclear waste [accept: no harmful waste but not “no waste”] • High concentration of energy e.g. per kilogram • No carbon emissions / use less non-renewables • Abundance of fuel / deuterium [and lithium] [not tritium \rightarrow sif] • Could be profitable soon <p>Bad</p> <ul style="list-style-type: none"> • Tritium from where / needs generation • Does not work yet / huge energy in for little out [needs slightly more than “hasn’t got to breakeven”] • Induced nuclear waste. • Set-up / research costs • Possible military use <p>Any 2 or 3 advantages and/or disadvantage \rightarrow 1</p> <p>4 statements with at least 1 of each (1)</p>	2
			[20]

Question		Marking details	Marks Available
SECTION C			
8	(a)	Laminated (or equivalent) (1) to prevent eddy currents (1) Suitable material for core (1) to avoid magnetising/hysteresis losses (1)	4
	(b) (i)	First mark for diagram with V_L , V_C , V_R perpendicular with V_L , opposite V_C [or impedances] (1) resultant reactive impedance is $\omega L - \frac{1}{\omega C}$ [or $V_{\text{react}} = V_L - V_C$], shown on the diagram(1) Resultant [justified] = $\sqrt{\quad}$ etc.(1) or $V = \sqrt{(V_L - V_C)^2 + V_R^2}$ and $V = \sqrt{\left(I\omega L - \frac{I}{\omega C}\right)^2 + I^2 R^2}$	3
	(ii)	$f = \frac{1}{2\pi} \sqrt{\frac{1}{LC}}$ or $\omega = \sqrt{\frac{1}{LC}}$ or $\omega L = \frac{1}{\omega C}$ (1) Convincing substitution and/or algebra (1)	2
	(iii)	$\left[I = \frac{V}{R} = \right] \frac{12}{280}$ (1) Since all voltage across R or V_L and V_C cancel (or X_L and X_C) (1)	2
	(iv)	Equation used e.g. $Q = \frac{\omega L}{R}$ or $\frac{1}{\omega CR}$ used (1) Answer = 2.97 or (3) (1)	2
	(v)	Attempt at substitution e.g. accept $\sqrt{\left(10.35 \times 64 - \frac{1}{10.35 \times 9.2}\right)^2 + 280^2}$ $Z = 1286 \Omega$ (1) $I = \frac{V}{Z}$ (1) [independent mark]= 9.3 mA (1)	4
(vi)	ωL doubled and $\frac{I}{\omega C}$ halved(1) X_C and X_L switched (1)(cf(v)) $(416 - 1671)^2 = (1671 - 416)^2$ or equivalent -ve number squared. (1) Alternative: $X_C = 1671$ and $X_L = 416$ and $R = 280$ [used or implied](1) $Z = 1286(\Omega)$ - clearly shown (1) 3 rd mark - noticing X_C and X_L swapped.(1)	3	
			[20]

Question		Marking details	Marks Available		
9	(a)	(i)	I. Studied reflected light (from glass plate) (1) Reflection from 2 nd plate depends on orientation (not just angle of inc.) / Light asymmetrical about direction of travel / Reflected light polarised (1)	2	
			II. Developed wave theory mathematically (1) accounted for polarisation by reflection or double refraction or diffraction patterns of various obstacles or why we cannot see around corners (1)	2	
		(ii)	• Requires stiff (or solid) medium (where light travels) (1)	2	
			• which would also support longitudinal waves but not observed or would prevent movement of 'ordinary' objects. (1)		
		(b)	(i)	Magnetic fields – rotating vortices (1)	2
				Electric fields – stress (or strain) in vortex material (1)	1
	(ii)		Density and stiffness	1	
	(iii)		His ether (or equations) predicted $c = \sqrt{\frac{1}{\epsilon_0 \mu_0}}$ (1) Experiment confirmed this (within uncertainties).(1)	2	
	(c)	(i)	Oscillating E and B fields. (1) E and B at right angles to each other and to the propagation direction. (1)	2	
			<i>Principle of Relativity</i> understood (either by statement or implied) (1) Not consistent as laws [of E-M] would have special form in this frame (also implies first mark). (1)	2	
		(ii)	I. 6.39 μs	1	
			II. $\Delta\tau = \Delta t \sqrt{1 - \frac{v^2}{c^2}}$ (1) = 0.625 μs (1) [65.3 $\mu\text{s} \rightarrow 0$ marks]	2	
(iii)		III. 0.706 μs (1) approximately 10% (or 13%) out (1) [or any other correct and relevant remark]	2		
			[20]		

Question		Marking details	Marks Available
10 A	(a)	(i) LCS – largest plastic deformation	1
		(ii) QAS – highest breaking stress	1
	(b)	All are same / similar from initial gradients.	1
	(c)	HCS has greater strength and stiffness (1) Carbon in (crystal) lattice (1) Hinders/opposes/stops dislocation movement (1) Hence more opposition to plastic deformation in HCS (1)	4
	(d)	(i) $\frac{1}{2}mv^2 = \frac{1}{2}Fx$ (1) $\times \frac{1}{4}$ (1) $m = \rho Al$ (1) + convincing algebra (1)	4
(ii) $\varepsilon = 0.002$ (1) $v = \frac{1}{2} \sqrt{\frac{700 \times 10^6 \times 0.002}{8000}} = 6.6 \text{ m s}^{-1}$ [answer] (1)		2	
(iii) Accept either LCS or QAS with sensible reason: e.g. LCS has a higher breaking speed (1) because the area under the graph is greater / ε at breaking is much bigger (1) or QAS has a higher speed (1) because the area under the graph in the elastic region is bigger (1)		2	
B	(a)	2.6 → 2.7 GPa from the graph (1) 8.3 → 8.65 kg (1)	2
	(b)	Thin fibres have fewer surface imperfections (1) Very thin fibres have no surface imperfections (1)	2
	(c)	Thin glass fibres encased in resin / epoxy / plastic material	1
			[20]

Question		Marking details	Marks Available
11	(a)	(i) Same shape, below and longer minimum λ_0 (1) peaks in same place (1)	2
		(ii) Peaks/spikes/line spectrum move .	1
		(iii) $eV = \frac{hc}{\lambda}$ (1) $\lambda = 1.66 \times 10^{-11} \text{ m}$ (1)	2
		(iv) $P = IV = 9375 \text{ W}$ (1) 99.5% heat = $0.995 \times 9375 = 9328 \text{ W}$ (1) Or comment that roughly all 9375W dissipated as heat.	2
	(b)	CT detector(s) rotate (1) about patient / analysis point. Multiple detectors output to computer (1) Series of 2D images obtained or 3D image obtained (1)	3
	(c)	Radio waves [2-100 MHz] (1) Resonate or Same/match frequency of [hydrogen] nuclear rotation [or precession]. (1) Causes them to flip/change (1) [Not just: change spin]	3
	(d)	(i) crystal deforms / vibrates [when alternating p.d. applied]	1
		(ii) $\frac{\Delta\lambda}{\lambda} = \frac{2v}{c}$ (1) $v = 0.9 \text{ m s}^{-1}$ (1) [e.c.f. on missing factor of 2]	2
	(e)	(i) Mention of free radicals (1) [or equivalent, e.g. produces chemicals/ions/atoms which react/are highly reactive]. Damages DNA/cells/molecules (1)	2
		(ii) Absorbed dose = energy (absorbed) per unit mass. Dose equivalent = absorbed dose \times Q[uality] factor.	2
			[20]

Question		Marking details	Marks Available
12	(a)	(i) Power = solar constant \times area [or by impl.] (1) = 1.0686×10^{10} W / 1.0686×10^7 kW / 10.7 GW or equiv (1).	2
		(ii) $P = \sigma AT^4$ understood [accept $5.67 \times 10^{-8} \times A \times 5778$] – i.e. 2 terms identified although missing (1) $P = 4\pi r^2$ quoted (1) $P = 3.85 \times 10^{26}$ W (1) Solar constant = $\frac{3.85 \times 10^{26}}{4\pi \times (1.496 \times 10^{11})^2}$ [=1368 W m ⁻²]	4
	(b)	Hours in one year = 24×365 [.25] [or by impl.] (1) Total units = $1.0686 \times 10^7 \times 24 \times 365 \times 0.4$ [or by impl.] (1) Money = units $\times 0.2 = \text{£}7.5$ billion / 7.5×10^{11} p / $\text{£}7.489 \times 10^9$ (1)	
	(c)	Volume = area \times thickness [or by impl.] (1) Mass = density \times volume [or by impl.] (1) [manip] Mass = 4.95×10^6 kg (1)	
	(d)	$4.95 \times 10^6 \div 2500 = 198$ missions [or by impl.] (1) [ecf from (c)] $\times 350 \times 10^6 = \text{£} 69.3$ bn [or equiv.] (1)	2
	(e)	Heat engines inefficient [or by impl.] (1) Since $1 - \frac{T_1}{T_2} \approx 1 - \frac{300}{400} \approx 0.25$ (1) “which is poor” implies first mark. NB. T_2 in range 373 – 1700 K and T_1 in range 273 – 900 K [$< T_2$]	2
(f)	Reasonable since costs recovered in 9/10 years (1) (ignoring time for 200 shuttle missions) + Any 3 \times (1) good points: <ul style="list-style-type: none"> • Not weather dependant ✓ • Solar power at night ✓ • Less/no atmospheric absorption by microwaves ✓ • Time for 200 shuttle missions ✓ • Shuttle program ended ✓ 	4	
			[20]