

# Mark Scheme Electric Fields Past Paper Questions

## Jan 2002—Jan 2010 (old spec)

- 4(a)(i)** (force) to the right ✓  
 (ii) electrons accelerate or speed increases ✓ (2)

**Q4 Jan 2002**

- (b)(i)** sketch to show path curving upwards in the field  
 (must not become vertical) ✓  
 (ii) horizontal component of velocity is unchanged ✓  
 vertical or upwards acceleration (or force) ✓  
 parabolic path described (or named) ✓ max (3)

(5)

$$\begin{aligned} \mathbf{3(a)(i)} \quad E \left( = \frac{Q}{4\pi\epsilon_0 r^2} \right) &= \frac{29 \times 1.6 \times 10^{-19}}{4\pi \times 8.85 \times 10^{-12} \times (1.15 \times 10^{-10})^2} \quad \checkmark \\ &= 3.15 \times 10^{12} \text{ V m}^{-1} \text{ (or N C}^{-1}\text{)} \quad \checkmark \end{aligned}$$

**Q3 Jan 2003**

$$\begin{aligned} \mathbf{(a)(ii)} \quad V \left( = -\frac{GM}{r} \right) &= (-) \frac{6.67 \times 10^{-11} \times 63 \times 1.66 \times 10^{-27}}{1.15 \times 10^{-10}} \quad \checkmark \\ &= (-) 6.07 \times 10^{-26} \quad \checkmark \quad -\text{sign and J kg}^{-1} \quad \checkmark \end{aligned} \quad (5)$$

- (b)** arrow pointing to the right ✓ (1)  
(6)

**2**

**(a)**

**Q2 Jan 2004**


quantity	SI unit	
(gravitational potential)	$\text{J kg}^{-1}$ or $\text{N m kg}^{-1}$	scalar
(electric field strength)	$\text{N C}^{-1}$ or $\text{V m}^{-1}$	vector
(magnetic flux density)	$\text{T}$ or $\text{Wb m}^{-2}$ or $\text{N A}^{-1} \text{ m}^{-1}$	vector

6 entries correct ✓✓✓  
 4 or 5 entries correct ✓✓  
 2 or 3 entries correct ✓ (3)

$$\begin{aligned} \mathbf{(b)(i)} \quad mg &= EQ \quad \checkmark \\ E \left( = \frac{mg}{Q} = \frac{4.3 \times 10^{-9} \times 9.81}{3.2 \times 10^{-12}} \right) &= 1.32 \times 10^4 (\text{V m}^{-1}) \quad \checkmark \end{aligned}$$

- (ii)** positive ✓ (3)  
(6)

<b>Question 5</b>		
(a) (i)	$E = \frac{V}{d} = \frac{1400}{15 \times 10^{-3}} \checkmark (= 9.3 \times 10^4 \text{ V m}^{-1})$	<b>Q5 Jan 2006</b>
(ii)	$t = \frac{l}{v} = \frac{30 \times 10^{-3}}{3.2 \times 10^7} = 9.38 \times 10^{-10} \text{ s} \checkmark$	5
(iii)	$ma_y = Ee \checkmark$ $a_y = \frac{9.3 \times 10^4 \times 1.60 \times 10^{-19}}{9.11 \times 10^{-31}} \checkmark (= 1.64 \times 10^{16} \text{ m s}^{-2})$ acceleration is upwards [or towards + plate] $\checkmark$	
(b)	$v_y (= a_y t) = 1.64 \times 10^{16} \times 9.38 \times 10^{-10} \checkmark (= 1.54 \times 10^7 \text{ m s}^{-1})$ $v = \sqrt{(1.54 \times 10^7)^2 + (3.2 \times 10^7)^2} = 3.55 \times 10^7 \text{ m s}^{-1} \checkmark$ at $\tan^{-1}\left(\frac{1.54}{3.2}\right) = 26^\circ$ above the horizontal $\checkmark$	3
	<b>Total</b>	8

<b>Question 3</b>		
(a) (i)	force per unit charge $\checkmark$ acting on a positive charge $\checkmark$	<b>Q3 Jun 2006</b>
(ii)	vector $\checkmark$	3
(b) (i)	$F = \frac{Q_1 Q_2}{4\pi\epsilon_0 r^2} = \frac{4.0 \times 10^{-9} \times 8.0 \times 10^{-9}}{4\pi \times 8.85 \times 10^{-12} \times (80 \times 10^{-3})^2} \checkmark$ $= 4.5(0) \times 10^{-5} \text{ N} \checkmark$	
(ii)	(use of $V = \frac{Q}{4\pi\epsilon_0 r}$ gives) $0 = \left(\frac{4.0 \times 10^{-9}}{4\pi\epsilon_0 x}\right) - \left(\frac{8.0 \times 10^{-9}}{4\pi\epsilon_0 (80 \times 10^{-3} - x)}\right)$ or $\frac{4}{x} = \frac{8}{80 - x} \checkmark$ $x = 26.7 \text{ mm} \checkmark$	4
(c)	 <p>correct directions for <math>E_4</math> and <math>E_8</math> <math>\checkmark</math>  <math>E_8</math> approx twice as long as <math>E_4</math> <math>\checkmark</math>  correct direction of resultant <math>R</math> shown <math>\checkmark</math></p>	3
	<b>Total</b>	10

<b>Question 4</b>		
(a) (i)	$E \left( = \frac{Q}{4\pi\epsilon_0 r^2} \right) = \frac{79 \times 1.6 \times 10^{-19}}{4\pi \times 8.85 \times 10^{12} \times (3.0 \times 10^{-14})^2} \checkmark$ <p style="text-align: center;">(gives <math>E = 1.3 \times 10^{20} \text{ V m}^{-1}</math> (<math>1.26 \times 10^{20}</math>))</p>	
(ii)	$F (= EQ') = 1.26 \times 10^{20} \times 2 \times 1.60 \times 10^{-19} \checkmark$ $= 40 \text{ N}$ (40.3) $\checkmark$	5
(iii)	$V \left( = \frac{Q}{4\pi\epsilon_0 r} \right) = \frac{79 \times 1.6 \times 10^{-19}}{4\pi \times 8.85 \times 10^{-12} \times 3.0 \times 10^{-14}} \checkmark$ <p style="text-align: center;">gives <math>V = 3.8 \times 10^6 \text{ V}</math> (or <math>\text{J C}^{-1}</math>) (<math>3.79 \times 10^6</math>) <math>\checkmark</math></p>	
(b) (i)	kinetic energy $\rightarrow$ electric potential energy $\rightarrow$ kinetic energy $\checkmark$	
(ii)	initial kinetic energy = potential energy at point P $\checkmark$ $= (2e)V \checkmark$ $= 2 \times 1.60 \times 10^{-19} \times 3.79 \times 10^6 = 1.21 \times 10^{-12} (\text{J}) \checkmark$	4
	<b>Total</b>	<b>9</b>

<b>Question 3</b>		
(a) (i)	force is perpendicular to initial velocity <b>or</b> acts in opposite direction to direction of electric field $\checkmark$ initial velocity component is maintained $\checkmark$ electron is accelerated in perpendicular direction $\checkmark$ parabolic path $\checkmark$	<b>Q3 Jan 2008</b>
(ii)	force is in opposite direction to initial velocity $\checkmark$ electron decelerated $\checkmark$ direction of motion may eventually be reversed $\checkmark$	max 5

(b) (i)	$E \left( \frac{V}{d} \right) = \frac{110}{44 \times 10^{-3}} = 2500 \text{ V m}^{-1}$ (or $\text{N C}^{-1}$ ) $\checkmark$	
(ii)	$F (= EQ) 2500 \times 1.6 \times 10^{-19} = 4.0 \times 10^{-16} \text{ N} \checkmark$	
(iii)	$E_k \text{ gained} (= E_p \text{ lost}) = eV \checkmark$ $= 1.6 \times 10^{-19} \times 110 = 1.7(6) \times 10^{-17} (\text{J}) \checkmark$ <b>[or</b> $E_k \text{ gained} = Fd \checkmark$ $= 4.0 \times 10^{-16} \times 44 \times 10^{-3} = 1.7(6) \times 10^{-17} (\text{J}) \checkmark$ <b>[or</b> use of $F = ma$ and $v^2 = u^2 + 2as$ gives $a = 4.39 \times 10^{14} (\text{m s}^{-2})$ and $v^2 = 3.86 \times 10^{13} (\text{m}^2 \text{s}^{-2}) \checkmark$ $E_k \text{ gained} = \frac{1}{2} mv^2 = \frac{1}{2} \times 9.11 \times 10^{-31} \times 3.86 \times 10^{13}$ $= 1.7(6) \times 10^{-13} \text{ J} \checkmark$ <b>]</b>	4
	<b>Total</b>	<b>9</b>

<b>Question 4</b>			
(a) (i)	$V = \frac{Q}{4\pi\epsilon_0 r}$ and $E_p = eV \checkmark$ gives $E_p \left( \frac{e^2}{4\pi\epsilon_0 r} \right) = \frac{(1.6 \times 10^{-19})^2}{4\pi \times 8.85 \times 10^{-12} \times 2.0 \times 10^{-15}} \checkmark$ $(= 1.15 \times 10^{-13} \text{ J})$	<b>Q4 Jan 2008</b>	5
(ii)	$2 \times \frac{1}{2} mv^2 = 1.15 \times 10^{-13} \checkmark$ gives $v^2 = \frac{1.15 \times 10^{-13}}{2 \times 1.67 \times 10^{-27}} \checkmark \therefore v = 5.8(7) \times 10^6 \text{ m s}^{-1} \checkmark$		
(b) (i)	$\Delta m = 2 \times (2.01355) - (3.01550 + 1.00728) \checkmark (= 4.32 \times 10^{-3} \text{ u})$ $E = 4.32 \times 10^{-3} \times 931.3 = 4.02 \text{ (MeV)} \checkmark$ $= 4.02 \times 10^6 \times 1.6 \times 10^{-19} = 6.4(4) \times 10^{-13} \text{ J} \checkmark$		
(ii)	energy per unit mass = $\frac{6.44 \times 10^{-13}}{4 \times 1.67 \times 10^{-27}}$ $= 9.6(4) \times 10^{13} (\text{J kg}^{-1}) \checkmark$ [denominator may be $2 \times 2.014 \times 1.66 \times 10^{-27}$ ]	4	
(c)	supply of fuel is almost unlimited (deuterium from sea water) $\checkmark$ fewer waste or radioactivity or environmental problems $\checkmark$ energy released per unit mass is (generally) greater $\checkmark$		<b>max 2</b>
		<b>Total</b>	<b>11</b>

<b>Question 4</b>			
(a) (i)	radial straight lines $\checkmark$ symmetrical in all directions $\checkmark$ directed inwards towards charge $\checkmark$ (marks could be taken from diagram)	<b>Q4 Jun 2009</b>	4
(ii)	line, labelled L, which is a circular arc (or a complete circle) centred on charge $\checkmark$		
(b) (i)	$E \left( = \frac{Q}{4\pi\epsilon_0 r^2} \right) = \frac{0.80 \times 10^{-9}}{4\pi \times 8.85 \times 10^{-12} \times (40 \times 10^{-3})^2} \checkmark$ $= 4.50 \times 10^3 (\text{Vm}^{-1}) \checkmark$		
(ii)	point marked at (40, 4.5) $\checkmark$ curve of decreasing gradient $\checkmark$ correct $E \propto (1/r^2)$ relationship shown by line drawn $\checkmark$	5	
		<b>Total</b>	<b>9</b>

Question 2		
(a)	graph B ✓	1
(b)	for graph A, $V \propto \frac{1}{r}$ ✓ for graph B, $E \propto \frac{1}{r^2}$ ✓ [if candidate correctly quotes equations for V and E only, with no further explanation in words, allow ✓ only] <b>[alternatively allow a fully correct reference to the – and + gradients of graphs A and B respectively in regions R for ✓✓]</b>	2
(c)	$E = \frac{Q}{4\pi\epsilon_0 r^2}$ and $V = \frac{Q}{4\pi\epsilon_0 r}$ give $E = \frac{V}{r}$ ✓ [no credit for using just $E = \frac{V}{d}$ ] $\therefore$ potential of the point $V = Er = 3.6 \times 10^4 \times 40 \times 10^{-3} = 1.4 \times 10^3 V$ ✓ (1440) (allow $J C^{-1}$ )	2
	<b>Total</b>	<b>5</b>