



Question number	Answer	Notes	Marks
1 (a)	<p>MP1. pitch is <u>frequency</u>;</p> <p>MP2. any one of:</p> <ul style="list-style-type: none"> • whether sound/note sounds high or low; • high sound has high frequency ORA; 	<p>allow 'it' for pitch</p> <p>ignore references to amplitude, wavelength</p> <p>allow vibrates more often / with shorter time period</p> <p>'high pitch has high frequency' ORA gains 2 marks</p>	2
(b) (i)	ruler / measuring tape; oscilloscope / mobile phone app / data logger / (guitar) tuner;	ignore microphone frequency meter frequency gauge frequency counter	2
(ii)	dependent – frequency / pitch; independent – length (of pipe);		2
(c)	<p>any three of:</p> <p>MP1. repeat AND average the readings;</p> <p>MP2. (measure a) larger range of values;</p> <p>MP3. (measure some) intermediate values;</p> <p>MP4. improved precision of a named variable / instrument;</p> <p>MP5. control a named variable (e.g. temperature);</p> <p>MP6. plot a graph of frequency and length;</p> <p>MP7. deal with anomalies;</p>	<p>accept 'measure more values' for 1 mark if NEITHER MP2 nor MP3 awarded</p> <p>e.g. 'use a cm ruler', 'measure frequency in mHz' etc.</p> <p>ignore references to accuracy</p> <p>allow 'blow with controlled apparatus'</p> <p>allow 'plot a graph of the results'</p> <p>allow 'identify anomalies'</p>	3

Total 9 marks

Question number	Answer	Notes	Marks												
2 a (i)	0.28 0.37	(both for 1 mark)	1												
	(ii) suitable scales; axes labelled; plotting of second and fifth points ;; line of best fit; 	Must use > half width and half height of grid no units on axis labels ignore orientation of graph to nearest ½ square, up to two marks available for this line – allow ecf from candidate’s third and fourth points <table border="1" data-bbox="1268 482 1667 705"> <thead> <tr> <th>sin i</th> <th>sin r</th> </tr> </thead> <tbody> <tr> <td>0.00</td> <td>0.00</td> </tr> <tr> <td>0.26</td> <td>0.17</td> </tr> <tr> <td>0.42</td> <td>0.28</td> </tr> <tr> <td>0.57</td> <td>0.37</td> </tr> <tr> <td>0.71</td> <td>0.47</td> </tr> </tbody> </table> If incorrect graph plotted (i° against r°) the only scales and line mark can be awarded (NB in this case can only get first MP in (a)(iii))	sin i	sin r	0.00	0.00	0.26	0.17	0.42	0.28	0.57	0.37	0.71	0.47	Max 5
sin i	sin r														
0.00	0.00														
0.26	0.17														
0.42	0.28														
0.57	0.37														
0.71	0.47														
	(iii) Attempt at gradient of line, seen on graph or in working; Value in range 1.48 to 1.54;	e.g. triangle or equivalent drawn on graph, rise/run bald correct answer is 1 mark only	2												

b	<p>Any two of -</p> <p>MP1. Idea that value relates to all the data collected;</p> <p>MP2. Idea that method allows for anomalies;</p> <p>MP3. Idea that effects of uncertainty/error can be reduced or accounted for;</p>	<p>Method checks reliability, anomalies can be seen</p> <p>graph is an averaging technique</p> <p>Ignore comments about accuracy</p>	2
---	---	--	---

(Total for Question 2 = 10 marks)

Question number	Answer	Accept	Reject	Marks
3 (a)	<p>Refraction into glass towards the normal ($r > 0$);</p> <p>Angle of incidence <u>and</u> angle of refraction both labelled correctly at the same surface;</p> <p>Refraction at the lower surface into air away from the normal;</p> <p>Emergent ray parallel to incident ray after correct refraction (by eye);</p> 	<p>Accept dotted lines Ignore any reflections</p> <p>Ignore a second incorrectly labelled pair</p>		4

Question number	Answer	Accept	Reject	Marks								
3 (b) (i)	One mark for either sin i or sin r correct; <table border="1" data-bbox="422 294 1037 534" style="margin-left: auto; margin-right: auto;"> <tr> <td style="text-align: center;"><i>i</i></td> <td style="text-align: center;">60°</td> </tr> <tr> <td style="text-align: center;"><i>r</i></td> <td style="text-align: center;">34°</td> </tr> <tr> <td style="text-align: center;">sin <i>i</i></td> <td style="text-align: center;">0.87</td> </tr> <tr> <td style="text-align: center;">sin <i>r</i></td> <td style="text-align: center;">0.56</td> </tr> </table>	<i>i</i>	60°	<i>r</i>	34°	sin <i>i</i>	0.87	sin <i>r</i>	0.56	sin i = 0.866; sin i = 0.8660; sin r = 0.559; sin r = 0.5592; Ignore degree sign Ignore any other values		1
<i>i</i>	60°											
<i>r</i>	34°											
sin <i>i</i>	0.87											
sin <i>r</i>	0.56											
(ii)	n = sin i ÷ sin r;	Accept refractive index = sin i ÷ sin r		1								
(iii)	Two marks for correct answer Refractive index = 1.55;; Or Refractive index = 1.6;; Or Refractive index = 1.5;;	Accept for one mark only any other value in the range 1.5 < n < 1.6; Any power of 10 error, e.g. 155.3		2								

Question number	Answer	Accept	Reject	Marks
3 (c)	<p>Any three of:</p> <p>MP1 any mention of repetition / take an average of readings;</p> <p>MP2 vary i to obtain more values ;</p> <p>MP3 plot a graph of <u>$\sin i$ against $\sin r$</u> ; OR Calculate/work out/ find n;</p> <p>MP4 find gradient of graph ; OR Calculate average of n;</p> <p>MP5 sensible experimental precaution / improvement to method (e.g. mark lines on paper, thinner beam, fix block firmly in position, remove anomalies, sharper pencil, use a more precise protractor e.g. $\frac{1}{2}^\circ$);</p>	<p>Ignore reference to critical angle</p> <p>Ignore second glass block</p> <p>Ignore different colours</p>		3



Total 11 marks

Question number	Answer	Notes	Marks
4 (a)	(i) set-up showing any two from-clear indication of equipment needed; correct refraction at one surface of glass block shown; protractor shown in use;	ray-box or pins Allow ruler for apparent depth method	2
	(ii) angle of incidence; angle of refraction; OR critical angle; idea of grazing emergence;	Allow apparent depth method, i.e. real depth; apparent depth;	2
	(iii) find $\sin i$ and $\sin r$; refractive index is the ratio of sines; OR find $\sin c$; refractive index is $1/\sin c$;	Accept for two marks <ul style="list-style-type: none"> • $(n =) \sin i/\sin r$ • $(n =) 1/\sin c$ • graph of $\sin i$ vs $\sin r$ Allow refractive index = real depth \div apparent depth for two marks	2
	(b) (i) Diagram – reflection at first back surface; reflection at second back surface;	judge by eye <ul style="list-style-type: none"> • straightness of ray and correctness of angle • emergent ray parallel to incident ray 	2
	(ii) Refracted / slows down / wavelength decreases	Ignore: direction change ideas it does nothing / nothing happens	1

Total 9 marks


Question number	Answer	Notes	Marks
5 (a)	<p>cooking – micro(waves) OR infrared (waves);</p> <p>treating cancer – ultraviolet OR x-rays OR gamma (rays);</p> <p>identifying broken bones - x-rays;</p>	if more than one example given for each use then reject mark if any incorrect	3
(b)	C - the same speed;		1
(c) (i)	<p>drawn ray shows refraction in the correct direction (downwards) at both surfaces;</p> <p>drawn ray is above yellow ray and diverges from it (if ray had entered at the original point);</p>	<p>judge by eye ignore arrows and labels dependent on previous</p> <p>allow if ray drawn enters parallel to original ray</p>	2
(ii)	A- black;		1

Total 7 marks


Question number	Answer	Notes	Marks												
6 (a)	B;		1												
(b) (i)	<p>MP1. Axes labelled with units; MP2. Correct scales (to occupy at least ¼ of the area of the graph and in sensible intervals); MP3. Plotting; MP4. Plotting; MP5. straight line of best fit which extends beyond given data points;</p> <div style="display: flex; align-items: center;">  <table border="1" data-bbox="949 859 1192 1069" style="margin-left: 20px;"> <thead> <tr> <th>Distance in m</th> <th>Time in ms</th> </tr> </thead> <tbody> <tr> <td>0.62</td> <td>1.8</td> </tr> <tr> <td>0.80</td> <td>2.4</td> </tr> <tr> <td>1.00</td> <td>3.0</td> </tr> <tr> <td>1.20</td> <td>3.8</td> </tr> <tr> <td>1.38</td> <td>4.2</td> </tr> </tbody> </table> </div>	Distance in m	Time in ms	0.62	1.8	0.80	2.4	1.00	3.0	1.20	3.8	1.38	4.2	<ul style="list-style-type: none"> ignore orientation of graph scale intervals on axes should be 2 or 5 or 10 points should be less than 0.5 sq in diameter -1 each incorrect plot to max of -2 tolerance = +/- ½ square if zero is not included, then line should go through all points except 3rd or 4th if zero included, look for balance of points 	5
Distance in m	Time in ms														
0.62	1.8														
0.80	2.4														
1.00	3.0														
1.20	3.8														
1.38	4.2														

(ii)	<p>Attempt to find slope or gradient of line ; AND evaluation of value; matching unit; e.g. = 0.6/0.0018 = 333 m/s</p>	<p>Δ seen or two lines from same axis seen or rise/run seen value in range of 310-350 allow 0.333 km/s 0.333 m/ms</p>	3
(iii)	<p>Any one specific variable from the experiment; e.g. hitting the block in the same place</p> <p>Use the same microphone/timer/wires</p> <p>Ensure there is no 'hammer bounce'</p>	<p>These must be specific to the experiment Accept same</p> <ul style="list-style-type: none"> • temperature • humidity • density • draughts • force • block <p>ignore</p> <ul style="list-style-type: none"> • 'keep everything the same' • use control variables • repeat experiment 	1
(iv)	<p>Any 2 suggestions from</p> <p>MP1. repeat the time readings (for each distance);</p> <p>MP2. measure the distance to the sensor of the microphone;</p> <p>MP3. use wider range of distance readings (<0.62 or >1.38);</p> <p>MP4. use intermediate distances (between points);</p>	<p>ignore imprecise suggestions e.g.</p> <ul style="list-style-type: none"> • 'be careful with timer' • 'change the distance' 	2

(Total for Question 6 = 12 marks)

Question number	Answer	Notes	Marks
7 (a)	standard definition of wavelength; e. <ul style="list-style-type: none"> • distance between two points on a wave/ two peaks/ two troughs • distance between each wavefront • distance travelled by wave in one time period 	allow: from clear diagram crest for peak ignore: <ul style="list-style-type: none"> • 'the length of a wave' • 'distance taken for 1 cycle' • distance between one wave and the next one 	1
7 (bi)	Speed of wave = frequency x wavelength;	allow: in any rearrangement $v = f \cdot \lambda$	1
(bii)	substitution into any form of the equation ; evaluation; e. $3(\text{m/s}) = 1.5(\text{Hz}) \times \lambda$ $(\lambda) = 2(\text{m});$	accept for 1 mark $\frac{3}{1.5}$	2

Question number	Answer	Notes	Marks
7 (ci)	Diffraction; And one of <ul style="list-style-type: none"> • The incoming wave spreads out at the gap; • The energy carried by the wave spreads out ; 	allow: <ul style="list-style-type: none"> • diffraction seen in (cii) • recognisable spelling for 'diffraction' ignore: <ul style="list-style-type: none"> • the wave gets bigger • wave is bent • (wavefront is) curved 	2
7 (cii)	idea that (diffraction only apparent when) λ and size of gap comparable/RA; wavelength of light is very small / smaller than water waves /smaller than the gap;	Allow RA	2
Total			9

Question number	Answer	Notes	Marks
8 (a)	idea that higher frequency gives higher pitch;	allow reverse argument condone idea of proportionality / linearity	1
(b) (i)	(wave) speed = frequency \times wavelength	allow abbreviation, e. $v = f \times \lambda$ or rearrangements	1
(ii)	substitution into correctly rearranged equation; evaluation; e. (v =) 340 / 160 (v =) 2.1 (m)	allow 2.125, 2.12, 2.13 or 2 (if supported)	2
(c) (i)	straight line of best fit drawn within indicated area; 	line does not need to be extended beyond data range for this mark	1
(ii)	line of best fit extended to 20°C; student's own value from graph \pm half a square;		2

(d)	any 2 from: MP1. speed (of sound) decreases (with temperature); MP2. frequency is constant; MP3. so wavelength decreases (with temperature);	allow 'sound slows down' ignore references to particle speed allow λ is smaller	2
-----	---	---	---

Total 9 marks